

**COVARIATES USED IN THE CASE MIX ADJUSTMENT  
OF PCS and MCS MEASURES**

	PCS AND MCS MODELS		
	One	Two	Three
<b>DEMOGRAPHICS COVARIATES</b>			
Age (Continuous)	✓	✓	✓
Gender (Male or Female)	✓	✓	✓
Race (White, Black, Other Minority)	✓	✓	✓
Education	✓	✓	
Marital Status	✓	✓	
Annual Household Income	✓		
<b>CHRONIC MEDICAL CONDITIONS</b>			
Hypertension or high blood pressure	✓	✓	
Angina pectoris or coronary artery disease	✓	✓	
Congestive heart failure	✓	✓	
Myocardial infarction or heart attack	✓	✓	
Other heart conditions, such as problems with heart valves or arrhythmias	✓	✓	
Stroke	✓	✓	
Emphysema, or asthma, or COPD (Chronic Obstructive Pulmonary Disease)	✓	✓	
Crohn's disease, ulcerative colitis, or inflammatory bowel disease	✓	✓	
Arthritis of the hip or knee	✓	✓	
Arthritis of the hand or wrist	✓	✓	
Sciatica	✓	✓	
Diabetes, high blood sugar, or sugar in the urine	✓	✓	
Any cancer (other than skin cancer)	✓	✓	
<b>HOS STUDY DESIGN VARIABLES</b>			
Who Completed Survey (Self or Other)	✓	✓	
Mode of Survey Administration (Mail or Telephone)	✓	✓	✓
CMS Plan Region	✓	✓	✓
Survey Vendor	✓	✓	✓

Note: Model One included all covariates listed in and was used for beneficiaries with completed data for all of the covariates. Model Two was used for beneficiaries with completed data for all of the covariates except annual household income. Model Three was limited to age, gender, race, mode of survey administration, CMS plan region, and survey vendor, and was used for beneficiaries who did not have enough completed data for Model One or Model Two. The variables included in Model Three were available for all participating beneficiaries.