

Part B. Collection of Information Employing Statistical Methods

B-1. *Description of universe and selection methods used.*

The Unemployment Insurance Data Validation (UI DV) methodology relies on two basic tests for the validity of the aggregate counts that are reported to the Department. The first is an independent reconstruction of the counts, based on a new extract and count from the underlying State database from which the reports were initially prepared. This step tests whether the State prepared its report extracts from the correct sources and whether its item-counting software works properly. In this step the validator also checks--and adjusts--for any duplicate counts that may have inflated the reported counts being validated. The second step checks for invalid elements in the transaction pool from which the elements are extracted for both the reported and validated (reconstructed) counts. If the State labels or classifies certain transactions incorrectly, both reported counts and reconstructed counts will be based on an unknown proportion of individual transactions which do not conform to Federal reporting definitions, and thus both sets of counts will be wrong. The UI DV methodology checks for invalid elements by specifying that samples be drawn from certain classes of transactions and the sampled transactions checked against original UI program documentation available in the State's system to determine whether the State is coding and classifying transactions appropriately.

UI DV relies on existing records from State UI databases and management information systems. As a result, traditional response rate issues do not arise in UI DV. However, states may not complete UI DV or submit reports timely for any of several reasons. See B-3, below.

Because UI DV's scope is very extensive, different sample designs are used for efficiency, to reduce the need for large samples required to estimate a specific proportion of incorrect transactions in the population. The sample types and their logic are as follows. Table B-1 gives the range of samples drawn for Benefits validation. Tax validation relies on an elaborate series of logic tests in building the extract file, supplemented by sorts and two-case samples to ensure that the extract file is built properly. All logic tests, sorts and samples for an extract file must be passed before the reconstructed count can be considered the valid standard for judging reported counts and thus that the reported counts can pass validation.

- **Random Samples.** In Benefits validation, the State draws 17 random samples for the most important types of reports data, e.g., those used to determine administrative funding or build key performance measures. Although random samples of 100 or 200 elements are drawn, only 30 or 60 elements are evaluated initially as acceptance samples; only if the result of the initial acceptance sample is inconclusive is the entire sample evaluated to estimate the underlying error rate.
- **Supplemental Samples for Missing Subpopulations.** These are samples of one transaction from any subpopulations not represented in the random samples of the broader populations which conceptually include them. These are reviewed simply to check that validation files are programmed properly by determining that the only reason the examined sample did not include a representative from the missing subpopulation is sampling variability--probability that rare elements in the population will not be included in the relatively small random samples.
- **Supplemental Samples to Examine Data Outliers.** The random and supplemental samples ensure that the population as a whole was defined properly but probably do not assess whether time-lapse measures or dollar transactions contain extreme values. UI DV addresses this issue by

sorting those populations and examining the five highest and five lowest values in each sorted population to ensure that there are no calculation and data errors. Although in the validation these are referred to as “samples” they are technically the selection of specific elements.

- **Supplemental Minimum Samples.** UI DV draws no random samples for some transactions considered of lower priority. UI DV simply ensures that the reporting software uses the correct field in the database to process and report the transactions. This is done by randomly selecting two cases per subpopulation.

TABLE B-1

Benefits Samples by Population, Type, and Size						
Benefits Population		Sample Name	Universe (subpops)	Sample Type	Size	
Number	Type of Transaction				How Selected	Total
1	Weeks Claimed	Intrastate Weeks Claimed	1.1-1.3	Random	60/200	60/200
		Interstate Liable Weeks Claimed	1.4-1.6	Random	30/100	30/100
		Inter Weeks Claimed filed fr Agent	1.7-1.9	Minimum	2 per subpop	6
2	Final Payments	Final Payments	2.1-2.4	Random	30/100	30/100
3	Claims	New Intra & Inter Liable Claims	3.1-3.18	Random	60/200	60/200
		New Intra & Inter Liable Claims	3.1-3.18	Missing Subpops	1 per subpop	≤17
		Interstate Filed from Agent	3.19-3.21	Minimum	2 per subpop	6
		Interstate Taken as Agent	3.22 -3.24	Minimum	2 per subpop	6
		Intra and Inter Transitional Claims	3.25-3.33	Random	30/100	30/100
		CWC Claims	3.34-3.39	Random	30/100	30/100
		CWC Claims	3.34-3.39	Missing Subpops	1 per subpop	≤5
		Monetary Sent w/o New Claim	3.40 3.45	Minimum	2 per subpop	12
		Entering Self Employment Pgm	3.46	Minimum	2	2
3a	Additional Claims	Intrastate Additional Claims	3a.1-3a.3	Random	30/100	30/100
		Interstate Liable Additional Claims	3a.4-3a.6	Minimum	2 per subpop	6
4	Payments	First Payments	4.1-4.16	Random		60/200
		First Payments	4.1-4.16	Missing Subpops	1 per subpop	≤15
		First Payments: Intrastate Outliers	4.1, 4.3, 4.5, 4.7, 4.9, 4.11, 4.13, 4.15	Outliers (TL)	5 highest, 5 lowest	10
		Continued Weeks total Payments	4.17-4.24	Outliers (TL)	5 highest, 5 lowest	10
		Continued Weeks Partial Payments	4.24-4.32	Random	30/100	30/100
		Adjusted Payments	4.33-4.42	Outliers (\$)	5 highest, 5 lowest	10
		Self-Employment Payments	4.43	Minimum	2	2
		CWC First Payments	4.44-4.45	Random	30/100	30/100
		CWC Continued Payments	4.46-4.47	Minimum	2 per subpop	4
		CWC Adjusted Payments	4.48-4.49	Minimum	2 per subpop	4
		CWC Prior Weeks Compensated	4.50-4.51	Minimum	2 per subpop	4

5	Nonmonetary Determinations	Single Claimant Nonmon Dets	5.1-5.60	Random	30/100	30/100
		Single Claimant Nonmon Dets	5.1-5.60	Missing Subpops	1 per subpop	≤59
		Single Claimant Nonmon Dets	5.1-5.60	Outliers (TL)	5 highest, 5 lowest	10
		UI Multi-Claimant Determinations	5.61-5.64	Minimum	2 per subpop	8
		Single Claimant Redeterminations	5.65-5.70	Random	30/100	30/100
6	Appeals Filed, Lower Authority	Appeals Filed, Lower Authority	6.1-6.2	Minimum	2 per subpop	4
	Appeals Filed, Higher Authority	Appeals Filed, Higher Authority	7.1-7.2	Minimum	2 per subpop	4
8	Lower Authority Appeals Decisions	Lower Authority Appeals Decisions	8.1-8.52; 8.54-8.55	Random	60/200	60/200
		Lower Authority Appeals Decisions	8.33-8.52; 8.54-8.55	Missing Subpops	1 per subpop	≤21
		Lower Authority Appeals Decisions	8.1-8.52; 8.54-8.55	Outliers (TL)	5 highest, 5 lowest	10
9	Higher Authority Appeals Decisions	Higher Authority Appeals Decisions	9.1-9.20; 9.22-9.23	Random	30/100	30/100
		Higher Authority Appeals Decisions	9.13-9.20; 9.22-9.23	Missing Subpops	1 per subpop	≤ 9
		Higher Authority Appeals Decisions	9.13-9.20; 9.22-9.23	Outliers (TL)	5 highest, 5 lowest	10
10	Appeals Case Aging, Lower Authority	Appeals Case Aging, Lower Auth	10.1-10.7	Outliers (TL)	5 highest, 5 lowest	10
11	Appeals Case Aging, Higher Authority	Appeals Case Aging, Higher Auth	11.1-11.6	Outliers (TL)	5 highest, 5 lowest	10
12	Overpayments Established	Overpayment \$ Established	12.1-12.7; 12.9-12.15	Random	60/200	60/200
		Overpayment \$ Established	12.1-12.7; 12.9-12.15	Missing Subpops	1 per subpop	≤13
		Overpayment \$ Established	12.1-12.7; 12.9-12.15	Outliers (\$)	5 highest, 5 lowest	10
13	Overpayment Reconciliation Activities	Overpayment Reconciliation Activities	13.1-13.34	Random	30/100	30/100
		Overpayment Reconciliation Activities	13.1-13.34	Missing Subpops	1 per subpop	≤33
		Overpayment Reconciliation Activities	13.1-13.34	Outliers (\$)	5 highest, 5 lowest	10
14	Aged Overpayments	Aged Overpayments	14.1-14.12	Random	30/100	30/100
		Aged Overpayments	14.1-14.12	Missing Subpops	1 per subpop	≤11
		Aged Overpayments	14.1-14.12	Outliers (\$)	5 highest, 5 lowest	10

NOTES: The software draws the larger number of Random samples; the first 30 or 60 are investigated as acceptance samples and the remaining 70/140 are only investigated if needed to produce an estimate after an ambiguous result.

Software selects Missing Subpopulation samples on the basis of subpopulations represented in the full 100-case or 200-case draw. Not all subpopulations may be investigated if only first 30 or 60 cases of random sample are reviewed.

Outlier samples may be based on sorts by time lapse (TL), or dollar amount (\$).

B-2. Procedures for the collection of information in which sampling is involved.

- *Statistical methodology for stratification and sample selection*
 - B-1 above indicates that 17 samples are random; 11 are size 30/100, six 60/200. The validation software draws samples of 100 or 200, as required; validators evaluate the first 30 of 100 (60 of 200) as acceptance samples. This often results in a clear pass or fail. If ambiguous findings result, the remaining 70 or 140 are evaluated to estimate underlying error rates.
 - Supplemental samples of size one or two are also drawn from all unrepresented sub-populations to check for the correctness of programming or to ensure that reporting software uses the correct fields in the database.
 - To check for extreme (outlying) values, the 5 highest and 5 lowest values in report elements classified by time lapse (e.g., 7 days and under, 8-14 days, over 70 days) or report fields containing dollars are evaluated.

- *Estimation Procedure*

- Validators must determine whether each underlying population error rate is $\leq 5\%$.
- The DV procedure specifies selection of random samples of 100 or 200, depending on the importance of the underlying transactions.
- The validator uses a sequential review procedure. The first 30 of the full 100, or 60 of the 200, sampled transactions are checked against agency documentation and the number of errors (i.e., those which fail to conform to Federal definitions) are noted.
- The first sequence treats the sampled transactions as acceptance samples of size 30 or 60 to determine whether a judgment can be made at that level or whether review of the remaining cases in the sample is called for. If the result is inconclusive, or the State wishes to estimate the probable underlying error in a population that has clearly failed in the first stage, the additional 60 or 140 sampled transactions are verified and a judgment is made from the 100- or 200-case estimation sample.

- The first stage procedure uses the following decision rules:

	<u>Pass</u>	<u>Fail</u>	<u>Inconclusive</u>
30 Cases	0 errors	≥ 5	1 - 4 errors (evaluate remaining 70 cases)
60 Cases	0 errors	≥ 7	1 - 6 errors (evaluate remaining 140 cases)

These decision rules (as well as those below for the full sample) assume that the samples of transactions are selected without replacement from a large population, and that each transaction in a sampled population of transactions has an equal chance of being selected into the main sample of 100 or 200 and into the subsample of 30 or 60 that is used for the first stage. Based on

these assumptions, the probabilities of any process passing or failing are computed using the binomial formula.¹

- *Degree of Accuracy Needed for Purpose Described in the Justification.*
 - The basic standard is that a reported element is considered to be reported with sufficient accuracy if no more than 5% of the underlying transactions are invalid, i.e., do not conform to Federal definitions for the report element. If the error rate is above 5%, the State's reported counts are considered invalid--even if the reported count equals the reconstructed count--for the report elements involved. This means the State will have to take action to correct the reporting procedure. The sampling procedure must balance the costs of conducting the validation review against the risks of (a) taking an unwarranted and probably expensive action to correct a process whose true underlying error rate is less than 5% and (b) allowing reporting errors to continue by failing to detect underlying populations whose error rates exceed 5%. The Department only *requires* a state to take action on the basis of the evidence of a random sample; the non-random benefits samples described in B-1 above provide diagnostic information but the Department does not require states to act on the findings.
 - The decision rules for the first stage are based on minimizing the chances of failing a sample when the true error rate is acceptable (≤ 0.05). In the first stage, a process passes only with zero errors, and fails if it has 5 or more errors ($n = 30$) or 7 or more errors ($n = 60$). To find these cut-off points (pass, fail)

¹The probability of exactly d events (in this case errors) occurring with n trials where the population prevalence of these events is p (in this case the error rate) is expressed as:

$$P(d) = \binom{n}{d} p^d (1-p)^{n-d} = \left(\frac{n!}{d!(n-d)!} \right) p^d (1-p)^{n-d}$$

The probability that no more than c events occurring is:

$$P(d \leq c) = \sum_{d=0}^c P(d)$$

for the first stage, we calculate the Type I and Type II error contributed from the first stage based on the Binomial distribution with the actual error rate $p = 0.05$.² The cut-off for failing at the first stage is labeled C_1 .

To minimize the Type II error contributed from the first stage we require that there be no error at all to pass the test at the first stage.

To find the optimal cutoff (C_1), we compared Type I errors for different levels of C_1 . The larger C_1 is, the smaller the type I error is. We want to choose C_1 such that the Type I error (p_1)

- is below the 0.05 threshold; and
- is not too close to 0.05 (or too close to 0)

Table 1 gives the type I errors contributed from stage one upon different C_1 's. From the table we can see that: for the sample size n_1 of 30, Type I error would be larger than 0.05 if we choose C_1 at 4. On the other hand, partial Type I error would be too small if we choose C_1 at 6. At $C_1=5$, it is 0.01564, a reasonable number given the criteria above. Hence we decide that the optimal cutoff for $n_1=30$ is 5 and similarly the optimal cutoff for $n_1=60$ is 7.

$$\begin{aligned}
 {}^2 p_1 &= \text{Type I error contributed from first stage} \\
 &= P(\text{rejecting the null when it is true}) \\
 &= P(\geq C_1 \text{ outcomes}) \\
 &= 1 - P(d \leq C_1 - 1) \\
 &= 1 - \sum_{m=0}^{C_1-1} \binom{n_1}{d} p^d (1-p)^{n_1-d}
 \end{aligned}$$

where d is the number of errors

$$\begin{aligned}
 p_2 &= \text{Type II error contributed from the first stage} \\
 &= P(\text{Accepting the null when it is false}) \\
 &= P(0 \text{ outcomes}) \\
 &= (1-p)^{n_1}
 \end{aligned}$$

since for any event d , since $0! = 1$ and $p^0 = 1$,

$$P(d_i = 0) = (1-p)^{n_1}$$

Table 1: Type I Errors From Stage One Upon Different Cutoffs at the First Stage

P	$n_I=30$			$n_I=60$		
	$C_1=4$	$C_1=5$	$C_1=6$	$C_1=6$	$C_1=7$	$C_1=8$
0.01	0.00022	0.00001	0	0.00003	0	0
0.02	0.00289	0.00030	0.000025	0.00127	0.00020	0.00003
0.03	0.01190	0.00185	0.000233	0.00914	0.00210	0.00042
0.04	0.03059	0.00632	0.001061	0.03251	0.00989	0.00262
0.05	0.06077	0.01564	0.003282	0.07872	0.02969	0.00979

- Failure occurs when the number of errors is at least $C_1 = 5$ for $n_I = 30$ and 7 when $n_I = 60$). So the probability of failing can be expressed as 1 minus the probability of not failing where the probability of not failing is the cumulative probability of having fewer than c_i errors.³ The probability of passing at the first stage is the probability of having zero errors. The probabilities of failing in the first stage when the true error rate is ≤ 0.05 and of passing at the first stage if the true error rate is > 0.05 are shown in the following two tables.

Probability of Failing When the Error Rate is ≤ 0.05 (Type I error for first stage of sequential sample)

True Error Rate	$n_I=60$	$n_I=30$
0.01	<.001	<.001
0.02	<.001	<.001
0.03	.002	.002
0.04	.010	.006
0.05	.030	.016

Probability of Passing When the Error Rate is > 0.05 (Type II error for first stage of double sample)

True Error Rate	$n_I=60$	$n_I=30$
0.05	.046	.215
0.06	.024	.156
0.07	.013	.113
0.08	.007	.082
0.09	.003	.059
0.10	.002	.042

³For a given true error rate (p), the probability of failing is:

$$1 - P(\text{not failing}) = 1 - P(d \leq C_1 - 1)$$

- As noted, if the result is inconclusive, the State must evaluate the additional 60 or 140 sampled transactions and make a judgment from the 100- or 200-case estimation sample. (The State may also wish do this to estimate the probable underlying error in a population which has clearly failed in the first stage.)
- In the first stage, the methodology emphasizes avoiding Type II error. In the second stage, it is structured to avoid Type I error. The cut-offs are set to ensure that if the underlying error rate is less than or equal to 5%, the probability that a sample will fail is $< .05$. If the underlying error rate is greater than 5%, probability that a sample will fail is $> .05$ and increases as the underlying rate increases. The Type I error and power probabilities are summarized in Table 2.

○ Thus the second stage decision rule is as follows:

	<u>Conclude Error Rate is</u>	
	<u>$\leq 5\%$</u>	<u>$> 5\%$</u>
Expanded Sample 100	≤ 9 errors	10+ errors
Expanded Sample 200	≤ 16 errors	17+ errors

In the second stage, there are only two outcomes: reject or fail to reject, so we only need to compute the probability of rejecting the null hypothesis knowing the true error rate is p . This probability is the probability of Type I error when the null hypothesis is true and is the power of the test when the null hypothesis is false.⁴

The value of the second stage failure cut-offs C_2 , is that where conditional on Type I error being below the 0.05 threshold, C_2 is such that the power of the test is the largest. Table 2 gives the Type I error and the power of the test for some potential cutoffs. From the table we can see that the optimal cutoff for 30/70 sample is **10** and the optimal cutoff for 60/140 sample is **17**.

⁴ $P(\text{rejecting}) = P(\text{rejecting, first } n1 \text{ conclusive}) + P(\text{rejecting, first } n1 \text{ inconclusive})$

$$\begin{aligned}
 &= p_1 + \sum_{d_1=1}^{C_1-1} P(\text{rejecting, } d_1 \text{ errors in the first } n1) \\
 &= p_1 + \sum_{d_1=1}^{C_1-1} \sum_{d_2=C_2}^{n1} P(d_2 \text{ errors in the second } n2, d_1 \text{ errors in the first } n1) \\
 &= p_1 + \sum_{d_1=1}^{C_1-1} \sum_{d_2=C_2}^{n1} \binom{n1}{d_1} p^{d_1} (1-p)^{n1-d_1} \times \binom{n2}{d_2} p^{d_2} (1-p)^{n2-d_2}
 \end{aligned}$$

Table 2: Type I Error and Power of the Test Upon Different Cutoffs in the Second Stage

Type I error						
P	n =100			n =200		
	C ₂ =9	C ₂ =10	C ₂ =11	C ₂ =16	C ₂ =17	C ₂ =18
	Type I error			Type I error		
0.01	0.000012	0.000012	0.000012	0.000002	0.000002	0
0.02	0.000465	0.000329	0.000305	0.000198	0.000196	0.00020
0.03	0.004622	0.002568	0.002015	0.002419	0.002196	0.00213
0.04	0.022540	0.011884	0.008021	0.015451	0.012147	0.01075
0.05	0.068876	0.038260	0.024241	0.064142	0.047050	0.03789
	Power			Power		
0.05	0.06888	0.03826	0.02424	0.06414	0.04705	0.03789
0.06	0.15310	0.09279	0.05930	0.17911	0.13402	0.10470
0.07	0.27197	0.18072	0.12097	0.36030	0.28608	0.22917
0.08	0.41082	0.29735	0.21151	0.56559	0.47959	0.40341
0.09	0.55088	0.42973	0.32548	0.74364	0.66785	0.59150
0.10	0.67648	0.56208	0.45148	0.86768	0.81414	0.75353

To compute the overall probability that the sample passes, one must take into account the ways in which the sample can pass. We denote the number of errors in the first stage as d_1 and the number from the second stage as d_2 , and the cut-off for the first sample as c_{1i} and for the second as c_{2i} . The smaller sample (30/70), where $c_1 = 5$ and $c_2 = 10$, can pass in any of five ways:

$$\begin{aligned}
 & d_1 = 0, \\
 & d_1 = 1 \text{ and } d_2 < 9 \\
 & d_1 = 2 \text{ and } d_2 < 8 \\
 & d_1 = 3 \text{ and } d_2 < 7 \\
 & d_1 = 4 \text{ and } d_2 < 6
 \end{aligned}$$

For the larger sample, (60/140) the ways the sample can pass follow the same pattern. More generally, the sample will pass if:

$$d_1 = h, \text{ and } d_2 < c_2 - h, \text{ where } h < c_1$$

Given this, we can compute the probability of passing for any underlying error rate, as:

$$p(\text{Pass}) = P(\text{Pass1}) + P(\text{Pass2})$$

$$P(\text{Pass1}) = P(d_1 = 0)$$

$$p(\text{Pass2}) = \sum_{h=1}^{c_1-1} P(d_1 = h)P(d_2 = c_2 - h)$$

The joint results of the two-stage process produce the following probabilities for the two

sample sizes:

<i>Error Rate</i>	<u>Failing a Measure that Should Fail</u>						<u>Failing a Measure that Should Pass</u>				
	<i>.10</i>	<i>.09</i>	<i>.08</i>	<i>.07</i>	<i>.06</i>	<i>.05</i>	<i>.05</i>	<i>.04</i>	<i>.03</i>	<i>.02</i>	<i>.01</i>
Sample											
30/70	.56	.43	.30	.18	.09	.04	.04	.01	.00	.00	.00
60/140	.81	.67	.48	.29	.13	.05	.05	.01	.00	.00	.00

States who fail may wish to examine a confidence region for their observed error rates. In the case where only the initial sample (30 or 60) has been examined, construction of a confidence region is straightforward. Where the full sample ($n = 100$ or 200) has been examined, the process is more complex. Below, lower confidence bounds are presented for states to use. Lower bounds are presented instead of confidence intervals, because states with high observed error rates are more likely to find this measure of sampling error useful.⁶

As discussed above, in determining whether a sample passed or failed the states will test for each sample the null hypothesis that the true error rate is less than or equal to 0.05. Constructing a lower confidence bound for an observed error rate (p^*) is analogous to the pass/fail determination. It can be thought of as testing a hypothesis. However, to construct the confidence bound, the test is of a different hypothesis: the true error rate equals the one observed (i.e., $p=p^*$) versus the alternative that the true error rate is less. Thus, the procedures for finding a lower confidence limit are analogous to those in determining the pass or fail cut-off points.

For constructing the confidence bounds the initial samples ($n = 30$ or 60) can be treated as simple random samples with size nl from a Binomial distribution.

Therefore for an observed number of errors d_o the corresponding lower confidence bound is determined by finding $p \in [0,1]$, such that

$$R(p^*) = P(d \leq d_o) \text{ errors} = \sum_{i=0}^{d_o} \binom{nl}{d_i} p^{d_i} (1-p)^{nl-d_i} = 1 - \alpha$$

⁶Confidence intervals or sets do not seem to be covered in industrial quality control, where the sequential sampling procedures described in this section are often used. In these settings, the concern is only with whether the batch or sample passed or failed, not with the precision of the observed error rate.

$R(p^*)$ is a decreasing function of p . For example, when $n_1=30$ and $d_o=4$, ($p^*=.133$). For $\alpha =0.05$, the corresponding solution is 0.069 so the lower 95 percent bound would be 0.069.

The following table gives the lower 95% confidence bound for $n_1=30$ and $n_1=60$ respectively.

Table 3: The Lower Confidence Bound for Simple Random Sampling

Errors	$n_1 = 30$		$n_1 = 60$		
	Error Rate	Lower Bound (95%)	Errors	Error Rate	
0	0.000	0.002	00.000	0.002	0
1	0.033	N/A	10.033	N/A	1
2	0.067	N/A	20.067	N/A	2
3	0.100	N/A	30.100	N/A	3
4	0.133	N/A	40.133	N/A	4
5	0.167	0.091	50.167	0.091	5
6	0.200	0.115	60.200	0.115	6
7	0.233	0.141	70.233	0.141	7
8	0.267	0.167	80.267	0.167	8
9	0.300	0.194	90.300	0.194	9
10	0.333	0.222	100.333	0.222	10
11	0.367	0.250	110.367	0.250	11
12	0.400	0.279	120.400	0.279	12
13	0.433	0.309	130.433	0.309	13
14	0.467	0.339	140.467	0.339	14
15	0.500	0.370	150.500	0.370	15
16	0.533	0.402	160.533	0.402	16
17	0.567	0.434	170.567	0.434	17
18	0.600	0.467	180.600	0.467	18
19	0.633	0.501	190.633	0.501	19
20	0.667	0.535	200.667	0.535	20

For $n_1 = 30$, 1 to 4 errors in the first sample will result in the second-stage sample ($n_2 = 70$) being selected and for $n_1 = 60$, 1 to 6 errors will result in the second-stage sample ($n_2 = 140$) being selected. Because in these instances the error rate will be based on the full sample ($n=100$ or $n= 200$), the lower confidence limits will be found in Table 5, and hence they are designated as N/A in this table.

When both samples are used, errors are observed from both samples and the samples are not independently selected (the second sample is used only if the first sample is inconclusive). So to construct a lower bound for this case we begin in a manner analogous to setting the cut off points for failing when the purpose is to determine whether the sample passes or fails.

Thus, the lower bound is the smallest value of p such that:

$$H_0 : P \leq p^*$$

is accepted (p^* is the observed error rate). With this criterion, one can define a decision rule for the sequential sampling. (The method for the decision rule has already been illustrated above.) For example for the 30/70 sample, Table 4 gives the optimal cutoff for some illustrative error rates.

Table 4: The Optimal Cutoff for p^* in Sequential Sampling

P	n1 = 30		n1 = 60		
	C_1	C_2	P	C_1	C_2
0.06	6	11	0.06	7	20
0.07	6	13	0.07	9	22
0.08	6	15	0.08	10	24
0.09	7	15	0.09	11	26
0.10	7	17	0.10	12	29
0.15	9	23	0.15	15	41
0.20	11	29	0.20	19	52

For each observed pair of errors, the lower 95% confidence bound is the first p that the null hypothesis is going to be accepted upon this p . For example, if there are 2 errors in the first stage and 5 errors overall, the smallest p such that the null is accepted upon p is 0.020. Table 5 gives the 95 percent lower bound for the case where both samples are used.

Unusual Problems Requiring Specialized Sampling Procedures

- o The discussion above indicates that the methodology uses specialized sampling procedures. Strictly speaking, none of these are required. However, because of the scope of UI DV, they are employed for efficiency. Most State UI management information systems are highly automated, and States are able to obtain most data elements they report to the Department of Labor as a byproduct of their ongoing functions of paying benefits and collecting taxes. Thus, the greatest risks to report validity are from systematic errors—incorrectly programmed functions which miss certain elements, double count other elements, obtaining counts of transactions which do not meet the Federal reporting requirements for the element being reported, or programming which

Table 5: The Lower (95%) Confidence Bound for Sequential Sampling

Errors			N=30/70		N=60/140	
Total	From n1	From n2	Error Rate	Lower Confidence Bound	Error Rate	Lower Confidence Bound
1	1	0	0.010	0.002	0.005	0.001
2	1	1	0.020	0.002	0.010	0.001
2	2	0	0.020	0.010	0.010	0.005
3	1	2	0.030	0.009	0.015	0.005
3	2	1	0.030	0.010	0.015	0.005
3	3	0	0.030	0.023	0.015	0.012
4	1	3	0.040	0.015	0.020	0.008
4	2	2	0.040	0.015	0.020	0.008
4	3	1	0.040	0.023	0.020	0.012
4	4	0	0.040	0.040	0.020	0.020
5	1	4	0.050	0.020	0.025	0.010
5	2	3	0.050	0.020	0.025	0.010
5	3	2	0.050	0.023	0.025	0.012
5	4	1	0.050	0.040	0.025	0.020
6	1	5	0.060	0.027	0.030	0.014
6	2	4	0.060	0.027	0.030	0.014
6	3	3	0.060	0.027	0.030	0.014
6	4	2	0.060	0.040	0.030	0.020
7	1	6	0.070	0.033	0.035	0.017
7	2	5	0.070	0.033	0.035	0.017
7	3	4	0.070	0.033	0.035	0.017
7	4	3	0.070	0.040	0.035	0.020
8	1	7	0.080	0.038	0.040	0.019
8	2	6	0.080	0.038	0.040	0.019
8	3	5	0.080	0.038	0.040	0.019
8	4	4	0.080	0.041	0.040	0.020
9	1,2,3,4	8,7,6,5	0.090	0.047	0.045	0.024
10	1,2,3,4	9,8,7,6	0.100	0.053	0.050	0.027
11	1,2,3,4	10,9,8,7	0.110	0.058	0.055	0.031
12	1,2,3,4	11,10,9,8	0.120	0.069	0.060	0.034
13	1,2,3,4	12,11,10,9	0.130	0.075	0.065	0.037
14	1,2,3,4	13,12,11,10	0.140	0.080	0.070	0.042
15	1,2,3,4	14,13,12,11	0.150	0.092	0.075	0.046
16	1,2,3,4	15,14,13,12	0.160	0.098	0.080	0.049
17	1,2,3,4	16,15,14,13	0.170	0.110	0.085	0.054
18	1,2,3,4	17,16,15,14	0.180	0.116	0.090	0.057
19	1,2,3,4	18,17,16,15	0.190	0.123	0.095	0.060
20	1,2,3,4	19,18,17,16	0.200	0.134	0.100	0.066

reflects a misinterpretation of Federal reporting requirements. Systematic problems normally affect all elements in a population grouping, so the examination of just a few is sufficient to identify the problem. A larger, random sample would of course identify the same problem but at much higher cost. Similarly, large random samples would probably detect the existence of outliers in time lapse data or data involving the reporting of dollar amounts. However, small samples of transactions from the extremes of an arrayed distribution do it much more efficiently.

- *Use of Periodic Data Collection to Reduce Burden*
 - UI DV employs a 3-year cycle to reduce burden. Only the components which fail validation (a discrepancy between a reported count and a reconstructed count greater than 2%, or quality samples showing more than a 5% rate of invalid cases in the population examined) must be revalidated in the following year.
 - The exception is the report cells used to calculate Government Employment and Results Act measures. These must be validated annually, and the reported count must be within $\pm 1\%$ of the reconstructed count.

B-3. *Methods to Maximize Response Rates.*

Although this collection is based on agency records, our experience to date does indicate non-response in the sense that some states have not been able to complete all or part of data validation. In some cases, state resources have precluded them from doing all or part of DV. In others, they have deferred part of DV pending the installation of new administrative data systems. There have been a few instances where the validation methodology cannot be applied because the state reports are not automated, or the state validators have concluded that their reports cannot pass validation or be completely validated because their data systems lack key information, e.g., the data a receivable was established. In all these instances, states are required to include corrective action plans to complete implementation of UIDV or to fix their reports and submit their UI DV reports as part of their annual performance management and budgeting plan (called the State Quality Service Plan). In the course of validations, states often discover that the documentation for certain reported transactions--e.g., nonmonetary determinations or benefit appeals--is missing. In considering which transactions have been reported accurately, validation does not distinguish between missing documentation and other forms of errors; if a transaction is not adequately documented it is considered an error.

B-4. *Tests of Procedures or Methods.*

- In 1998, three States--Massachusetts, Minnesota, and North Carolina--pilot tested the UI DV methodology. A technical support contractor, who employed as a

subcontractor the person who developed the UI DV methodology, provided oversight of the pilot test. The contractor's evaluation indicated that the methodology functioned as intended and enabled the States to detect, and correct, reporting errors. The cost data from the pilot were the basis for the burden estimates in the original request. In the first three years of authorization, most states have completed at least parts of validation requirements. Burden estimates for this request are based on estimates provided by states that have completed validations, and reflect assumptions consistent with a new software environment.

B-5. Individuals Consulted on Statistical Aspects of the Design

William S. Borden
Senior Fellow
Mathematica Policy Research, Inc.
(609) 275-2321

John Hall
Senior Statistician
Mathematica Policy Research, Inc.
(609) 799-3535

Walter Corson
Senior Fellow
Mathematica Policy Research, Inc.
(609) 799-3535

Alan Dorfman
Senior Mathematical Statistician
Office of Survey Methods Research
Bureau of Labor Statistics
(202) 691-7378

Steven Marcus
President
Sparhawk Group, Inc.
(617) 787-0388

Andrew Spisak
Mathematical Statistician
Office of Workforce Security
Employment and Training Administration
(202) 693-3196