

Attachment C

Bioterrorism and Mass Casualty Preparedness in Hospitals: United States: 2003, Advance Data No. 364

Bioterrorism and Mass Casualty Preparedness in Hospitals: United States, 2003

by Richard W. Niska, M.D., M.P.H., and Catharine W. Burt, Ed.D., Division of Health Care Statistics

Abstract

Objectives—This study examined the content of hospital terrorism preparedness emergency response plans; whether those plans had been updated since September 11, 2001; collaboration of hospitals with outside organizations; clinician training in the management of biological, chemical, explosive, and nuclear exposures; drills on the response plans; and equipment and bed capacity.

Methods—The National Hospital Ambulatory Medical Care Survey (NHAMCS) is an annual survey of a probability sample of approximately 500 non-Federal general and short-stay hospitals in the United States. A Bioterrorism and Mass Casualty Supplement was included in the 2003 survey and provided the data for this analysis.

Results—Almost all hospitals have plans for responding to natural disasters (97.3 percent). Most have plans for responding to chemical (85.5 percent), biological (84.8 percent), nuclear or radiological (77.2 percent), and explosive incidents (76.9 percent). About three-quarters of hospitals were integrated into community-wide disaster plans (76.4 percent), and 75.9 percent specifically reported a cooperative planning process with other local health care facilities. Despite these plans, only 46.1 percent reported written memoranda of understanding with these facilities to accept inpatients during a declared disaster. Hospitals varied widely in their plans for re-arranging schedules and space in the event of a disaster. Training for hospital incident command and smallpox, anthrax, chemical, and radiological exposures was ahead of training for other infectious diseases. The percentage of hospitals training their staff in any exposure varied from 92.1 percent for nurses to 49.2 percent for medical residents. Drills for natural disasters occurred more often than those for chemical, biological, explosive, nuclear, and epidemic incidents. More hospitals staged drills for biological attacks than for severe epidemics. Despite explosions being the most common form of terrorism, drills for these were staged by only one-fifth of hospitals. Hospitals collaborated on drills most often with emergency medical services, fire departments, and law enforcement agencies.

Keywords: bioterrorism • emergency response plans • medical training • mass casualty drills • medical equipment

Introduction

In response to the terrorism incidents of September 2001, the Office of the Assistant Secretary of Planning and Evaluation, Department of Health and Human Services, requested that the Centers for Disease Control and Prevention's (CDC) National Center for Health Statistics (NCHS) survey hospitals regarding their preparedness for treating patients from bioterrorism attacks or mass casualty incidents. It provided funding to field a special supplement to the 2003 National Hospital Ambulatory Medical Care Survey (NHAMCS).

Data from the 2003 NHAMCS supplement will serve as a baseline for hospital terrorism preparedness because there were no major funding programs directed toward hospitals for this purpose prior to the start of the Bioterrorism Hospital Preparedness Program of the Health Resources and Services Administration (HRSA) in fiscal year (FY) 2002. Even these funds had not yet been fully awarded to hospitals at the time this survey was fielded in 2003 (1,2). Prior to the inception of the HRSA program, the Domestic Preparedness Program of the Department of Defense had not allocated any funding for hospitals, although it funded emergency response agencies (3).



Because the NHAMCS terrorism supplement was also repeated in 2004, it will be possible to study changes in preparedness as a function of the funding made available through major Federal programs. Information about strengths and limitations of terrorism preparedness in the Nation's hospitals will be crucial for appropriate Federal agencies that are charged with planning how future funding should be used to improve domestic defensive posture.

Hospitals would be among the first institutions affected after deployment of a weapon of mass destruction. They would treat those injured in terrorist attacks in ways not unlike those used for other incidents where large numbers of patients are involved. After a biological attack, emergency departments might be the first to note changes in the epidemiology of an infectious disease, to initiate a public health department referral to identify the causative organism, and to treat patients affected by the exposure. After a chemical or radiological attack, patients might require decontamination at the hospital if they have bypassed the firefighters or emergency medical technicians at the exposure site. After an explosion, large numbers of injured patients might be brought to emergency departments for treatment. In any of these scenarios, hospitals would experience an influx of patients who may have been exposed and have medical or psychological issues to be addressed.

Because of the heavy demand placed on their services at the time of an attack, hospitals need to be prepared to handle the workload. Because the most common terrorist attacks to date have been explosive or incendiary (car bombs, airplanes full of fuel, etc.), hospitals must be prepared to treat an influx of trauma cases. Hospitals must also be prepared to diagnose and treat diseases caused by CDC Class A bioterrorism agents (smallpox, anthrax, plague, botulism, tularemia, and hemorrhagic fever). Although hospitals are required to have disaster response plans to be accredited by the Joint Commission on Accreditation of Healthcare Organizations (JCAHO), the

standard elements of these plans are still quite general regarding terrorism (4).

The 2003 NHAMCS Bioterrorism and Mass Casualty Preparedness Supplement surveyed hospitals on their preparedness for such attacks by asking about the content of emergency response plans, training for terrorism response, experiences with internal and external disaster drills, and availability of specialized equipment such as decontamination showers, personal protective suits, and negative pressure isolation rooms.

This report provides a descriptive summary of the responses to the supplement from which national estimates of preparedness can be generated for the overall universe of U.S. hospitals. Other current research efforts involving the survey go beyond the descriptive work reported here and delve into hospital characteristics associated with greater preparedness in the areas of emergency response planning, training and practicing for terrorism, and equipment and specialized care units for terrorism-related incidents.

Methods

The NHAMCS is an annual probability sample survey of non-Federal general and short-stay hospitals (those with an average length of stay of less than 30 days) within the 50 States and the District of Columbia. Children's hospitals were included in the sampling frame. Federal hospitals, hospital units of institutions, and hospitals with less than six beds were ineligible for the NHAMCS. Hospitals with a 24-hour emergency department (ED) or outpatient department (OPD) supervised by a physician were included. Hospitals without an ED or whose OPD was comprised solely of ancillary clinics (e.g., radiology, laboratory services, physical rehabilitation, renal dialysis) were ineligible to participate. The sampling frame consisted of hospitals listed in the 1991 Verispan Hospital Database (VHD), which was updated using the 2000 VHD to allow the inclusion of hospitals that opened or changed their eligibility status since the previous sample in 1991. The VHD was formerly known as the SMG Hospital

Database. An additional sample of 66 hospitals (identified as proprietary or as located in nonmetropolitan statistical areas) was selected from the 2002 VHD. Hospitals were randomly assigned to 1 of 13 4-week reporting periods scattered over the entire year. The data were weighted according to the inverse probability of hospital selection and a nonresponse adjustment factor. Thus, estimates are considered representative of similar health care facilities over the entire Nation.

Terrorism preparedness items were added to the 2003 NHAMCS in a special supplement (see "Technical Notes"). This was administered to the same hospitals as those sampled for the NHAMCS. The supplement used the first 2 stages of the regular NHAMCS design involving 112 geographic primary sampling units (PSU) and hospitals that have emergency or outpatient departments within PSUs. For this analysis, the supplement included questions about updates of the hospital's emergency response plan since September 11, 2001; whether that plan addressed natural disasters and biological, chemical, nuclear-radiological, and explosive-incendiary terrorism incidents; collaboration with outside organizations for planning; key elements of the plans such as establishment of alternate care sites; clinician training in the management of exposures to weaponizable infectious diseases, chemicals, and nuclear materials; drills on aspects of the response plans; and equipment and bed capacity available at the hospital. The content of the questionnaire was based on the JCAHO list of elements that are important to have in an emergency response plan, training questions about the CDC Class A bioterrorism agents, and drill scenarios suggested by bioterrorism researchers at HRSA, Johns Hopkins University, George Washington University, the American Hospital Association, the Office of Emergency Preparedness at the Department of Health and Human Services, the Government Accountability Office, the Office of Management and Budget, and the Centers for Medicare and Medicaid Services. The collection form was pilot tested in October 2002 and refined and

implemented in December 2002. During the 2003 NHAMCS induction interview with the administrator of each sampled hospital, the supplement was left for the bioterrorism coordinator to complete and return to the interviewer. Of 546 hospitals sampled in 2003, 462 were in scope for the NHAMCS. Data were received from 399 hospitals, for a response rate of 86.4 percent. This represented a weighted national total of approximately 5,000 hospitals. The percentage of missing values (re-coded into "no" or "blank" responses for any one variable) was less than 3%, except where noted in the tables. Descriptive statistics with their corresponding measures of sampling error were generated using SUDAAN-9.0 software (RTI International), which takes into account the complex sample design of the NHAMCS (5). Analyses involving comparisons across subgroups used a chi-square test statistic evaluated at a 0.05 level of significance.

Results

Emergency Response Plans

Nearly all hospitals had emergency terrorism response plans. The majority had revised their plans since September 11, 2001 (92.3 percent), but 6.3 percent had not. Few hospitals had no response plan (1.5 percent). Table 1 shows the percent distribution of all hospitals (with or without plans) and the percentage of hospitals with revised plans by selected hospital characteristics. Only teaching hospital status was associated with a greater likelihood of having a revised plan ($p < 0.05$). Among hospitals with a response plan, 97.3 percent addressed natural disasters, 85.5 percent addressed chemical terrorism, 84.8 percent addressed biological terrorism, 77.2 percent addressed nuclear or radiological terrorism, 76.9 percent addressed explosive or incendiary incidents, and 63.5 percent addressed all five of these incident types (table 2).

Almost all hospitals (95.4 percent) had provisions in their plans to contact outside entities. The most frequent outside entities were State or local public health departments (81.9 percent), State or local law enforcement

(77.7 percent), emergency medical services (71.9 percent), and fire departments (65.7 percent) (table 3). About three-quarters of hospitals (76.4 percent) defined their role in community-wide planning in their response plans, and 75.9 percent provided for cooperative planning with other area health care facilities. However, only 46.1 percent of hospitals had memoranda of understanding with outlying hospitals to accept inpatients during a declared disaster (table 4).

In the area of internal disaster planning, 73.1 percent of hospitals planned to cancel elective procedures and admissions during an emergency, 66.8 percent had plans for establishing an alternate care site, and 60.8 percent had plans for medical utilization of nonclinical hospital space. However, only 36.6 percent of hospitals planned to convert their postanesthesia care unit (where postoperative patients are monitored as they awake from anesthesia) to augment intensive care capacity. Only 27.3 percent planned to activate decommissioned ward space (i.e., opening closed hospital bed capacity) during an emergency. The majority of hospitals (60.1 percent) provided for coordinated supply-chain management of critical supplies and pharmaceuticals in their response plans, which includes such methods as vendor-managed pharmaceutical inventories to ensure that supplies are replaced before becoming outdated. Less than half (44.1 percent) of hospitals had plans to stockpile antibiotics and other supplies (table 4).

Most hospitals (87.4 percent) were members of a regional interagency disaster preparedness task force. Only about one-third of hospitals (34.3 percent) were designated to receive patients through the National Disaster Medical System (NDMS). Another one-third of hospitals were unaware of their NDMS designation status (table 5).

Training in Emergency Response

Three-quarters (75.0 percent) of hospitals had key hospital personnel trained to implement a formal incident command system, which is an organized procedure for managing resources and personnel during an emergency (table 5).

Staff members at most hospitals (88.5 percent) had received training since September 11, 2001, in the identification, diagnosis, and treatment of any of the biological agents studied in this survey (smallpox, anthrax, plague, botulism, tularemia, viral hemorrhagic fever, viral encephalitis) (figure 1). Training on individual diseases or conditions received by hospital staff physicians, interns, residents, physician assistants, registered nurses, licensed practical nurses, laboratory staff, and others was most frequent for smallpox (86.7 percent), anthrax (83.7 percent), and chemical exposures (80.2 percent) (table 6 and figures 1 and 2). Data on training by professional category are shown in table 7 and figure 3. The most frequent sources of preparedness training were hospitals (61.4 percent) and State or




Figure 1. Percentage of hospitals that trained their staff in emergency response, by selected subject areas: United States, 2003




Figure 2. Percentage of hospitals that trained their staff in bioterrorism response, by biological agent: United States, 2003

local public health departments (44.8 percent) (table 8).

Mass Casualty Drills

Nine out of 10 hospitals (88.4 percent) participated in an internal mass casualty drill since September 11, 2001, and 87.1 percent conducted a mass casualty drill in collaboration with other organizations. For the internal drills, the most common scenario was a

general disaster and emergency response (69.7 percent). Other scenarios included chemical release (44.9 percent), biological attack (37.5 percent), explosive or incendiary attack (21.3 percent), and nuclear or radiological attack (15.4 percent). In contrast to biological attacks, drills involving severe epidemics were staged by only 7.1 percent of hospitals (table 9). Scenarios for collaborative drills followed a similar pattern. The




Figure 3. Percentage of hospitals that trained their staff in terrorism response, by professional category: United States, 2003

organizations with which hospitals collaborated most often for mass casualty drills were emergency medical services (70.9 percent), fire departments (67.1 percent), and State or local law enforcement (64.5 percent) (table 10).

Mass Casualty Resources

Hospitals were asked to quantify certain available resources that would be useful in a mass casualty incident. Among hospitals with mechanical ventilators, there was an average of 7.8 per 100 beds (table 11). Among hospitals with personal protective hazardous materials suits, there was an average of 9.4 per 100 beds. On average, there were 4.3 negative pressure isolation rooms and 13.8 critical care beds per 100 beds, including those in adult and pediatric intensive care, coronary care, and postanesthesia care units. Among hospitals with decontamination showers, there was an average of 1.5 per 100 beds, but the capacity of these showers was not ascertained in this survey.

Discussion

Almost all hospitals have natural disaster plans because it is a JCAHO accreditation requirement (4) to hold annual disaster drills on scenarios such as floods, hurricanes, earthquakes, tornados, or others that have stressed health care systems on a regular basis. Although it is natural to focus on natural disasters during preparedness planning, some gaps in preparedness for man-made disasters would be expected because terrorism is not within the usual experience of U.S. hospitals. In a convenience sample of 30 hospitals from West Virginia, Pennsylvania, Maryland, Virginia, and the District of Columbia, Treat et al. (6) found that only 27 percent had addressed weapons of mass destruction in their disaster plans. In this pre-September 11 survey, none of the rural hospital respondents believed that their hospitals were prepared for biological, chemical, or nuclear disasters. Most of the urban hospital respondents believed their hospitals were only somewhat prepared. In a 1998 survey of 186 emergency

departments in Alaska, Idaho, Oregon, and Washington, Wetter et al. (3) found that 80 percent of the hospitals had hazardous materials response plans. However, only 17 percent had chemical weapons plans, and only 12 percent had plans for biological weapons response.

The NHAMCS supplement was conducted throughout 2003 and reflects a heightened level of urgency due to the airline terrorist attacks of September 11, 2001, and the anthrax outbreak of 2001. Planning for biological scenarios was quite common despite the rarity of these events in actual experience. The Government Accountability Office (GAO) surveyed 1,482 urban hospitals between May and September 2002 as to their preparedness for biological terrorism. Approximately 81 percent of these hospitals had a written emergency response plan that addressed bioterrorism (7). The NHAMCS results for 2003 show a further increase to 94 percent for urban hospitals having revised emergency response plans for terrorism in general (table 1). Because funding from the Bioterrorism Hospital Preparedness Program of the Health Resources and Services Administration was focused exclusively on biological preparedness until FY 2003, this may have also played a role in the emphasis on this contingency (8). However, because funds from that program had not been fully distributed to hospitals by the time the 2003 NHAMCS was fielded, it will be interesting to follow up on this issue in the 2004 iteration of this survey (1,2).

Planning for chemical attacks is similar to that for biological attacks and ahead of nuclear-radiological exposures. Many communities have hazardous materials teams because of the high potential for industrial or farming exposures. Also, there has been experience in recent history with chemical attacks, such as the sarin incident in Tokyo (9). Wetter et al. (3) found that hospitals within 35 miles of a military chemical weapons depot were five times as likely as other hospitals to have a chemical weapons response plan, but no more likely to have a plan for biological weapons.

Plans for explosive or incendiary incidents were reported the least

frequently of all the scenarios in this survey. However, one would expect planning for explosions to rank much higher than for other forms of terrorism because this is the most frequent means employed by terrorists, both domestically (e.g., World Trade Center) and internationally (e.g., Iraq, Madrid, and London). But the \$125 million of funding from the HRSA Bioterrorism Hospital Preparedness Program did not cover explosions or fires in FY 2002, and separate HRSA funding for Statewide trauma systems nationwide has been level at approximately \$2 million per year from FY 2001 to FY 2004 (10).

The majority of hospitals had provisions in their response plans for cooperation with appropriate outside entities. However, less than half reported having actual written memoranda of understanding with other facilities on at least one dimension of mutual aid—being able to transfer patients elsewhere when their surge capacity is exceeded. This replicated the pre-September 11 findings of Treat et al. (6), who found that although 97 percent of their hospitals had patient overflow plans involving other facilities, none of them had specific agreements in place for mass disaster casualties. The post-September 11 GAO study was more optimistic for urban hospitals in that 69 percent of their sample had agreements to share resources during a biological incident (7). One area for future research might be to frame the questions in terms of actual arrangements in place instead of plans to do so.

One interesting anomaly in the results was the high percentage of hospitals that were unaware of whether they were designated by NDMS to receive patients in a disaster. It is possible that the NDMS designation had been granted at some time in the distant past, but never actually tested because of the low prevalence of mass casualty disasters. Given that, it may not be surprising that the institutional memory of such a designation might have been lost, especially if the hospital had undergone changes in ownership or administration since it was granted. This finding suggests that hospital drills

might be crafted to reflect the NDMS designation status, thus encouraging the hospital to review its status.

It is encouraging that such a high percentage of registered and licensed practical nurses had received training in at least one area of terrorism preparedness, and that staff physicians were not far behind. However, it is puzzling that interns and residents lag so far behind their senior colleagues because one might expect that physicians in training would be at the forefront of acquiring new knowledge. The high percentage of nurses who received training is reassuring because the pre-2001 survey done by Treat et al. (6) found that staff at less than one-quarter of hospitals in the States surrounding Washington, DC, had received education on weapons of mass destruction. Wetter et al. (3) found that only 23 percent of hospitals in the Northwest had offered training before September 11 on chemical weapons response, and only 19 percent had offered biological weapons response training.

In their 2002 survey, the GAO found that 70 percent of urban hospitals had trained their staff for all six of the CDC Class A biological agents (smallpox, anthrax, plague, botulism, tularemia, and hemorrhagic fever) (7). However, in a separate analysis of urban hospitals responding to the 2003 NHAMCS supplement, none of the sample had trained their staff in all six agents. It is possible that differing methodologies might be partially responsible for this discrepancy. The GAO used a mailed survey with followup reminders by mail and telephone, and the NHAMCS survey was presented to the bioterrorism coordinators by on-site field representatives during personal interviews. However, although the GAO report was more optimistic than the NHAMCS for five of the six agents individually (smallpox being equal), the discrepancies were not as marked. For example, at the high end of the scale, GAO found that 93 percent of urban hospitals had offered training for anthrax, and 86 percent of the NHAMCS urban sample had. At the low end, GAO found that 71 percent of

urban hospitals had offered training for viral hemorrhagic fever, and only 53 percent of the NHAMCS urban sample had.

The top training priorities that are demonstrated in this study are interesting because they are a mix of more likely occurrences (such as chemical incidents) and less likely, but high profile occurrences (such as smallpox and anthrax). Training for the rest of the infectious diseases is perhaps more reflective of their rarity. Again, this demonstrates a post-September 11 shift in thinking. The Treat et al. (6) survey found that only 10 percent of hospitals drilled on chemical or nuclear attacks, and only 3 percent drilled on biological weapons.

It is interesting to note the discrepancy between hospital drills for biological attacks and those for severe epidemics. The percentage of hospitals drilling on biological scenarios more closely approximates that for chemical attacks. Future research might define more precisely what skills are being drilled in a biological attack response, especially whether biological response is being thought of in terms of acute decontamination of a known aerosol exposure rather than in terms of an infectious disease epidemic beginning some days after an apparent exposure.

In the aftermath of the devastation wrought by Hurricane Katrina, this report provides information that may assist in the formulation of future policies and procedures on preparedness for such natural disasters. Hospitals reported that their drills lagged behind their written response plans, formal patient transfer arrangements lagged behind cooperative planning with other hospitals, and drills that included public health departments and volunteer organizations lagged behind drills that included emergency medical services and fire departments. Knowledge of these shortfalls may offer opportunities for future improvement efforts. As hospitals review their plans in light of the problems experienced during this disaster, the lessons learned may also affect future data collection on the

ability of hospitals to respond to all types of emergencies.

References

1. Hearne SA, Davis M, Segal LM, et al. Ready or not? Protecting the public's health in the age of bioterrorism. Trust for America's Health. December 2003. www.healthyamericans.org.
2. McHugh M, Staiti AB, Felland LE. How prepared are Americans for public health emergencies? Twelve communities weigh in. *Health Aff*; 23:201-9; May-June 2004.
3. Wetter DC, Daniell WE, Treser CD. Hospital preparedness for victims of chemical or biological terrorism. *Am J Public Health*. 91:710-6; May 2001.
4. Joint Commission on Accreditation of Healthcare Organizations. Emergency management standards — EC.2.9.1. Joint Commission Resources, Oakbrook Terrace, Illinois. January 1, 2003.
5. Research Triangle Institute SUDAAN (Release 9.0.1) [Computer Software]. Research Triangle Park, NC: Research Triangle Institute. 2005.
6. Treat KN, Williams JM, Furbee PM, et al. Hospital preparedness for weapons of mass destruction incidents: an initial assessment. *Ann Emerg Med*. 38:562-5; November 2001.
7. General Accounting Office. Hospital preparedness — most urban hospitals have emergency plans but lack certain capacities for bioterrorism response. GAO-03-924. Washington. U.S. General Accounting Office. 2003.
8. Health Resources and Services Administration. National bioterrorism hospital preparedness program cooperative agreement guidance. 2003.
9. Arizona Department of Health Services. History of biowarfare and bioterrorism. August 3, 2004. www.azdhs.gov/phs/edc/edrp/es/bthistor2.htm.
10. Health Resources and Services Administration. FY 2004 State trauma project abstracts and funding profiles. www.hrsa.gov/trauma/states/.

Table 1. Percent distribution of all hospitals and percentage of hospitals having revised emergency response plans since September 11, 2001 (with standard errors), by hospital characteristics: United States, 2003

| Hospital characteristic | Percent distribution | Standard error | Percent with plan revised since September 11, 2001 | Standard error |
|--|----------------------|----------------|--|----------------|
| All hospitals | 100.0 | ... | 92.3 | 2.3 |
| Geographic region | | | | |
| Northeast | 16.3 | 1.6 | 92.3 | 3.5 |
| Midwest | 28.3 | 2.8 | 91.3 | 4.5 |
| South | 37.7 | 2.9 | 90.5 | 4.6 |
| West | 17.8 | 2.7 | 97.5 | 1.7 |
| Metropolitan status | | | | |
| Metropolitan statistical area | 60.0 | 3.6 | 94.2 | 2.5 |
| Nonmetropolitan statistical area | 40.0 | 3.6 | 89.4 | 4.3 |
| Ownership | | | | |
| Voluntary, nonprofit | 62.0 | 4.3 | 94.1 | 2.1 |
| Government | 26.3 | 4.2 | 84.5 | 6.2 |
| Proprietary | 11.7 | 2.1 | 99.8 | 0.2 |
| Teaching hospital | | | | |
| Yes | 6.3 | 0.8 | 100.0 | 0.0 |
| No | 93.7 | 0.8 | 91.7 | 2.4 |
| Number of hospital beds | | | | |
| Less than 100 | 59.0 | 2.7 | 88.6 | 3.7 |
| 100–199 | 20.4 | 2.4 | 97.7 | 2.0 |
| 200–299 | 10.0 | 1.6 | 97.6 | 1.8 |
| 300 or more | 10.7 | 1.2 | 97.0 | 2.2 |
| Has a 24-hour emergency department | | | | |
| Yes | 92.7 | 2.9 | 91.7 | 2.4 |
| No | *7.3 | 2.9 | 99.2 | 0.9 |

... Category not applicable.

0.0 Quantity more than zero but less than 0.05.

* Figure does not meet standards of reliability or precision.

NOTE: Numbers may not add to totals because of rounding.

Table 2. Percentage (with standard errors) of hospitals with emergency response plans for selected types of incidents: United States, 2003

| Type of incident | Percent of hospitals | Standard error |
|--|----------------------|----------------|
| Natural disaster | 97.3 | 1.0 |
| Chemical | 85.5 | 2.4 |
| Biological | 84.8 | 2.6 |
| Nuclear or radiologic exposure | 77.2 | 3.5 |
| Explosive or incendiary exposure | 76.9 | 4.0 |
| All types of incidents | 63.5 | 4.3 |

Table 3. Percentage (with standard errors) of hospitals having provisions in their emergency bioterror response plans for contacting outside entities, by type of outside entity to be contacted: United States, 2003

| Type of outside entity contact | Percent of hospitals | Standard error |
|--|----------------------|----------------|
| Any entity | 95.4 | 1.6 |
| State or local health departments | 81.9 | 3.4 |
| State or local police | 77.7 | 2.9 |
| Emergency medical services | 71.9 | 3.9 |
| Fire department | 65.7 | 4.1 |
| Other State or local government | 59.9 | 3.5 |
| Hazardous materials teams | 58.3 | 3.9 |
| Centers for Disease Control and Prevention | 52.4 | 4.1 |
| Other hospitals | 50.7 | 4.0 |
| Federal Bureau of Investigation | 42.5 | 4.1 |
| Utilities | 32.0 | 3.6 |
| Vendors | 28.9 | 4.1 |
| Local political official | 19.7 | 2.4 |
| Other Federal agency | 11.7 | 1.7 |
| Other laboratories | 8.4 | 2.2 |

Table 4. Percentage (with standard errors) of hospitals having specific components in their emergency response plans, by type of plan component: United States, 2003

| Type of plan component | Percent of hospitals | Standard error |
|--|----------------------|----------------|
| Definition of role in community planning | 76.4 | 3.5 |
| Cooperative planning with other health care facilities | 75.9 | 3.8 |
| Cancellation of elective procedures and admissions | 73.1 | 4.2 |
| Establishment of an alternate care site | 66.8 | 3.0 |
| Medical utilization of nonclinical space | 60.8 | 3.7 |
| Coordinated supply-chain management | 60.1 | 3.8 |
| Memoranda of understanding with hospitals to accept inpatients | 46.1 | 4.1 |
| Stockpiling antibiotics and supplies | 44.1 | 3.5 |
| Conversion of post-anesthesia unit to intensive care unit | 36.6 | 3.6 |
| Activation of decommissioned wards | 27.3 | 2.7 |

Table 5. Percent distribution (with standard errors) of hospitals involved in interagency task forces, designated by the National Disaster Medical System, and trained in incident command: United States, 2003

| Hospital characteristic | Percent distribution | Standard error |
|--|----------------------|----------------|
| All hospitals | 100.0 | ... |
| Interagency task force | | |
| Member | 87.4 | 4.0 |
| Not a member | *8.0 | 3.7 |
| Unknown | *4.6 | 1.9 |
| National Disaster Medical System | | |
| Designated to receive patients | 34.3 | 2.9 |
| Not designated | 32.2 | 4.1 |
| Unknown | 33.5 | 3.3 |
| Key personnel in incident command system | | |
| Trained | 75.0 | 3.8 |
| Not trained | 20.8 | 3.7 |
| Unknown | *4.2 | 1.6 |

... Category not applicable.

* Figure does not meet standards of reliability or precision.

NOTE: Numbers may not add to totals because of rounding.

Table 6. Percentage (with standard error) of hospitals whose staff have received training, by disease or exposure: United States, 2003

| Disease or exposure | Percent of hospitals | Standard error |
|--|----------------------|----------------|
| Smallpox | 86.7 | 2.4 |
| Anthrax | 83.7 | 3.1 |
| Chemical exposure | 80.2 | 2.7 |
| Nuclear or radiologic exposure | 66.4 | 3.3 |
| Botulism | 65.0 | 3.7 |
| Plague | 61.0 | 4.0 |
| Viral encephalitis | 53.2 | 3.7 |
| Tularemia | 51.3 | 3.9 |
| Viral hemorrhagic fever | 51.2 | 3.4 |

NOTE: Staff members include staff physicians, interns or residents, physician assistants or nurse practitioners, nurses, laboratory staff, and others.

Table 7. Percentage (with standard errors) of hospitals that trained their staff in any terrorism-related disease or condition, by type of profession: United States, 2003

| Type of clinician | Percent of hospitals | Standard error |
|---|----------------------|----------------|
| Nurses | 92.1 | 2.2 |
| Staff physicians | 82.5 | 3.0 |
| Laboratory staff | 72.9 | 3.8 |
| Physician assistants or nurse practitioners | 53.1 | 5.2 |
| Residents or interns | 49.2 | 5.6 |

NOTE: Terrorism-related diseases or conditions include smallpox, anthrax, plague, botulism, tularemia, viral hemorrhagic fever, viral encephalitis, and chemical, nuclear, and radiologic exposures.

Table 8. Percentage (with standard errors) of hospitals that used selected terrorism-preparedness training sources, by training source: United States, 2003

| Training source | Percent of hospitals | Standard error |
|---|----------------------|----------------|
| Hospital | 61.4 | 3.3 |
| State or local public health department | 44.8 | 3.7 |
| Professional association | 33.9 | 3.8 |
| Other State or local government agency | 24.5 | 3.5 |
| Federal agency | 10.1 | 2.2 |
| Private vendor | 8.4 | 1.7 |
| Insurance organization | *0.1 | 0.1 |
| Other | 19.3 | 3.8 |

* Figure does not meet standards of reliability or precision.

Table 9. Percentage (with standard errors) of hospitals participating in internal and external mass casualty drills, by type of drill scenario: United States, 2003

| Scenario | Internal drills | | External drills | |
|--|----------------------|----------------|----------------------|----------------|
| | Percent of hospitals | Standard error | Percent of hospitals | Standard error |
| Hospitals staging drills | 88.4 | 2.7 | 87.1 | 2.8 |
| General disaster or emergency response | 69.7 | 3.7 | 66.4 | 4.0 |
| Chemical release | 44.9 | 3.9 | 44.1 | 3.5 |
| Biologic attack | 37.5 | 3.5 | 33.9 | 3.2 |
| Explosive or incendiary attack | 21.3 | 3.2 | 24.2 | 3.4 |
| Nuclear or radiologic attack | 15.4 | 2.1 | 15.2 | 2.2 |
| Severe epidemic | 7.1 | 1.4 | 4.8 | 0.9 |

Table 10. Percentage (with standard errors) of hospitals collaborating in external mass casualty drills, by type of collaborating organization: United States, 2003

| Type of collaborating organization | Percent of hospitals | Standard error |
|--|----------------------|----------------|
| Emergency medical services | 70.9 | 4.0 |
| Fire department | 67.1 | 3.9 |
| State or local law enforcement | 64.5 | 3.7 |
| Other State or local government agencies | 52.0 | 4.0 |
| Other hospitals or hospital associations | 45.9 | 3.6 |
| State or local public health department | 45.6 | 4.0 |
| Hazardous materials teams | 43.8 | 3.5 |
| American Red Cross | 21.3 | 2.4 |
| Other volunteer organizations | 15.6 | 1.7 |
| Key vendors of medical supplies | *7.0 | 3.0 |
| Other | 17.5 | 3.9 |

* Figure does not meet standards of reliability or precision.

Table 11. Mean number and rate (with standard errors) of selected specialized equipment per responding hospital: United States, 2003

| Type of equipment | Mean | Standard error | Mean per 100 hospital beds | Standard error |
|-------------------------------------|------|----------------|----------------------------|----------------|
| Intensive care beds | 20.3 | 1.4 | 13.8 | 0.9 |
| Personal protective suits | 12.7 | 1.5 | 9.4 | 1.3 |
| Ventilators | 11.0 | 0.8 | 7.8 | 0.9 |
| Isolation rooms | 6.0 | 0.4 | 4.3 | 0.3 |
| Decontamination showers | 1.5 | 0.1 | 1.5 | 0.2 |

Technical Notes

Form Approved OMB No. 0920-0278 Exp. Date 04/30/2005

FORM **NHAMCS-905**
(4-21-2003)

U.S. DEPARTMENT OF COMMERCE
Economics and Statistics Administration
U.S. CENSUS BUREAU
ACTING AS DATA COLLECTION AGENT FOR THE
U.S. Department of Health and Human Services
Centers for Disease Control and Prevention
National Center for Health Statistics

NOTICE - Public reporting burden of this collection of information is estimated to average 25 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDA/ATSDR Reports Clearance Officer, 1600 Clifton Road, MS D-24, Atlanta, GA 30333, ATTN: PRA (0920-0278).

**NATIONAL HOSPITAL AMBULATORY
MEDICAL CARE SURVEY
BIOTERRORISM AND MASS CASUALTY
PREPAREDNESS SUPPLEMENT
2003 PANEL**

Assurance of confidentiality - All information which would permit identification of an individual, a practice, or an establishment will be held confidential, will be used only by persons engaged in and for the purpose of the survey and will not be disclosed or released to other persons or used for any other purpose without consent of the individual or the establishment in accordance with section 308(d) of the Public Health Service Act (42 USC 242m).

BACKGROUND INFORMATION

| | | | |
|---------------------------------|--|--------------------------------------|--------------------|
| A. Hospital name | | B. Hospital number | |
| C. Hospital contact name | | D. Hospital contact telephone | Area code Number |
| E. Census contact name | | F. Census contact telephone | Area code Number |

This year we are conducting a special survey supplement on bioterrorism and mass casualty preparedness in hospitals. Please answer the following questions. We appreciate your time on this important public health concern.

| | | | |
|---|---|---|---|
| 1. Has your hospital's emergency/bioterror response plan been revised since September 11, 2001? | | <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> <i>Mark (X) this box if hospital has no emergency/bioterror response plan and SKIP to item 5.</i> | |
| 2. Does your emergency/bioterror response plan specifically address each of the following types of incidents? If "No" - Indicate whether or not your hospital is currently developing a response for the type of incident. Mark (X) one box for each type of incident. | Does your plan address this type of incident? | Are you currently developing a response for the incident? | |
| | | (1) | (2) |
| a. Biological | <input type="checkbox"/> Yes <input type="checkbox"/> No → | <input type="checkbox"/> Yes <input type="checkbox"/> No | <input type="checkbox"/> Yes <input type="checkbox"/> No |
| b. Chemical | <input type="checkbox"/> Yes <input type="checkbox"/> No → | <input type="checkbox"/> Yes <input type="checkbox"/> No | <input type="checkbox"/> Yes <input type="checkbox"/> No |
| c. Nuclear/Radiologic | <input type="checkbox"/> Yes <input type="checkbox"/> No → | <input type="checkbox"/> Yes <input type="checkbox"/> No | <input type="checkbox"/> Yes <input type="checkbox"/> No |
| d. Explosive/Incendiary | <input type="checkbox"/> Yes <input type="checkbox"/> No → | <input type="checkbox"/> Yes <input type="checkbox"/> No | <input type="checkbox"/> Yes <input type="checkbox"/> No |
| e. Natural disaster | <input type="checkbox"/> Yes <input type="checkbox"/> No → | <input type="checkbox"/> Yes <input type="checkbox"/> No | <input type="checkbox"/> Yes <input type="checkbox"/> No |
| f. Other - Please specify | <input type="checkbox"/> Yes <input type="checkbox"/> No → | <input type="checkbox"/> Yes <input type="checkbox"/> No | <input type="checkbox"/> Yes <input type="checkbox"/> No |

PLEASE CONTINUE WITH QUESTION 3 ON NEXT PAGE

7. Have key personnel in your hospital been trained in how to implement a formal incident command system (e.g., HEICS [Hospital Emergency Incident Command System] or comparable platform) during emergencies?

- 1 Yes
- 2 No
- 3 Don't know

8. TRAINING

a. Have your hospital staff received special training (e.g., in-service or other courses, CME, Grand Rounds, or self-guided study) since September 11, 2001 in the identification, diagnosis, and treatment of the following diseases/conditions?

Type of personnel who received training
Mark (X) appropriate columns OR mark (X) N/A box, if your hospital does not have this type of personnel.

| | <input type="checkbox"/> N/A | <input type="checkbox"/> N/A | <input type="checkbox"/> N/A | <input type="checkbox"/> N/A | <input type="checkbox"/> N/A | <input type="checkbox"/> N/A |
|---|------------------------------|--|------------------------------|------------------------------|------------------------------|------------------------------|
| | Staff physicians (a) | House officers (intern/resident) (b) | PA/NP (c) | RN/LPN (d) | Lab staff (e) | Other (f) |
| (1) Smallpox | | | | | | |
| (2) Anthrax | | | | | | |
| (3) Plague | | | | | | |
| (4) Botulism | | | | | | |
| (5) Tularemia | | | | | | |
| (6) Viral Hemorrhagic Fever | | | | | | |
| (7) Viral Encephalitis (WNV, SLE, EEE, VEE, etc.) | | | | | | |
| (8) Chemical exposure | | | | | | |
| (9) Nuclear/Radiologic exposure | | | | | | |

NOTE If no training received – Please SKIP to item 9. Otherwise, continue with item b.

b. Indicate who conducted the training.

Mark(X) all that apply.

- 1 Professional association (e.g., medical, physician assistant, nursing, laboratory)
- 2 State or local public health department
- 3 Other state or local government agency
- 4 Federal agency
- 5 Hospital
- 6 Insurance organization
- 7 Private vendor
- 8 Other –Specify

PLEASE CONTINUE WITH QUESTION 9 ON NEXT PAGE

Suggested citation

Niska RW, Burt CW. Bioterrorism and mass casualty preparedness in hospitals: United States, 2003. Advance data from vital and health statistics; no 364. Hyattsville, MD: National Center for Health Statistics. 2005.

Copyright information

All material appearing in this report is in the public domain and may be reproduced or copied without permission; citation as to source, however, is appreciated.

National Center for Health Statistics

Director
Edward J. Sondik, Ph.D.
Acting Co-Deputy Directors
Jennifer H. Madans, Ph.D.
Michael H. Sadagursky

U.S. DEPARTMENT OF
HEALTH & HUMAN SERVICES

Centers for Disease Control and Prevention
National Center for Health Statistics
3311 Toledo Road
Hyattsville, Maryland 20782

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE, \$300

To receive this publication regularly, contact
the National Center for Health Statistics by
calling 1-866-441-NCHS (6247)
E-mail: nchsquery@cdc.gov
Internet: www.cdc.gov/nchs

DHHS Publication No. (PHS) 2005-1250
05-0429 (8/05)

| |
|--|
| FIRST CLASS POSTAGE & FEES PAID CDC/NCHS PERMIT NO. G-284 |
|--|