

Attachment J

- (1) NCI. Risks Associated with Smoking Cigarettes with Low Machine-Measured Yields of Tar and Nicotine. Smoking and Tobacco Monograph No. 13, NIH Pub. No. 02-5074. 2001. Bethesda, MD, U.S. Department of Health and Human Services, National Institutes of Health, National Cancer Institute.
Ref Type: Report
- (2) Djordjevic MV, Stellman SD, Zang E. Doses of nicotine and lung carcinogens delivered to cigarette smokers. *J Natl Cancer Inst.* 2000;92:106-111.
- (3) Bridges RB, Combs JG, Humble JW, Turbek JA, Rehm SR, Haley NJ. Puffing topography as a determinant of smoke exposure. *Pharmacol Biochem Behav.* 1990;37:29-39.
- (4) Kolonen S, Tuomisto J, Puustinen P, Airaksinen MM. Effects of smoking abstinence and chain-smoking on puffing topography and diurnal nicotine exposure. *Pharmacol Biochem Behav.* 1992;42:327-332.
- (5) Federal Trade Commission. Cigarette Report for 2001. 2003.
Ref Type: Report
- (6) Djordjevic MV, Fan J, Ferguson S, Hoffmann D. Self-regulation of smoking intensity. Smoke yields of the low-nicotine, low-'tar' cigarettes. *Carcinogenesis.* 1995;16:2015-2021.
- (7) Fischer S, Spiegelhalter B, Preussmann R. Influence of smoking parameters on the delivery of tobacco-specific nitrosamines in cigarette smoke--a contribution to relative risk evaluation. *Carcinogenesis.* 1989;10:1059-1066.
- (8) Bernert, J. T. Biomarkers of environmental tobacco smoke exposure. 2004.
Ref Type: Slide
- (9) Russell MA, Jarvis M, Iyer R, Feyerabend C. Relation of nicotine yield of cigarettes to blood nicotine concentrations in smokers. *Br Med J.* 1980;280:972-976.
- (10) National Center for Chronic Disease Prevention and Health Promotion. Cigarette Smoking-Related Mortality. Tobacco Information and Prevention Source (TIPS) . 2006.
Ref Type: Electronic Citation
- (11) Mendelson JH, Sholar MB, Goletiani N, Siegel AJ, Mello NK. Effects of low- and high-nicotine cigarette smoking on mood states and the HPA axis in men. *Neuropsychopharmacology.* 2005;30:1751-1763.
- (12) Pickworth WB, Moolchan ET, Berlin I, Murty R. Sensory and physiologic effects of menthol and non-menthol cigarettes with differing nicotine delivery. *Pharmacol Biochem Behav.* 2002;71:55-61.

- (13) Lee EM, Malson JL, Moolchan ET, Pickworth WB. Quantitative comparisons between a nicotine delivery device (Eclipse) and conventional cigarette smoking. *Nicotine Tob Res.* 2004;6:95-102.
- (14) Pickworth W.B., Houlgate, P., Schorp, M., Dixon, M., Borgerding, M. F., and Zaatari, G. A review of human smoking behavior data and recommendations for a new ISO standard for the machine smoking of cigarettes. Report of the *AD HOC* WG9 Smoking Behaviour Review Team to ISO/TC 126 WG9. 8-10-2005.
Ref Type: Report
- (15) Surgeon General. Tobacco use as drug dependence. Centers for Disease Control and Prevention. Chapter 4, 145-239. 1988.
Ref Type: Report
- (16) Caraballo RS, Giovino GA, Pechacek TF et al. Racial and ethnic differences in serum cotinine levels of cigarette smokers: Third National Health and Nutrition Examination Survey, 1988-1991. *JAMA.* 1998;280:135-139.
- (17) US Department of Commerce, Census Bureau. National Cancer Institute Sponsored Tobacco Use Supplement to the Current Population Survey (2001-2002). Data files. 2004.
Ref Type: Report
- (18) Cohen J. *Statistical power analyses for the behavioral sciences.* 2nd ed. New York: Erlbaum; 1988.
- (19) Ahijevych K, Parsley LA. Smoke constituent exposure and stage of change in black and white women cigarette smokers. *Addict Behav.* 1999;24:115-120.
- (20) Clark PI, Gautam S, Gerson LW. Effect of menthol cigarettes on biochemical markers of smoke exposure among black and white smokers. *Chest.* 1996;110:1194-1198.
- (21) Eissenberg T, Adams C, Riggins EC, III, Likness M. Smokers' sex and the effects of tobacco cigarettes: subject-rated and physiological measures. *Nicotine Tob Res.* 1999;1:317-324.
- (22) Perez-Stable EJ, Herrera B, Jacob P, III, Benowitz NL. Nicotine metabolism and intake in black and white smokers. *JAMA.* 1998;280:152-156.
- (23) Tabachnick BG, Fidell LS. *Using multivariate statistics.* 4th ed. Boston: Allyn and Bacon; 2001.
- (24) Hecht SS, Carmella SG, Chen M et al. Quantitation of urinary metabolites of a tobacco-specific lung carcinogen after smoking cessation. *Cancer Res.* 1999;59:590-596.
- (25) Benowitz NL, Jacob IP. Individual differences in Nicotine kinetics and metabolism in humans. *Pharmacokinetics, metabolism, and pharmaceuticals of drugs of abuse.* NIDA Research Monograph 173, NIH Publication No. 97-4141 ed. Bethesda, MD: U.S.

Department of Health and Human Services, National Institutes of Health, National Institute on Drug Abuse; 1997:48-64.

- (26) Vivometrics. Vivometrics LifeShirt® system receives U.S. Food and Drug Administration 510(K) market clearance.
http://www.vivometrics.com/site/press_pr20020422.html Press Release. 4-22-2002.
Ref Type: Generic
- (27) Paakko P, Kokkonen P, Anttila S, Kalliomaki PL. Cadmium and chromium as markers of smoking in human lung tissue. *Environ Res.* 1989;49:197-207.
- (28) Navas-Acien A, Selvin E, Sharrett AR, Calderon-Aranda E, Silbergeld E, Guallar E. Lead, cadmium, smoking, and increased risk of peripheral arterial disease. *Circulation.* 2004;109:3196-3201.
- (29) Friberg L, Piscator M, Nordberg GF, Kjellstrom T. *Cadmium in environment.* 2nd ed. Boca Raton, FL.: CRC Press; 1974.
- (30) Institute of Medicine. *Clearing the smoke: Assessing the science base for tobacco harm reduction.* Washington DC: National Academy Press; 2001.
- (31) Murphy SE, Link CA, Jensen J et al. A comparison of urinary biomarkers of tobacco and carcinogen exposure in smokers. *Cancer Epidemiol Biomarkers Prev.* 2004;13:1617-1623.
- (32) Benowitz NL, Jacob P, III. Metabolism of nicotine to cotinine studied by a dual stable isotope method. *Clin Pharmacol Ther.* 1994;56:483-493.
- (33) Bernert JT, Jr., McGuffey JE, Morrison MA, Pirkle JL. Comparison of serum and salivary cotinine measurements by a sensitive high-performance liquid chromatography-tandem mass spectrometry method as an indicator of exposure to tobacco smoke among smokers and nonsmokers. *J Anal Toxicol.* 2000;24:333-339.
- (34) Benowitz NL, Jacob P, III. Trans-3'-hydroxycotinine: disposition kinetics, effects and plasma levels during cigarette smoking. *Br J Clin Pharmacol.* 2001;51:53-59.
- (35) Benowitz NL, Jacob P, III. Trans-3'-hydroxycotinine: disposition kinetics, effects and plasma levels during cigarette smoking. *Br J Clin Pharmacol.* 2001;51:53-59.
- (36) Hecht SS. Biochemistry, biology, and carcinogenicity of tobacco-specific N-nitrosamines. *Chem Res Toxicol.* 1998;11:559-603.
- (37) Hecht SS, Hoffman D. N-nitroso compounds and man: sources of exposure, endogenous formation and occurrence in body fluids. *Eur J Cancer Prev.* 1998;7:244-246.
- (38) Hecht SS. Human urinary carcinogen metabolites: biomarkers for investigating tobacco and cancer. *Carcinogenesis.* 2002;23:907-922.

- (39) Hecht SS. Tobacco smoke carcinogens and lung cancer. *J Natl Cancer Inst.* 1999;91:1194-1210.
- (40) Hoffmann D, Hoffmann I, El Bayoumy K. The less harmful cigarette: a controversial issue. a tribute to Ernst L. Wynder. *Chem Res Toxicol.* 2001;14:767-790.
- (41) Carmella SG, Akerkar S, Hecht SS. Metabolites of the tobacco-specific nitrosamine 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in smokers' urine. *Cancer Res.* 1993;53:721-724.
- (42) Carmella SG, Akerkar SA, Richie JP, Jr., Hecht SS. Intraindividual and interindividual differences in metabolites of the tobacco-specific lung carcinogen 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) in smokers' urine. *Cancer Epidemiol Biomarkers Prev.* 1995;4:635-642.
- (43) Karreth S, Lenk W. The metabolism of 4-aminobiphenyl in rat. I. Reaction of N-hydroxy-4-aminobiphenyl with rat blood in vivo. *Xenobiotica.* 1991;21:417-428.
- (44) Linhart I, Frantik E, Vodickova L, Vosmanska M, Smejkal J, Mitera J. Biotransformation of acrolein in rat: excretion of mercapturic acids after inhalation and intraperitoneal injection. *Toxicol Appl Pharmacol.* 1996;136:155-160.
- (45) U.S.Environmental Protection Agency. Toxicological review of acrolein (CAS No. 107-02-8): In Support of Summary Information on the Integrated Risk Information System (IRIS). EPA/635/R-03/003. 2003. Washington, D.C.
Ref Type: Report
- (46) Agency for Toxic Substances and Disease Registry (ATSDR). Toxicological profile for 1,3 butadiene. Public Health Service. 1992. Atlanta, GA, US Department of Health and Human Services.
Ref Type: Generic
- (47) U.S.Environmental Protection Agency. Health Assessment of 1,3-Butadiene. EPA/600/P-98/001F. 2002. Washington, DC, National Center for Environmental Assessment.
Ref Type: Report
- (48) Melikian AA, Qu Q, Shore R et al. Personal exposure to different levels of benzene and its relationships to the urinary metabolites S-phenylmercapturic acid and trans,trans-muconic acid. *J Chromatogr B Analyt Technol Biomed Life Sci.* 2002;778:211-221.
- (49) Qu Q, Cohen BS, Shore R et al. Benzene exposure measurement in shoe and glue manufacturing: a study to validate biomarkers. *Appl Occup Environ Hyg.* 2003;18:988-998.
- (50) Qu Q, Shore R, Li G et al. Validation and evaluation of biomarkers in workers exposed to benzene in China. *Res Rep Health Eff Inst.* 2003;1-72.

- (51) Qu Q, Melikian AA, Li G et al. Validation of biomarkers in humans exposed to benzene: urine metabolites. *Am J Ind Med.* 2000;37:522-531.
- (52) Bayliss, D., Chen, C., Sonawane, B., and Valcovic, L. Carcinogenic effects of benzene: An update. EPA/600/P-97/001A. 1997. Washington, DC, U.S. Environmental Protection Agency, National Center for Environmental Assessment.
Ref Type: Report
- (53) Brzeznicki S, Jakubowski M, Czerski B. Elimination of 1-hydroxypyrene after human volunteer exposure to polycyclic aromatic hydrocarbons. *Int Arch Occup Environ Health.* 1997;70:257-260.
- (54) Jongeneelen FJ, Bos RP. Excretion of pyrene and hydroxypyrene in urine. *Cancer Lett.* 1990;51:175-179.
- (55) Buckley TJ, Liroy PJ. An examination of the time course from human dietary exposure to polycyclic aromatic hydrocarbons to urinary elimination of 1-hydroxypyrene. *Br J Ind Med.* 1992;49:113-124.
- (56) van Schooten FJ, Jongeneelen FJ, Hillebrand MJ et al. Polycyclic aromatic hydrocarbon-DNA adducts in white blood cell DNA and 1-hydroxypyrene in the urine from aluminum workers: relation with job category and synergistic effect of smoking. *Cancer Epidemiol Biomarkers Prev.* 1995;4:69-77.
- (57) Watson C, McCraw J, Polzin G, Ashley D. Development of a method to assess cigarette smoke intake. *Environ Sci Technol.* 2004;38:248-253.
- (58) Jenkins RA, Guerin MT, Tomkins BA. *The chemistry of environmental tobacco smoke: Composition and measurement.* Boca Raton, FL: CRC Press LLC; 2000.