

Rheutopia [Roo-TOH-pee-ah] (also known as arixalate)

Uses

- Rheumatoid arthritis in adults. Rheutopia reduces painful and swollen joints, slows joint damage, and improves mobility and the ability to do physical activities.
- Polyarticular juvenile rheumatoid arthritis in children at least 4 years old who did not have good results from other medicines. Rheutopia reduces pain, improves mobility, and decreases the number of painful joints.
- Ankylosing spondylitis. Rheutopia reduces back pain, swelling, and improves mobility.
- Plaque psoriasis in adults who may benefit from taking medicine or receiving phototherapy (using ultraviolet light). Rheutopia improves or clears up areas of skin with psoriasis.

Call Your Doctor

Stop using Rheutopia and tell your doctor right away if you develop:

- Fever, cough, flu-like symptoms, skin infection (red, warm, painful skin or open sores).
- Numbness, tingling, weakness, vision problems, or dizziness.
- Chills, swollen lymph nodes, night sweats, fever, or weight loss.
- Bruising, bleeding, and pale skin.
- Shortness of breath, swelling of ankles or feet, or sudden weight gain.
- Chest discomfort or pain, shortness of breath, joint pain or a rash on your cheeks or arms.

Important Warning: Serious Infections

- Rheutopia affects the immune system. It can lower your ability to fight infections. Do not use Rheutopia if you have an active infection.
- People taking Rheutopia have gotten serious infections including tuberculosis (TB) and infections caused by viruses, fungi, or bacteria. Some people have died from these infections.

Common Side Effects

- Redness, rash, swelling, itching or bruising where the shot was given.
- Headache
- Runny nose

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Tell Your Doctor

Before using Rheutopia, tell your doctor if you:

- have an infection, are being treated for an infection, or think you have an infection (such as a cold, flu or skin infection).
- have TB or have been near someone who has TB.
- have any nervous system or heart problems.
- have lived in or traveled to other countries.
- have been recently been vaccinated or are scheduled to receive a vaccination (including a flu shot).
- are taking the medicine Kineret (anakinra).

Directions for Use

- Rheutopia is an injection (shot). Do not use Rheutopia until your doctor has shown you how to give a shot.
- Store Rheutopia in the refrigerator. Do not shake or freeze.
- If you forget to take a dose, take it as soon as you remember. Take your next dose at your regularly scheduled time.
- Your doctor will tell you how often to use Rheutopia. Do not use Rheutopia more often than prescribed.

Rheutopia [Roo-TOH-pee-ah] (also known as arixalate)

Uses

- Rheumatoid arthritis in adults. Rheutopia reduces painful and swollen joints, slows joint damage, and improves mobility and the ability to do physical activities.
- Polyarticular juvenile rheumatoid arthritis in children at least 4 years old who did not have good results from other medicines. Rheutopia reduces pain, improves mobility, and decreases the number of painful joints.
- Ankylosing spondylitis. Rheutopia reduces back pain, swelling, and improves mobility.
- Plaque psoriasis in adults who may benefit from taking medicine or receiving phototherapy (using ultraviolet light). Rheutopia improves or clears up areas of skin with psoriasis.

Important Warning: Serious Infections

- Rheutopia affects the immune system. It can lower your ability to fight infections. Do not use Rheutopia if you have an active infection.
- People taking Rheutopia have gotten serious infections including tuberculosis (TB) and infections caused by viruses, fungi, or bacteria. Some people have died from these infections.

Tell Your Doctor

Before using Rheutopia, tell your doctor if you:

- have an infection, are being treated for an infection, or think you have an infection (such as a cold, flu or skin infection).
- have TB or have been near someone who has TB.
- have any nervous system or heart problems.
- have lived in or traveled to other countries.
- have been recently been vaccinated or are scheduled to receive a vaccination (including a flu shot).
- are taking the medicine Kineret (anakinra).

Call Your Doctor

Stop using Rheutopia and tell your doctor right away if you develop:

- Fever, cough, flu-like symptoms, skin infection (red, warm, painful skin or open sores).
- Numbness, tingling, weakness, vision problems, or dizziness.
- Chills, swollen lymph nodes, night sweats, fever, or weight loss.
- Bruising, bleeding, and pale skin.
- Shortness of breath, swelling of ankles or feet, or sudden weight gain.
- Chest discomfort or pain, shortness of breath, joint pain or a rash on your cheeks or arms.

Common Side Effects

- Redness, rash, swelling, itching or bruising where the shot was given.
- Headache
- Runny nose

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Directions for Use

- Rheutopia is an injection (shot). Do not use Rheutopia until your doctor has shown you how to give a shot.
- Store Rheutopia in the refrigerator. Do not shake or freeze.
- If you forget to take a dose, take it as soon as you remember. Take your next dose at your regularly scheduled time.
- Your doctor will tell you how often to use Rheutopia. Do not use Rheutopia more often than prescribed.