

Supporting Statement B for

NIH Toolbox for Assessment of
Neurological and Behavioral Function (NIA)

April 11, 2011

Molly Wagster, Ph.D.,
Contracting Officer Technical
Representative
NIH Toolbox
National Institute on Aging

Gateway Building, 350
7201 Wisconsin Ave
Bethesda, MD 20892

Telephone: 301-496-9350
Fax: 301-496-1494
E-mail: wagsterm@nia.nih.gov

TABLE OF CONTENTS

B.	COLLECTION OF INFORMATION EMPLOYING STATISTICAL METHODS...	6
B.1	RESPONDENT UNIVERSE AND SAMPLING METHODS.....	6
B.2	PROCEDURES FOR THE COLLECTION OF INFORMATION.....	8
B.3	METHODS TO MAXIMIZE RESPONSE RATES AND DEAL WITH NONRESPONSE.....	10
B.4	TEST OF PROCEDURES OR METHODS TO BE UNDERTAKEN.....	11
B.5	INDIVIDUALS CONSULTED ON STATISTICAL ASPECTS AND INDIVIDUALS COLLECTING AND/OR ANALYZING DATA.....	12

List of Attachments

BATTERY OF TOOLBOX MEASURES

Sensation domain

- 1 Automated Audiometry from the NIH Toolbox (AANT)
- 2 Hearing Handicap Inventory
- 3 Words-in-Noise (WIN)
- 4 Odor Identification Ages 10-85
- 5 Odor Identification Ages 3-9
- 6 University of Pennsylvania Smell Identification Test (UPSIT)
- 7 Pain Assessment Proxy Ages 3-7
- 8 Pain Assessment Ages 8-12
- 9 Pain Assessment Ages 13-17
- 10 Pain Assessment Ages 18-85
- 11 Sucrose Preference Test
- 12 Toolbox Taste Test
- 13 Dynamic Visual Acuity (DVA)
- 14 Vision Health-Related Quality of Life Survey
- 15 Brief Kinesthesia
- 16 Pediatric Pain Scale
- 17 Tactile Discrimination Test

Cognition domain

- 18 Dimensional Change Card Sort
- 19 Bateria III Picture Vocabulary
- 20 Flanker
- 21 Imitation Based Assessment of Memory (IBAM)
- 22 List Sorting
- 23 Oral Symbol Digit Modalities Test
- 24 Pattern Comparison
- 25 Rey Auditory Verbal Learning Test (RAVLT)
- 26 Reading Recognition
- 27 Determining Handedness
- 28 Vocabulary Comprehension
- 29 Spanish Word Accentuation Test

Emotion domain

- 30 Instructions for Emotional Health Questionnaire
- 31 Emotional Health Questionnaires for Toolbox
- 32 Supplemental Emotional Health Questionnaires

Motor functioning domain

- 33 9 Hole Pegboard
- 34 Endurance 2 Minute Walk Test (2MWT)
- 35 Locomotion 4 Meter Walk Test
- 36 Grip Strength Dynamometry
- 37 Strength HHD Knee Extension
- 38 Balance Accelerometer Measure (BAM)

ADDITIONAL MEASURES

- 39 Initial Questionnaire - ADULT
- 40 Initial Questionnaire - PROXY and CHILD
- 41 Additional Somatosensation Questions
- 42 Additional Audition Olfaction Taste Questions
- 43 Cognition Information Form
- 44 Health Care Access and Utilization
- 45 Instrumental Activities of Daily Living (IADL) - Adults
- 46 Applied Cognition - Adult
- 47 Pediatric Perceived Cognitive Function (PCF) - Child
- 48 Pediatric Perceived Cognitive Function (PCF) - Proxy
- 49 PROMIS Pediatric Physical Function Upper Extremity
- 50 PROMIS Pediatric Physical Function Mobility
- 51 Pediatric Functional Assessment of Chronic Illness Therapy – Fatigue (FACIT-F)
- 52 PROMIS 29 - Participant
- 53 School Performance and Activities
- 54 Sociodemographic Form – Parent
- 55 Test Anxiety Scale (TAS)
- 56 Falls Efficacy Scale - International (Short FES-I)
- 57 EtOH Questionnaire
- 58 Education Adult Self-Report
- 59 Education Proxy Report for Children

RECRUITMENT MATERIALS

- 60 Recruitment Instructions, Scripts and Letters

REGULATORY DOCUMENTS

- 61 Certificate of Confidentiality
- 62 Northwestern University IRB Approval Letter
- 63 Consent Documents

OTHER DOCUMENTS

- 64 Team Member Contact List
- 65 NIH Privacy Act Review Memo
- 66 NIH Privacy Act Systems of Record 09-25-0200

B. COLLECTION OF INFORMATION EMPLOYING STATISTICAL METHODS

B.1 Respondent Universe and Sampling Methods

The instruments in the NIH Toolbox are designed primarily for use in a **Target Population** with the following characteristics:

1. General non-institutionalized population dwelling in the community
2. Ages 3-85 years
3. Availability of a proxy respondent for pediatric population
4. Capable of following instructions given in English or Spanish and understanding the informed consent process
5. Adequate visual, auditory, vestibular, or motor functioning or availability of assistance or assist devices is required to complete some of the items in the test battery

This target population is estimated to include approximately all 283,764,466 civilian noninstitutionalized individuals ages 3-85 years in the United States. A total of 8,400 individuals and 4,500 parent-proxies will be sampled from this population. Distribution of the sample and the respondent universe across the strata are presented in Table B.1.1. The accrual vendor's existing recruitment database of 492,008 subjects will be randomly sampled and screened for household members' age, gender, race/ethnicity, education and primary language. We expect that approximately 15% of the eligible individuals will agree to participate and 85% of the selected participants will actually complete the tests successfully.

Table B.1.1. Distribution of Respondent Universe And Proposed Sample Across Strata

	Universe¹	Sample
Total	283,764,466	8400
Males	139,361,273	4200
Females	144,403,193	4200
English-speaking ²	232,686,862	5850
Spanish-speaking ²	31,214,091	2550
Age 3	4,168,945	300
Age 4	4,122,392	300
Age 5	4,133,901	300
Age 6	4,072,774	300
Age 7	4,060,824	300
Age 8	4,104,423	300
Age 9	3,921,722	300
Age 10	3,922,622	300
Age 11	3,908,905	300
Age 12	3,946,112	300
Age 13	4,032,341	300
Age 14	4,071,443	300
Age 15	4,106,574	300
Age 16	4,191,736	300
Age 17	4,250,817	300
Age 18-29	49,965,595	600
Age 30-39	39,581,409	600
Age 40-49	43,459,875	600
Age 50-59	40,494,881	600
Age 60-69	27,275,851	600
Age 70-85	21,971,324	900

¹ US Census Bureau Civilian Noninstitutionalized Population estimates for 5/1/2009

² Estimated using percentages reported in Census 2000 Brief: Language Use and English-Speaking Ability: 2000

The NIH Toolbox sample frame (see tables below) will be drawn from the Delve sample. The sample frame was derived from the 2010 Census (respective to each age group). The Delve database is updated in a process involving continuous recruitment. The Delve sample has been evaluated over the last two weeks and has been confirmed to meet the needs of our sample frame (including oversampling required to insure final completes) without further revision.. No revisions to the sample or frame are predicted given the very short accrual period to follow (<3 months).

Adults:

* Race / ethnicity quotas apply to English-speaking group only, it is expected that the Spanish-speaking group will be almost entirely Hispanic. H=Hispanic, B=Black, A=Asian, O=Other, includes American Indian, Alaska Native, Native Hawaiian or Other Pacific Islander and individuals with 2 or more races.

Age		Race / Ethnicity*	Educational Attainment
18 - 29		19% H, 15% B, 4% A, 3% O	16% Less, 29% HS, 55% More
30 - 39		19% H, 14% B, 6% A, 3% O	12% Less, 27% HS, 61% More
40 - 49		14% H, 13% B, 5% A, 2% O	11% Less, 31% HS, 58% More
50 - 59		10% H, 12% B, 4% A, 2% O	11% Less, 31% HS, 58% More
60 - 69		8% H, 10% B, 4% A, 2% O	13% Less, 32% HS, 54% More
70 - 79			
80 - 85		7% H, 9% B, 4% A, 1% O	24% Less, 37% HS, 39% More
Total			

Children:

Age		Race / Ethnicity*	Highest Educated Parent's Educational Attainment
3		25% H, 15% B, 5% A, 5% O	16% Less, 29% HS, 55% More
4		25% H, 15% B, 5% A, 5% O	16% Less, 29% HS, 55% More
5		23% H, 14% B, 5% A, 5% O	12% Less, 27% HS, 61% More
6		23% H, 14% B, 5% A, 5% O	12% Less, 27% HS, 61% More
7		23% H, 14% B, 5% A, 5% O	12% Less, 27% HS, 61% More
8		23% H, 14% B, 5% A, 5% O	12% Less, 27% HS, 61% More
9		23% H, 14% B, 5% A, 5% O	12% Less, 27% HS, 61% More
10		20% H, 15% B, 4% A, 4% O	12% Less, 27% HS, 61% More
11		20% H, 15% B, 4% A, 4% O	12% Less, 27% HS, 61% More

12		20% H, 15% B, 4% A, 4% O	12% Less, 27% HS, 61% More
13		20% H, 15% B, 4% A, 4% O	12% Less, 27% HS, 61% More
14		20% H, 15% B, 4% A, 4% O	12% Less, 27% HS, 61% More
15		19% H, 16% B, 4% A, 4% O	11% Less, 31% HS, 58% More
16		19% H, 16% B, 4% A, 4% O	11% Less, 31% HS, 58% More
17		19% H, 16% B, 4% A, 4% O	11% Less, 31% HS, 58% More
Total			

B.2 Procedures for the Collection of Information

Existing recruitment databases will be randomly sampled and screened for household members' age, gender, race/ethnicity, education and primary language. This will be done by Delve, the marketing research firm which will conduct the data collection activities. The goal is to accrue 150 participants for each of the adult cells, defined by age, gender, and language; and 75 participants in each of the child cells, as shown in Tables B2.1 and B2.2. Since very few school age children speak Spanish as their dominant language, this sample will not include Spanish speakers age 8-17, but additional English speaking children will be recruited to maintain the overall sample size across ages. The single-year age increments in the children are designed to

capture the rapid developmental change that occurs during childhood, while the adult age groups were similarly designed to correspond with accepted periods of transition and decline. The ultimate goal of this project is to provide normative reference values that will be acceptable to the epidemiological and clinical research community. . The data collected in this study will provide needed normative values for sensory, motor, cognitive, and emotional function that are relevant and generalizable to community based samples of children, adolescents, adults and older adults of both genders, with varying educational levels, from urban and suburban locales, for both English or Spanish preferred language populations. Therefore, this sample size per cell was selected based on the precedent of other norming studies, in particular the Wechsler Adult Intelligence Scale. This sample size also provides sufficient precision in the estimates of within-cell normative values. In many instances precision will be improved by combining English and Spanish-speaking cells and/or combining males and females within an age group. For certain tests, such as the Reading and Vocabulary components of the Cognition domain, it will be important to provide normative values separately for English and Spanish-speakers, whereas for Motor domain measures this distinction is not necessary and the language groups may be combined for increased precision. Similarly, some instruments may warrant gender-specific normative values while other instruments will not. For the smallest cells of size 75, the 95% confidence interval for the cell mean will be estimated with a precision of ± 0.23 standard deviation units. Cells of size 150 will have precision of ± 0.16 , size of 300 corresponds to precision of ± 0.11 , size of 600 corresponds to precision of ± 0.08 , and size of 900 corresponds to precision of ± 0.06 . This provides us with precision of ± 0.20 or better for all but the smallest cell size. It is widely accepted in the minimally important difference literature on patient-reported outcomes that differences less than 0.20 standard deviation units (a small effect size according to

accepted criteria) are unlikely to be meaningful differences. The cells that are rarest in the U.S. population are the Spanish-speaking child and elderly cells.

Table B2.1. Target Completes for Children by Cell

Age	English		Spanish		Total
	Male	Female	Male	Female	
3	75	75	75	75	300
4	75	75	75	75	300
5	75	75	75	75	300
6	75	75	75	75	300
7	75	75	75	75	300
8	150	150	0	0	300
9	150	150	0	0	300
10	150	150	0	0	300
11	150	150	0	0	300
12	150	150	0	0	300
13	150	150	0	0	300
14	150	150	0	0	300
15	150	150	0	0	300
16	150	150	0	0	300
17	150	150	0	0	300
Total	1875	1875	375	375	4500

Table B2.2. Target Completes for Adults by Cell

Age	English		Spanish		Total
	Male	Female	Male	Female	
18-29	150	150	150	150	600
30-39	150	150	150	150	600
40-49	150	150	150	150	600
50-59	150	150	150	150	600
60-69	150	150	150	150	600
70-79	150	150	150	150	900
80-85	150	150			
Total	1050	1050	900	900	3,900

We plan to stratify the recruitment databases into strata corresponding to testing cells defined by age, sex, and language. We will select a sample of persons from those databases for participation in the study. We will select a conditionally equal probability sample of persons from each stratum. Interviewers will contact the households to recruit the selected persons into the study.

We expect to potentially recruit multiple persons from some households, particularly from larger households containing members of the rarer cells, although we are not explicitly controlling for the number of persons per household.

Participants will complete a small portion of the assessment on paper prior to their scheduled visit. Trained Delve technicians will then administer the Toolbox measures to adult participants, child participants, and proxies, as appropriate. For one-week test-retest reliability testing, 375 children (and their parents for proxy measures) and 375 adults will be randomly selected and asked to schedule another appointment at which the Toolbox measures will be re-administered. Another 375 children (and their parents for proxy measures) and 375 adults will be randomly selected to be contacted by telephone to schedule a 3-month follow-up appointment to evaluate change in the Toolbox measures.

B.3 Methods to Maximize Response Rates and Deal with Nonresponse

We have assumed that approximately 85% of the selected participants will actually complete the tests successfully. To achieve this response rate, in addition to offering geographically appropriate compensation, we will provide scheduled respondents with a detailed packet of information regarding what they can expect on the day of testing, directions to the testing site, and a telephone number to call if they have questions regarding the study (see Attachment 60). They will also receive a telephone call reminding them of their appointment 2-3 days in advance and offering them the opportunity to reschedule if the scheduled appointment is no longer convenient. Respondents who are unable to travel to the testing location due to the financial burden or unavailability of transportation will be provided compensation for travel. In addition, selected respondents who do not keep their testing appointment will receive a call offering them

the opportunity to reschedule. These techniques combined will serve to maximize respondent cooperation. The 15% that do not complete the study may drop out in a variety of ways—not completing the telephone questionnaire and confirming eligibility, not giving parental consent, not showing up for the tests, etc. The tests require the participants to go to a testing site for about three hours of tests, which is not conducive to a higher response rate, even with compensation. This response rate is consistent with the OMB minimum acceptable response rate, and furthermore, the design provides an adequate number of respondents in rare population cells to accurately norm the measures.

We will do a nonresponse bias analysis. The sample will be selected disproportionately to achieve the target numbers within each cell, and the cases for a cell will not be combined with cases from other cells (with the possible exception of Spanish-speaking elderly). Therefore, the nonresponse bias of interest is strictly within each cell. We will compare the weighted respondents to the total U.S. for the same age, gender, and language for a very limited set of demographics available, such as geographical region, household income, and household size.

B.4 Test of Procedures or Methods to be Undertaken

All of the measures administered as part of the assessment have undergone extensive review as part of their development and no additional testing of the procedures is planned under the scope of this proposed survey. The Toolbox measures are located in Attachments 1-38, and the additional measures are located in Attachments 39-59.

B.5 Individuals Consulted on Statistical Aspects and Individuals Collecting and/or Analyzing Data

Role	Name	Affiliation	Telephone number
Consultation on design aspects	Jennifer Beaumont, MS	NU-MSS	312-503-9805
	Richard Havlik, MD, MPH	Westat	301-652-3427
	Ron Hays, PhD	UCLA	310-794-2294
	Kay Savio	Delve	314-851-3116
	Kim Reale	Delve	314-851-3124
Interviewing contractor	Judy Golas	Delve	740-392-9436
Data analysis	Jennifer Beaumont, MS	NU-MSS	312-695-1324
	Richard Havlik, MD, MPH	Westat	301-503-9805
	Ron Hays, PhD	UCLA	310-794-2294
	Devin Peipert, MSW	NU-MSS	312-503-3647 end_of_the_skype_highlighting
	Katy Wortman, MSW	NU-MSS	312-503-3609 end_of_the_skype_highlighting
	James Burns, MS	NU-MSS	312-503-3380 end_of_the_skype_highlighting
	Seung Choi, PhD	NU-MSS	312-503-9803 end_of_the_skype_highlighting
	Jin-Shei Lai, PhD, OTR/L	NU-MSS	312-503-3370 end_of_the_skype_highlighting
	Richard Gershon, PhD	NU-MSS	312-503-3453 end_of_the_skype_highlighting
	Natalie McKinney, BA	NU-MSS	312-503-3909 end_of_the_skype_highlighting
	Tracy Podrabsky, BS	NU-MSS	312-503-3908 end_of_the_skype_highlighting
	Jamie Griffith, PhD	NU-MSS	312-503-3538 end_of_the_skype_highlighting
	Elizabeth Hahn, MA	NU-MSS	312-503-9804 end_of_the_skype_highlighting

