

## FORM C-2 Soybean Pre-Harvest Lab Determinations 2011



	YEAR, CROP, FORM, MMDD (1 – 7)		
	125		
	Date Sample Received in Lab:		
WI	EIGHT and NUMBER of UNTHRESHED PODS		
1.	Weight of Unit 1 pods and beans removed from bag	Grams to Tenths	503
2.	Weight of Unit 2 pods and beans removed from bag	Grams to Tenths	504
CO	OUNT of PODS from ONE UNIT (BAG) ONLY		
3.	Unit used (Always use pods from Unit 1, if possible)	Unit Code	512
4.	Number of pods with developed beans	Number	513
	(Developed beans are at least 50% of the mass of normal beans in that field. Generally, they are thicker than a nickel.)		
5.	Number of pods with undeveloped beans	Number	514
WI	EIGHT and MOISTURE of THRESHED BEANS		
	resh and hull only pods with developed beans from both units. If pods are too wet to thresh easily, ds should be dried for a short period at no more than 70 degrees C and then threshed.		
6.	Weight of all threshed beans from both units immediately before moisture test	Grams to Tenths	507
7.	Moisture content <sup>1/</sup>	Percent	508
I al	b Technician Date Analyzed		
∟dl	Date Affalyzed	MM	DD

<sup>&</sup>lt;sup>1</sup>If the sample weight is too small or too dry for a moisture test, follow the procedures on the back of this form to complete the moisture test.

If the sample weight is too small for moisture test, sufficient grains of known moisture content (use same class and stage of maturity) will be added to the sample so that a moisture test can be made. The moisture content of the sample can then be derived using the following formula:

$$E = \frac{(A+B)D - (BxC)}{A}$$

			•
Where	A = Weight of small or dry soybean sample	•—	Grams
	B = Weight of additional beans required for moisture test		Grams
	C = Moisture percent of B	•	Percent
	D = Moisture percent of A + B combined	•	Percent
	<b>E = Result : Moisture percent of small or dry soybean sample</b> ( <i>enter in item 7</i> )		Percent