



FORM C – 2R

Soybean Pre-Harvest Lab Determinations

Soybean Research Project

2011



| | |
|---------------------------------------|--|
| YEAR, CROP, FORM, MMDD (1 – 7) | |
| 1 2 5 ___ ___ ___ ___ | |

Date Sample Received in Lab: _____

WEIGHT and NUMBER of UNTHRESHED PODS

1. Weight of Unit 1 pods and beans removed from bag
2. Weight of Unit 2 pods and beans removed from bag

| | |
|------------------------|--------|
| | 503 |
| Grams to Tenths | . ____ |
| Grams to Tenths | 504 |
| | . ____ |

COUNT of PODS

3. Unit used (*Always use pods from Unit 1, if possible.*)
4. Number of pods with developed beans.
(Developed beans are at least 50% of the mass of normal beans in that field. Generally, they are thicker than a nickel.)
5. Number of pods with undeveloped beans

| | |
|------------------|-----|
| | 512 |
| Unit Code | |
| Number | 513 |
| Number | 514 |

WEIGHT and MOISTURE of THRESHED BEANS

Thresh and hull only pods with developed beans from both units. If pods are too wet to thresh easily, pods should be dried for a short period at no more than 70 degrees C and then threshed.

- Number of seeds (all threshed beans) from pods **Unit 1**.
- Number of seeds (all threshed beans) from pods **Unit 2**.
- Weight of threshed beans from **only Unit 1** immediately before moisture test.
6. Weight of all threshed beans from both units immediately before moisture test
7. Moisture content ^{1/}

| | |
|------------------------|--------|
| | Number |
| Number | . ____ |
| Grams to Tenths | |
| Grams to Tenths | 507 |
| Grams to Tenths | . ____ |
| Percent | 508 |
| | . ____ |

Lab Technician _____ Date Analyzed _____

MM DD

^{1/} If the sample weight is too small or too dry for a moisture test, follow the procedures on the back of this form to complete the moisture test.

FORM C-2R: SOYBEANS - continued

If the sample weight is too small for moisture test, sufficient grains of known moisture content (use same class and stage of maturity) will be added to the sample so that a moisture test can be made. The moisture content of the sample can then be derived using the following formula:

$$E = \frac{(A + B) D - (B \times C)}{A}$$

- Where
- A = Weight of small or dry soybean sample
 - B = Weight of additional beans required for moisture test
 - C = Moisture percent of B
 - D = Moisture percent of A + B combined
 - E = Result : Moisture percent of small or dry soybean sample (enter in item 7)

| | |
|--------|---------|
| . ____ | Grams |
| . ____ | Grams |
| . ____ | Percent |
| . ____ | Percent |
| . ____ | Percent |