This survey is being sent to a selected sample of health centers and providers. Please do not distribute to others for completion.

Physician

Physician assistant

Form Approved OMB Number: 0920-XXXX Expiration Date: XX/XX/XXXX

2012–2013 SURVEY of HEALTH CARE PROVIDERS

Public reporting burden of this collection of information is estimated to average 15 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Reports Clearance Officer, 1600 Clifton Road, MS D-74, Atlanta, GA 30333, ATTN: PRA (0920-xxxx).

I. PROVIDER, PATIENT and PRACTICE/ HEALTH CENTER CHARACTERISTICS

Please answer each of the following questions as they relate to you, your patients, and the practice/health center at which you received this survey. Which of the following describes the setting of Other (please specify) this practice/health center? (select all that apply) Community health center What is your primary clinical focus at this practice/health center? (select one) Family planning clinic Adolescent health or pediatrics Health department (state or local) Family medicine **HMO** or Hospital Obstetrics/gynecology or family Indian Health Service planning/reproductive health Planned Parenthood affiliate Primary (general health) care Private practice Other (please specify) School based health clinic Sexually transmitted infection clinic How many years has it been since you completed your most recent formal clinical training (e.g., University clinic medical/nursing school, residency/practicum/ Other (please specify) clinical)? Less than 5 years Does this practice/health center receive any non-5-14 years fee-for-service income to support family planning **Services?** (select all that apply) 15-24 years None 25 or more years Private grant(s) What is your gender? State appropriations Section 308 of Public Health Service Act Male Female Title V (MCH Block Grant) Title X (Family Planning) On average, how many female patients of Don't know reproductive age do you see per week? Other To approximately what percent of your **female** 10. In what state is your practice/health center patients of reproductive age do you provide located? family planning services*? 0% In this practice/health center, how many health 1-24% care providers, including you, provide family 25-49% planning services*? 50-74% What is your role as a health care provider? 75% or more (select one) Certified nurse midwife * For the purpose of this survey, a family planning service is any service related to postponing or preventing pregnancy. Family planning Nurse practitioner services may include a medical examination related to provision of a Nurse method, contraceptive counseling, method prescription or supply visits. A

patient may receive a family planning service even if the primary purpose

of her visit is not for contraception.

during the following time periods?	<u>sertion</u> or the	, , , , , , , , , , , , , , , , , , , ,	, commuco				
3	Trained to insert during routine care		Trained immed postp	- 1	Trained to insert immediately post-abortion		
	Yes	No	Yes	No	Yes	No	
Copper intrauterine device (Cu-IUD or ParaGard®)?							
Levonorgestrel-releasing intrauterine device (LNG-IUD or Mirena®)?							
Contraceptive implant (Implanon®/Nexplanon®)?			N/A	N/A	N/A	N/A	
12. Approximately what percentages of your <u>fem</u> characteristics? If unsure, give your best esti		<u> </u>					
)-24%	25-49%	≥	:50%	
Pay for their visit using Medicaid or other state or feder	ral assistance	?					
Are racial or ethnic minorities?							
Have limited English proficiency?							
Are adolescents?							
Are 35 years of age or older?							
, ,							
	_						
II. HEALTH CARE PROVIDER ATTITUDES	<u> </u>						
Please answer each of the following question	ns as thev r	elate to v	our attitu	des when p	rovidina f	amilv	
planning services. Please do not consult an							
	-	•		•	•		
13. How <u>safe</u> do you consider <u>combined oral con</u>	<u>itraceptives</u>	COCs) to	be for the	ollowing gro	oups?		
13. How <u>safe</u> do you consider <u>combined oral con</u>	itraceptives (COCs) to Ver saf	y Safe		Very unsafe	Don't know	
13. How <u>safe</u> do you consider <u>combined oral consider combined oral consider safe</u> . Breastfeeding women ≥ 1 month postpartum without of for venous thromboembolism (VTE)	·	Ver saf	y Safe		Very		
Breastfeeding women ≥ 1 month postpartum without of	·	Ver saf	y Safe		Very		
Breastfeeding women ≥ 1 month postpartum without of for venous thromboembolism (VTE) Smokers 35 years of age or older	·	Ver saf	y Safe		Very		
Breastfeeding women ≥ 1 month postpartum without of for venous thromboembolism (VTE) Smokers 35 years of age or older Obese women (BMI ≥30 kg/m²) Women with a history of bariatric surgery via restrictive	ther risk facto	Ver saf	y Safe		Very		
Breastfeeding women ≥ 1 month postpartum without of for venous thromboembolism (VTE) Smokers 35 years of age or older Obese women (BMI ≥30 kg/m²) Women with a history of bariatric surgery via restrictive (e.g., vertical banded gastroplasty) Women with a history of bariatric surgery via malabsor	ther risk facto	Ver saf	y Safe		Very		
Breastfeeding women ≥ 1 month postpartum without of for venous thromboembolism (VTE) Smokers 35 years of age or older Obese women (BMI ≥30 kg/m²) Women with a history of bariatric surgery via restrictive (e.g., vertical banded gastroplasty) Women with a history of bariatric surgery via malabsor procedures (e.g., Roux-en-Y gastric bypass)	ther risk facto	Ver saf	y Safe		Very		
Breastfeeding women ≥ 1 month postpartum without of for venous thromboembolism (VTE) Smokers 35 years of age or older Obese women (BMI ≥30 kg/m²) Women with a history of bariatric surgery via restrictive (e.g., vertical banded gastroplasty) Women with a history of bariatric surgery via malabsor procedures (e.g., Roux-en-Y gastric bypass) Women with rheumatoid arthritis Women with inflammatory bowel disease (i.e., ulcerative)	ther risk facto e procedures ptive	Ver saf	y Safe		Very		
Breastfeeding women ≥ 1 month postpartum without of for venous thromboembolism (VTE) Smokers 35 years of age or older Obese women (BMI ≥30 kg/m²) Women with a history of bariatric surgery via restrictive (e.g., vertical banded gastroplasty) Women with a history of bariatric surgery via malabsor procedures (e.g., Roux-en-Y gastric bypass) Women with rheumatoid arthritis	ther risk facto e procedures ptive ve colitis, Cro	Ver saf	Safe	Unsafe	Very unsafe		
Breastfeeding women ≥ 1 month postpartum without of for venous thromboembolism (VTE) Smokers 35 years of age or older Obese women (BMI ≥30 kg/m²) Women with a history of bariatric surgery via restrictive (e.g., vertical banded gastroplasty) Women with a history of bariatric surgery via malabsor procedures (e.g., Roux-en-Y gastric bypass) Women with rheumatoid arthritis Women with inflammatory bowel disease (i.e., ulcerative disease) without other risk factors for VTE	ther risk facto e procedures ptive ve colitis, Cro	Ver saf	Safe	Unsafe	Very unsafe	know	
Breastfeeding women ≥ 1 month postpartum without of for venous thromboembolism (VTE) Smokers 35 years of age or older Obese women (BMI ≥30 kg/m²) Women with a history of bariatric surgery via restrictive (e.g., vertical banded gastroplasty) Women with a history of bariatric surgery via malabsor procedures (e.g., Roux-en-Y gastric bypass) Women with rheumatoid arthritis Women with inflammatory bowel disease (i.e., ulcerative disease) without other risk factors for VTE	ther risk facto e procedures ptive ve colitis, Cro	Ver saf	Safe	Unsafe	Very unsafe		
Breastfeeding women ≥ 1 month postpartum without of for venous thromboembolism (VTE) Smokers 35 years of age or older Obese women (BMI ≥30 kg/m²) Women with a history of bariatric surgery via restrictive (e.g., vertical banded gastroplasty) Women with a history of bariatric surgery via malabsor procedures (e.g., Roux-en-Y gastric bypass) Women with rheumatoid arthritis Women with inflammatory bowel disease (i.e., ulcerative disease) without other risk factors for VTE	ther risk facto e procedures ptive ve colitis, Cro	Ver saf	Safe	Unsafe	Very unsafe	know	
Breastfeeding women ≥ 1 month postpartum without of for venous thromboembolism (VTE) Smokers 35 years of age or older Obese women (BMI ≥30 kg/m²) Women with a history of bariatric surgery via restrictive (e.g., vertical banded gastroplasty) Women with a history of bariatric surgery via malabsor procedures (e.g., Roux-en-Y gastric bypass) Women with rheumatoid arthritis Women with inflammatory bowel disease (i.e., ulcerative disease) without other risk factors for VTE 14. How safe do you consider intrauterine devices	ther risk facto e procedures ptive ve colitis, Crol	Ver saf	Safe	Unsafe	Very unsafe	know	
Breastfeeding women ≥ 1 month postpartum without of for venous thromboembolism (VTE) Smokers 35 years of age or older Obese women (BMI ≥30 kg/m²) Women with a history of bariatric surgery via restrictive (e.g., vertical banded gastroplasty) Women with a history of bariatric surgery via malabsor procedures (e.g., Roux-en-Y gastric bypass) Women with rheumatoid arthritis Women with inflammatory bowel disease (i.e., ulcerative disease) without other risk factors for VTE 14. How safe do you consider intrauterine devices Adolescents Immediately postpartum women (less than 10 minutes)	ther risk facto e procedures ptive ve colitis, Crol es (Cu-IUD or	Ver saf	Safe	Unsafe	Very unsafe	know	
Breastfeeding women ≥ 1 month postpartum without of for venous thromboembolism (VTE) Smokers 35 years of age or older Obese women (BMI ≥30 kg/m²) Women with a history of bariatric surgery via restrictive (e.g., vertical banded gastroplasty) Women with a history of bariatric surgery via malabsor procedures (e.g., Roux-en-Y gastric bypass) Women with rheumatoid arthritis Women with inflammatory bowel disease (i.e., ulcerative disease) without other risk factors for VTE 14. How safe do you consider intrauterine devices Adolescents Immediately postpartum women (less than 10 minutes placenta) Postpartum women (10 minutes after delivery of placer	ther risk facto e procedures ptive ve colitis, Crol es (Cu-IUD or	Ver saf	Safe	Unsafe	Very unsafe	know	
Breastfeeding women ≥ 1 month postpartum without of for venous thromboembolism (VTE) Smokers 35 years of age or older Obese women (BMI ≥30 kg/m²) Women with a history of bariatric surgery via restrictive (e.g., vertical banded gastroplasty) Women with a history of bariatric surgery via malabsor procedures (e.g., Roux-en-Y gastric bypass) Women with rheumatoid arthritis Women with inflammatory bowel disease (i.e., ulcerative disease) without other risk factors for VTE 14. How safe do you consider intrauterine devices Adolescents Immediately postpartum women (less than 10 minutes placenta) Postpartum women (10 minutes after delivery of placer 4 weeks postpartum) Nulliparous women	ther risk facto e procedures ptive ve colitis, Crol es (Cu-IUD or	Ver saf	Safe	Unsafe	Very unsafe	know	
Breastfeeding women ≥ 1 month postpartum without of for venous thromboembolism (VTE) Smokers 35 years of age or older Obese women (BMI ≥30 kg/m²) Women with a history of bariatric surgery via restrictive (e.g., vertical banded gastroplasty) Women with a history of bariatric surgery via malabsor procedures (e.g., Roux-en-Y gastric bypass) Women with rheumatoid arthritis Women with inflammatory bowel disease (i.e., ulcerative disease) without other risk factors for VTE 14. How safe do you consider intrauterine devices Adolescents Immediately postpartum women (less than 10 minutes placenta) Postpartum women (10 minutes after delivery of placer 4 weeks postpartum)	ther risk facto e procedures ptive ve colitis, Crol es (Cu-IUD or	Ver saf	Safe	Unsafe	Very unsafe	know	

		Very safe	Safe	Unsafe	Very unsafe	Don't know
Adolescents						
Breastfeeding women <1 month postpartum						
Breastfeeding women ≥ 1 month postpartum						
Smokers 35 years of age or older						
Obese women (BMI ≥30 kg/m²)						
Women with a history of bariatric surgery						
Women with rheumatoid arthritis not on immunosuppressiv	e therapy					
Women with inflammatory bowel disease	1 7					
Women with complicated diabetes (i.e., nephropathy, retino	opathy,					
neuropathy, other vascular disease or diabetes of >20 year duration)						
16. For each of the following contraceptive methods, her visit regardless of the timing of her menses (pregnant? Please answer for both adolescents a	'Quick Sta	rt') <u>if you</u>				
		Adolescen	ts			
	Safe	Unsafe	Don't know	Safe	Unsafe	Don't know
Combined hormonal contraceptives (COCs, patch, ring)						
DMPA						
Contraceptive implant						
Intrauterine devices (Cu-IUD or LNG-IUD)						
III. HEALTH CARE PROVIDER PRACTICES Please answer each of the following questions a when providing family planning services. 17. In the past month, when counseling your typical often have you (or your clinical team) done the form	female pa	_				
	J	_	Very often	Often	Not often	Never
Assessed the patient's reproductive life plan (i.e., asked ab						
intentions regarding the number and timing of pregnancies their personal values and life goals)					Ш	
Presented information regarding potential contraceptive memost effective methods presented first (tiered approach)						
Helped the patient think about potential barriers to using th method correctly and develop a plan to deal with these bar		d				
Used a method-specific informed consent form						
Informed adolescents that long-acting reversible contracep effective options	tives are s	afe and				

How safe do you consider DMPA (Depo-Provera®) to be for the following groups?

15.

18. In the past year, how often have you (or your clinical team) provided <u>DMPA</u> to <u>adolescents</u>?

	Very often or often	□ }	Go to quest	ion #19.									
	Not often or never	□ }	If "not often	or never" please	indicate why. (sele	ct all that apply)							
		a.	I rarely have adolescents as patients										
		b.	DMPA is unavailable in my practice/health center										
		C.	I am concerr	I am concerned about the safety of DMPA for adolescents									
		d.	I am concerr	am concerned about side effects that may lead to discontinuation									
		e.	My adolesce	y adolescent patients generally prefer a different method									
		f.	My practice/l	health center protoc	ol does not allow it								
		g.	Other reason	ns (please specify) _									
19.	In the past year, ho women ≥ 1 month					cribed <u>COCs</u> to <u>br</u>	<u>eastfeeding</u>						
	Very often or often	□ }	Go to quest	ion #20.									
	Not often or never		If "not often	or never" please	indicate why. (sele	ct all that apply)							
		a.	I rarely have	postpartum women	as patients								
		b.			of COCs for breast								
		C.			se in breast milk pro								
		d.	My postpartu	ım patients generall	ly prefer a different i	method							
		e.	My practice/l	health center protoc	ol does not allow it								
		f.	Other reasor	ns (please specify) _									
	Very often or often	women'	Go to quest										
	Not often or never	} }	If "not often	or never" please	indicate why. (sele	ct all that apply)							
		a.	-	nulliparous women	•								
		b.		•									
		C.		•	of IUDs for nullipar	ous women	IUDs are generally unavailable in my practice/health center I am concerned about the safety of IUDs for nulliparous women						
		d.	I am concerr	ned about the effect:	I am concerned about the effects on future fertility								
			I am concerned about difficult insertion										
		e.		ned about difficult in	sertion								
		e. f.	My nulliparo	ned about difficult in us patients generall	sertion y prefer a different r	method							
		_	My nulliparou My practice/l	ned about difficult in us patients generall health center protoc	sertion y prefer a different r col does not allow it								
		f. g. h.	My nulliparou My practice/l Cost barriers	ned about difficult in us patients generall health center protoc s prevent me from p	sertion y prefer a different r								
		f. g.	My nulliparou My practice/l Cost barriers	ned about difficult in us patients generall health center protoc	sertion y prefer a different r col does not allow it								
21.	When initiating the require the following	f. g. h. i. e followi n	My nulliparou My practice/I Cost barriers Other reason g contracept	ned about difficult in us patients generally health center protoc s prevent me from p ns (please specify)_ ive methods, pleas	sertion y prefer a different r col does not allow it roviding IUDs to nul se indicate if you o Please check all e	lliparous women or your practice/he exams and tests th	at apply.						
21.		f. g. h. i. e followi n	My nulliparou My practice/I Cost barriers Other reason g contracept	ned about difficult in us patients generally health center protoc s prevent me from p ns (please specify)_ ive methods, pleas	sertion y prefer a different r col does not allow it roviding IUDs to nucleon se indicate if you o	lliparous women							
		f. g. h. i. e followi n	My nulliparou My practice/I Cost barriers Other reason g contracept s and tests fo Blood	ned about difficult in us patients generally health center protocks prevent me from pass (please specify)_tive methods, please a healthy client. Clinical breast	sertion y prefer a different recol does not allow it roviding IUDs to number of the see indicate if you of Please check all elements.	or your practice/heexams and tests the Cervical cytology	nat apply. Chlamydia <i>l</i> gonorrhea						
COC	require the following	f. g. h. i. e followin ng <u>exam</u>	My nulliparou My practice/I Cost barriers Other reason g contracept s and tests fo Blood	ned about difficult in us patients generally health center protocks prevent me from pass (please specify)_tive methods, please a healthy client. Clinical breast	sertion y prefer a different recol does not allow it roviding IUDs to number of the see indicate if you of Please check all elements.	or your practice/heexams and tests the Cervical cytology	nat apply. Chlamydia <i>l</i> gonorrhea						
COC	require the followings/patch/ringestin-only pills (POPs	f. g. h. i. e followin ng <u>exam</u>	My nulliparou My practice/I Cost barriers Other reason g contracept s and tests fo Blood	ned about difficult in us patients generally health center protocks prevent me from pass (please specify)_tive methods, please a healthy client. Clinical breast	sertion y prefer a different recol does not allow it roviding IUDs to number of the see indicate if you of Please check all elements.	or your practice/heexams and tests the Cervical cytology	nat apply. Chlamydia <i>l</i> gonorrhea						
COC Prog DMP	require the followings/patch/ringestin-only pills (POPs	f. g. h. i. e followin ng <u>exam</u>	My nulliparou My practice/I Cost barriers Other reason g contracept s and tests fo Blood	ned about difficult in us patients generally health center protocks prevent me from pass (please specify)_tive methods, please a healthy client. Clinical breast	sertion y prefer a different recol does not allow it roviding IUDs to number of the see indicate if you of Please check all elements.	or your practice/heexams and tests the Cervical cytology	nat apply. Chlamydia <i>l</i> gonorrhea						
COC Prog DMP	require the followings/patch/ring estin-only pills (POPs	f. g. h. i. e followin ng <u>exam</u>	My nulliparou My practice/I Cost barriers Other reason g contracept s and tests fo Blood	ned about difficult in us patients generally health center protocks prevent me from pass (please specify)_ive methods, please a healthy client. Clinical breast	sertion y prefer a different recol does not allow it roviding IUDs to number of the see indicate if you of Please check all elements.	or your practice/heexams and tests the Cervical cytology	nat apply. Chlamydia <i>l</i> gonorrhea						

	did you <u>start a woman on the day of h</u> reasonably certain she was not pregna							rt') <u>if you</u>
	(22a) Adolescents					2b) Adult		
Very often or often	Go to question #22b		Very often		<u> </u>	uestion		
Not often	If "not often or never" please		or often Not often		If "not	often or	never" pleas	
or never a.	I do not think it is safe	(y)	or never				select all that ap	oply)
			a.	I do	not think it	is safe		
b.	I have liability concerns		b.	I hav	e liability o	concerns		
C.	I do not have enough training I do not think it is appropriate for		C.		not have e	_	_	
d.	adolescents		d.	I do . aduli	not think it ts	is approp	oriate for	
e.	My practice/health center protocol does not allow it		e.	Му р		alth cente	er protocol do	oes
f.	Other (please specify)		f.		anow it er (please :	snecify)		
the ti	e past year, when providing <u>DMPA</u> , <u>hov</u> ming of her menses ('Quick Start') <u>if yo</u> oth adolescents and adults.		l you <u>start a </u>	woma	n on the o	day of he		
	(23a) Adolescents				(23	Bb) Adult	:s	
Very often or often	Go to question #23b		Very often or often		Go to q	uestion	#24	
Not often	If "not often or never" please		Not often or never				never" pleas select all that ap	
or never a.	I do not think it is safe	(V)	a.	I do	not think it		seiect air triat ap	рріу)
b.	I have liability concerns		b.		e liability o			
C.	I do not have enough training		Б. С.		not have e			
	I do not think it is appropriate for				not think it	_	_	
d.	adolescents	_ ⊔	d.	aduli	ts			
e.	My practice/health center protocol does not allow it		e.		ractice/ne allow it	aith cente	er protocol do	bes
f.	Other (please specify)		f.	Othe	er (please s	specify) _		
	initiating the following methods, pleas follow-up visit.	e indicate	when you ac	dvise	healthy a	dult patie	ents to come	back
101 α	ionow-up visit.						Only	y if she
		4-6 week	s 3 month	ıs 6	months	mont	has p	roblems uestions
COCs, patc	h, ring							
POPs								
DMPA (rout	tine follow-up other than for re-injection)							
Implant	, , , , , , , , , , , , , , , , , , ,							
Intrauterine	device (Cu-IUD or LNG-IUD)							
25. In the	nest year how often have you or you	r aliniaal t	nom dono the	o follo	wing?			
23. In the	e past year, <u>how often</u> have you or you	r Cillical te	eam done un			Often	Not often	Never
	advance prescription for emergency con	traception	(EC) to a	very			NOT OILEIT	INEVE
	specifically seeking EC	no oificelle :	poolsing FC					
	n <u>advance supply</u> of EC to a woman not s	· -	~	L				
	prescribed a contraceptive at the same to Cu-IUD as EC	me you pro	Ivided EC	L				
rioviueu a	Cu-IUD as EC			L				

In the past year, <u>how often</u> did you or your clinical team <u>dispense</u> a year's supply of pills (COCs or POPs) <u>at</u>

one visit? Please answer for both new and continuing users.

In the past year, when providing or prescribing combined hormonal contraceptives (COCs, patch, ring), how

22.

			(26a) NEW USERS	(26b) CONTINUING USERS					
		ery often Go to question #26b			Very often Go to question #27				
	Not of			Not often					
	or nev	, , , , , , , , , , , , , , , , , ,		or never	_	ndicate why. (select	all that apply)		
		a.	I do not think it is safe	a.		I do not think it is safe			
		b. My practice/health center does not dispense pills		b.	dispens	My practice/health center does not dispense pills			
	c. My practice/health center protocol does not allow it		C.	not allo	My practice/health center protocol does not allow it				
		d.	I have liability concerns	d.		I have liability concerns			
		e.	There is not enough supply in my practice/health center	e.	practice	s not enough supply i health center			
		f.	It is too expensive for my practice/health center	f.	center	expensive for my pra			
		g.	I am concerned about wasting pill packs if the woman discontinues	g.		ncerned about wastir nan discontinues	ng pill packs if		
		h.	Other (please specify)	h.	Other (p	olease specify)			
		27. For routine health care, at what age do you or your practice/health center recommend that a woman begin routine cervical cancer screening? (select all that apply)		28. For routine health care, how often do you provide cervical cancer screening for a sexually active, 25-year old patient with previously normal results?					
	When	ever s	he becomes sexually active	Every visit					
				Annually					
	Starting at age 18 Starting at age 21		Every 2 years						
	Don't	_	9	Every 3 years Don't know					
			se specify)			Δ			
	Outci	(picus	Se speelify	Other (plea	se specify	/)			
29.			neral, how important to you are the following so ices related to contraception? Please answer for			rmed about recomn	nended clinical		
			•		rtant	Minor Source	Not Used		
				Sou	ırce				
		rence							
			education programs	L					
			with colleagues	L					
	Intern	et sea	rches/online resources						
	Institu	itional	practice protocols						
	Journ	als							
	Medic	ation _l	oackage inserts						
	Profes	ssiona	l organization publications or notifications						
	Textb	ooks (e.g., Contraceptive Technology)						
	U.S. N	Medica	Il Eligibility Criteria for Contraceptive Use (MEC)	Γ					
	WHO		, ,						
			ted Practice Recommendations for Contraceptive L	lse [
			se specify):						
	001	₁ 0.000	· _[]],						

V. AWARENESS OF GUIDELINES

We want to know about your awareness of CDC's 2010 U.S. Medical Eligibility Criteria for Contraceptive Use (U.S. MEC).

30.	How did you learn about CDC's 2010 U.S. MEC? (select all that apply)	
	I did not know about CDC's 2010 U.S. MEC before participation in this survey	
	Conference attendance	
	Continuing medical education programs	
	Discussions with colleagues	
	Internet searches/online resources	
	Institutional practice protocol	
	Journals	
	Professional organization publications or notifications	
	Textbooks (e.g., Contraceptive Technology)	
	Other (please specify)	
31.	Have you <u>used</u> any of the following U.S. MEC materials?	
	U.S. MEC website	
	U.S. MEC color-coded summary chart in English	
	U.S. MEC color-coded summary chart in Spanish	
	U.S. MEC wheel	
	U.S. MEC PDA application	
	U.S. MEC 2011 update with revised recommendations for postpartum contraceptive use	
	U.S. MEC 2012 update with revised recommendations for the use of hormonal contraception among women at high risk for HIV infection or infection with HIV	
32.	What <u>additional medical conditions or patient characteristics</u> would you like to see recommendathe U.S. MEC?	ations for in
	(please specify)	· · · · · · · · · · · · · · · · · · ·
	(please specify)	
	(please specify)	
P	Please share any additional comments that you may have in the space below.	

Thank you for completing this survey!
Please return using the enclosed postage paid envelope.