C7. Arctic Fetal Alcohol Spectrum Disorder Regional Training Center Intro to FASDs Workshop Pre-Test

For each item, please choose the response that best completes the sentence or answers the question.

- 1. The best public health message related to alcohol consumption by women of childbearing age is...
 - a. Women who know they are pregnant should not consume alcohol.
 - b. Women who are pregnant or could become pregnant should not consume alcohol.
 - c. All sexually active women should not consume alcohol.
 - d. Women of childbearing age should not consume alcohol.
 - e. Alcohol use, in moderation, is safe for all women.
- 2. An episode of binge or risky drinking for women of childbearing age (18-44) is defined as more than ______ standard drinks in about a two hour period.
 - a. 2
 - b. 3
 - c. 4
 - d. 5
 - e. 6
- 3. Which one of these is NOT considered a standard drink?
 - a. 12 ounces of beer
 - b. 12 ounces of wine cooler
 - c. 5 ounces of wine
 - d. 3 ounces of vodka
 - e. All of the above are standard drinks
- 4. Positive responses given to items on the TWEAK or the T-ACE screening instruments indicate...
 - a. a diagnosis of alcoholism
 - b. the likelihood that the woman will seek treatment for alcohol abuse
 - c. the need for the health care provider to discuss the risks associated with drinking alcohol, especially when the woman is pregnant
 - d. the difference between binge drinking and problem drinking
 - e. the need for immediate admission to a treatment facility for further evaluation
- 5. Scientific studies of the effects of alcohol on the fetus have shown that...
 - a. Alcohol has no effect on the development of the fetus in the third trimester.
 - b. Alcohol can only affect fetal development in the first trimester.
 - c. A small to moderate amount of alcohol is safest during the second trimester.
 - d. Of all the different types of alcohol, wine is the safest for a pregnant woman to consume.
 - e. None of the above is correct.

The public reporting burden of this collection of information is estimated to average 15 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to - CDC/ATSDR Reports Clearance Officer; 1600 Clifton Road NE, MS D-74, Atlanta, Georgia 30333 ATTN: PRA (XXXX-XXXX)

- 6. Although specific deficits experienced by individuals with an FASD vary widely, behavioral deficits are likely to include which of the following?
- a. memory problems, underactivity, and self-injurious behavior
- b. inattention, memory problems and hyperactivity
- c. particular mannerisms, hoarding, and memory problems
- d. emotion regulation, weight loss, fatigue, irritability
- e. auditory hallucinations, disorganized speech, and flat affect
- 7. Which of the following include all three facial abnormalities associated with fetal alcohol syndrome?
 - a. Low set ears, large palpebral fissures, large forehead
 - b. Small palpebral fissures, smooth philtrum, thin upper lip
 - c. Large palpebral fissures, smooth philtrum, thin upper lip
 - d. Large forehead, flattened cheeks, small palpebral fissures
 - e. Thin upper lip, distinct philtrum, ectropion
- 8. Which of the following is NOT required to confirm a diagnosis of fetal alcohol syndrome (FAS)?
 - a. Facial dysmorphia
 - b. Growth retardation
 - c. Central nervous system (CNS) abnormalities
 - d. Documentation of prenatal alcohol exposure
 - e. All of the above are required for a diagnosis of FAS
- 9. A variety of strategies have been found to be effective when working with individuals with an FASD. Which one of the following may not be as effective?
 - a. Concise, explicit instructions
 - b. Stable routines
 - c. Social cues
 - d. Visual aids
 - e. Repetition
- 10. Which organ system is primarily susceptible to damage or alteration caused by prenatal alcohol exposure?
- a. The circulatory system
- b. The skeletal system
- c. The central nervous system
- d. The reproductive system
- e. The digestive system
- 11. Growth deficit, central nervous system abnormalities, and a specific set of craniofacial features are all necessary for a diagnosis of which of the following conditions?
 - a. Fetal alcohol syndrome
 - b. Fetal alcohol spectrum disorder
 - c. Alcohol-related neurodevelopmental disorder
 - d. Fetal alcohol effects
 - e. Static encephalopathy/alcohol exposed
- 12. How is the University of Washington 4-Digit Diagnostic Code System used to identify an FASD?

- a. A physician diagnoses based on medical history and anthropomorphic measurements.
- b. A multidisciplinary team diagnoses based on the individual's dysmorphic features, growth parameters, and neurodevelopmental functioning.
- c. A neurologist diagnoses based on MRI results.
- d. Either a physician or psychologist diagnoses based on evidence of prenatal alcohol exposure.
- e. A psychologist diagnoses based on results from a battery of neuropsychological tests.
- 13. Which of the following best describes common impairments of the central nervous system among individuals who experience FASDs?
 - a. Abstract reasoning, eye-to-eye gaze, body postures, repetitive motor mannerisms
 - b. Poor organization, trouble with transitions, visual-spatial deficits, clumsiness
 - c. Hypervigilance, concentration, smooth handwriting, developmental milestone achievement
 - d. Verbal ability, underactivity, capacity to self-monitor, appreciation of nonverbal cues
 - e. All of the above
- 14. When caring for an individual with a known FASD, it helpful to consider _____ rather than _____ in order to better understand ability and behavior.
 - a. Developmental age; chronological age
 - b. Strengths; weaknesses
 - c. Protective factors; risk factors
 - d. Environment; mood
 - e. Chronological; adaptive
- 15. Which one of the following is NOT correct?

Persons with an FASD ...

- a. May experience problems in school and getting along with other children.
- b. Do not typically need support and services to live and work independently.
- c. Can benefit from obtaining an early diagnosis.
- d. May need support and services throughout their lives.
- e. May be at an increased risk for victimization and high risk behaviors.
- 16. Which of the following are considered protective factors against secondary disabilities associated with FASDs?
 - a. Living in stable and nurturing home environments
 - b. Early diagnosis (before 6 years of age)
 - c. Consistency in caregivers
 - d. Eligibility for social and educational services
 - e. All of the above

Please respond to the following questions about your professional experiences.

1. In your current position, do you... provide services to women of childbearing age? On a scale from 0 to 10 where 0 means "Not confident in my skills" and 10 means "Totally confident in my skills," how confident are you in your skills to do the following? (circle one number per row)

	<u>Not</u> confident in my skills										<u>Totally</u> <u>confident in</u> <u>my skills</u>
Screen women for risky or hazardous drinking	0	1	2	3	4	5	6	7	8	9	10
Educate pregnant women about the effects of alcohol on their babies	0	1	2	3	4	5	6	7	8	9	10
Conduct brief interventions for reducing alcohol consumption	0	1	2	3	4	5	6	7	8	9	10
Utilize resources to refer patients who need formal treatment for alcohol abuse	0	1	2	3	4	5	6	7	8	9	10
Identify persons with possible FAS or other prenatal alcohol-related disorders	0	1	2	3	4	5	6	7	8	9	10
Diagnose persons with possible FAS or other prenatal alcohol-related disorders	0	1	2	3	4	5	6	7	8	9	10
Utilize resources to refer patients for diagnosis and/or treatment services	0	1	2	3	4	5	6	7	8	9	10
Manage/coordinate the treatment of persons with FASDs	0	1	2	3	4	5	6	7	8	9	10

Are you (Circle one):	If you are a PROFESSIONAL, please	If you are a STUDENT OR				
Male	circle the one that best represents your	RESIDENT , please circle all that				
Female	current position:	apply:				
Transgender						
	ALLIED HEALTH PROFESSIONAL	ALLIED HEALTH STUDENT				
	Counselor	Human Services				
Are you Hispanic or Latino(a)?	OT/PT/SLP	OT/PT/SLP				
Yes	Psychologist	Psychology				
No	Social Worker	Social Work				
	Other Allied Health Professional:	Other Allied Health Student:				
	Specify:	Specify:				
How do you describe your race?						
(Select all that apply)	DENTAL PROFESSIONAL					
	Dental Assistant	HEALTH STUDENT				
American Indian/Alaska Native	Dental Hygienist	Dental Student				
Asian	Dentist	Medical Student				
Black or African American		Nursing Student				
Native Hawaiian/Pacific Islander	MEDICAL PROFESSIONAL					
White	Family Physician					
	Internal Medicine Physician	MEDICAL RESIDENT				
	Nurse (NP, RN, LPN)	Family Physician				
	OB/GYN	Internist				
	Pediatrician	OB/GYN				
	Physician Assistant	Pediatrician				
	Psychiatrist	Psychiatrist				
	Other Medical Professional	Other Resident				
I am the parent/caregiver	Specify:	Specify:				
of a child with FAS/FASD						
	OTHER	OTHER STUDENT				
	Administrator	Counseling Education				
	Corrections	Education				
	Educator, PreK-12	Other Student				
	Educator, PreK-12 SpED	Specify:				
	Educator, Other	1 5				
	Specify:					
	Lawyer/Judge					
	Public Health					
	Scientist					
	Other Professional:					
	Specify:					