

**C10. Arctic Fetal Alcohol Spectrum Disorder Regional Training Center
Train-the-Trainer Workshop Pre-Test**

For each item, please choose the response that best completes the sentence or answers the question.

1. The best public health message related to alcohol consumption by women of childbearing age is...
 - a. Women who know they are pregnant should not consume alcohol.
 - b. Women who are pregnant or could become pregnant should not consume alcohol.
 - c. All sexually active women should not consume alcohol.
 - d. Women of childbearing age should not consume alcohol.
 - e. Alcohol use, in moderation, is safe for all women.

2. An episode of binge or risky drinking for women of childbearing age (18-44) is defined as more than ____ standard drinks in about a two hour period.
 - a. 2
 - b. 3
 - c. 4
 - d. 5
 - e. 6

3. Which one of these is NOT considered a standard drink?
 - a. 12 ounces of beer
 - b. 12 ounces of wine cooler
 - c. 5 ounces of wine
 - d. 3 ounces of vodka
 - e. All of the above are standard drinks

4. Positive responses given to items on the TWEAK or the T-ACE screening instruments indicate...
 - a. a diagnosis of alcoholism
 - b. the likelihood that the woman will seek treatment for alcohol abuse
 - c. the need for the health care provider to discuss the risks associated with drinking alcohol, especially when the woman is pregnant
 - d. the difference between binge drinking and problem drinking
 - e. the need for immediate admission to a treatment facility for further evaluation

5. Which of these is NOT part of the addictive disease process?
 - a. Craving and compulsion
 - b. Violent behavior
 - c. Loss of control
 - d. Continued use despite adverse consequences
 - e. All of the above are part of the addictive disease process

6. Scientific studies of the effects of alcohol on the fetus have shown that...
 - a. Alcohol has no effect on the development of the fetus in the third trimester.
 - b. Alcohol can only affect fetal development in the first trimester.
 - c. A small to moderate amount of alcohol is safest during the second trimester.
 - d. Of all the different types of alcohol, wine is the safest for a pregnant woman to consume.
 - e. None of the above is correct.

7. Although specific deficits experienced by individuals with an FASD vary widely, behavioral deficits are likely to include which of the following?
 - a. memory problems, underactivity, and self-injurious behavior
 - b. inattention, memory problems and hyperactivity
 - c. particular mannerisms, hoarding, and memory problems
 - d. emotion regulation, weight loss, fatigue, irritability
 - e. auditory hallucinations, disorganized speech, and flat affect

8. Which of the following include all three facial abnormalities associated with fetal alcohol syndrome?
 - a. Low set ears, large palpebral fissures, large forehead
 - b. Small palpebral fissures, smooth philtrum, thin upper lip
 - c. Large palpebral fissures, smooth philtrum, thin upper lip
 - d. Large forehead, flattened cheeks, small palpebral fissures
 - e. Thin upper lip, distinct philtrum, ectropion

9. Which of the following is NOT required to confirm a diagnosis of fetal alcohol syndrome (FAS)?
 - a. Facial dysmorphia
 - b. Growth retardation
 - c. Central nervous system (CNS) abnormalities
 - d. Documentation of prenatal alcohol exposure
 - e. All of the above are required for a diagnosis of FAS

10. A variety of strategies have been found to be effective when working with individuals with an FASD. Which one of the following may not be as effective?
 - a. Concise, explicit instructions
 - b. Stable routines
 - c. Social cues
 - d. Visual aids
 - e. Repetition

11. Which one of the following statements is NOT correct?
 - a. Some women expose their fetus to alcohol because of alcohol dependence or addiction.
 - b. Health providers seek the best outcomes for women and fetuses by efforts to increase foreseeable benefits and minimize foreseeable harms.
 - c. Most medical societies advocate coercive methods as a first intervention for women who expose a fetus to alcohol.
 - d. Health providers have an ethical duty to respect their patients.
 - e. Confidentiality is an essential aspect of the health provider-patient relationship.

12. Which of the following is a validated screening tool commonly used to ascertain alcohol use among pregnant women?
 - a. SASSI-3
 - b. T-ACE
 - c. Addiction Severity Index
 - d. MMPI
 - e. DSM Screener

13. Which organ system is primarily susceptible to damage or alteration caused by prenatal alcohol exposure?
 - a. The circulatory system
 - b. The skeletal system
 - c. The central nervous system
 - d. The reproductive system
 - e. The digestive system

14. Preventing the harms of prenatal alcohol exposure includes which of the following proactive measures?
 - a. Education
 - b. Substance abuse treatment
 - c. Alcohol screening and referral services
 - d. Encouragement of preconception abstinence from alcohol
 - e. All of the above

15. Growth deficit, central nervous system abnormalities, and a specific set of craniofacial features are all necessary for a diagnosis of which of the following conditions?
 - a. Fetal alcohol syndrome
 - b. Fetal alcohol spectrum disorder
 - c. Alcohol-related neurodevelopmental disorder
 - d. Fetal alcohol effects
 - e. Static encephalopathy/alcohol exposed

16. How is the University of Washington 4-Digit Diagnostic Code System used to identify an FASD?
 - a. A physician diagnoses based on medical history and anthropomorphic measurements.
 - b. A multidisciplinary team diagnoses based on the individual's dysmorphic features, growth parameters, and neurodevelopmental functioning.
 - c. A neurologist diagnoses based on MRI results.
 - d. Either a physician or psychologist diagnoses based on evidence of prenatal alcohol exposure.
 - e. A psychologist diagnoses based on results from a battery of neuropsychological tests.

17. Which of the following best describes common impairments of the central nervous system among individuals who experience FASDs?
 - a. Abstract reasoning, eye-to-eye gaze, body postures, repetitive motor mannerisms

- b. Poor organization, trouble with transitions, visual-spatial deficits, clumsiness
 - c. Hypervigilance, concentration, smooth handwriting, developmental milestone achievement
 - d. Verbal ability, underactivity, capacity to self-monitor, appreciation of nonverbal cues
 - e. All of the above
18. Which of the following is an exception to confidentiality in the health provider-patient relationship?
- a. A case where the provider learns the patient has had a prior child who was prenatally exposed to alcohol.
 - b. When a sexually active woman of childbearing age reveals a binge-drinking pattern and the provider also learns that she is not using contraception.
 - c. When a pregnant woman admits to her health care provider that she has a recent history of alcohol addiction.
 - d. A case where the provider learns of alcohol exposure by a pregnant patient, indicating serious bodily harm to herself and potentially her fetus.
 - e. There are never exceptions to confidentiality in health care.
19. Punitive approaches towards pregnant women who are at risk for having an alcohol-exposed pregnancy...
- a. Build trust in the practitioner/patient relationship.
 - b. Violate principals of autonomy and justice.
 - c. Have been shown to decrease the incidence rate of FASDs.
 - d. Enhance maternal-child attachments.
 - e. Provide revenue for the legal system.
20. Which of the following includes an ethical and effective response to the risks of an FASD?
- a. Recognition that addiction is an illness that requires treatment.
 - b. Understanding that women do not intend to expose their fetuses to the risks of compulsive and abusive drinking.
 - c. Consideration that forced restraint from alcohol upon a woman can contribute to mistrust of health care providers when prenatal care is needed.
 - d. Acknowledging that women who abuse alcohol during pregnancy may not be able to readily stop, nor are they morally culpable.
 - e. All of the above.

Please respond to the following questions about your professional experiences.

1. In your current position, do you...
 provide services to women of childbearing age? YES NO
 provide services to individuals who may be at risk of an FASD? YES NO

If you, in your current professional position, interact with women of childbearing age for whom alcohol screening is appropriate, please respond to the following two items.

i. What factors discourage you from screening women of childbearing age for alcohol use?

ii. What factors encourage you to screen women of childbearing age for alcohol use?

On a scale from 0 to 10 where 0 means “Not confident in my skills” and 10 means “Totally confident in my skills,” how confident are you in your skills to do the following? (circle one number per row)

	<u>Not confident in my skills</u>											<u>Totally confident in my skills</u>
Screen women for risky or hazardous drinking	0	1	2	3	4	5	6	7	8	9	10	
Educate pregnant women about the effects of alcohol on their babies	0	1	2	3	4	5	6	7	8	9	10	
Conduct brief interventions for reducing alcohol consumption	0	1	2	3	4	5	6	7	8	9	10	
Utilize resources to refer patients who need formal treatment for alcohol abuse	0	1	2	3	4	5	6	7	8	9	10	
Identify persons with possible FAS or other prenatal alcohol-related disorders	0	1	2	3	4	5	6	7	8	9	10	
Diagnose persons with possible FAS or other prenatal alcohol-related disorders	0	1	2	3	4	5	6	7	8	9	10	
Utilize resources to refer patients for diagnosis and/or treatment services	0	1	2	3	4	5	6	7	8	9	10	
Manage/coordinate the treatment of persons with FASDs	0	1	2	3	4	5	6	7	8	9	10	

Please take a moment to tell us about yourself:

<p>Are you (Circle one): Male Female Transgender</p> <p>Are you Hispanic or Latino(a)? Yes No</p> <p>How do you describe your race? (Select all that apply)</p> <p>American Indian/Alaska Native Asian Black or African American Native Hawaiian/Pacific Islander White</p> <p><input type="checkbox"/> I am the parent/caregiver of a child with FAS/FASD</p>	<p>If you are a PROFESSIONAL, please circle the one that best represents your current position:</p> <p>ALLIED HEALTH PROFESSIONAL Counselor OT/PT/SLP Psychologist Social Worker Other Allied Health Professional: Specify: _____</p> <p>DENTAL PROFESSIONAL Dental Assistant Dental Hygienist Dentist</p> <p>MEDICAL PROFESSIONAL Family Physician Internal Medicine Physician Nurse (NP, RN, LPN) OB/GYN Pediatrician Physician Assistant Psychiatrist Other Medical Professional Specify: _____</p> <p>OTHER Administrator Corrections Educator, PreK-12 Educator, PreK-12 SpED Educator, Other Specify: _____ Lawyer/Judge Public Health Scientist Other Professional: Specify: _____</p>	<p>If you are a STUDENT OR RESIDENT, please circle all that apply:</p> <p>ALLIED HEALTH STUDENT Human Services OT/PT/SLP Psychology Social Work Other Allied Health Student: Specify: _____</p> <p>HEALTH STUDENT Dental Student Medical Student Nursing Student</p> <p>MEDICAL RESIDENT Family Physician Internist OB/GYN Pediatrician Psychiatrist Other Resident Specify: _____</p> <p>OTHER STUDENT Counseling Education Education Other Student Specify: _____</p>
--	--	--