



Physicians and other health care providers with questions about diagnosis and treatment of malaria cases can call CDC's Malaria Hotline:

- Monday – Friday, 9:00 am to 5 pm, EST: call 770-488-7788 (Fax: 770-488-4206)

- Off-hours, weekends, and federal holidays: call 770-488-7100 and ask to have the malaria clinician on call paged.

Information on malaria risk, prevention, and treatment is available at:

CDC's Malaria Web site <http://www.cdc.gov/malaria>

Part II (to be complete 4 weeks after treatment)

1 prescription and over the counter medicines the patient had taken during the 2 weeks **before** starting their treatment for malaria.

1 prescription and over the counter medicines the patient had taken during the 4 weeks **after** starting their treatment for malaria.

Medicine for malaria treatment taken as prescribed? No, doses missed Yes, no doses missed Unknown

Do signs or symptoms of malaria resolve without any additional treatment within 7 days after treatment start?

Yes No Unknown

If yes, did the patient experience a recurrence of signs or symptoms of malaria during the 4 weeks after starting malaria treatment?

Yes No Unknown

Did the patient experience any adverse events within 4 weeks after receiving the malaria treatment? Yes No Unknown

Event description	Relationship to treatment suspected*	Time to Onset since treatment start	Fatal?	Life-Threatening?	Other Seriousness?***
_____	<input type="checkbox"/>	_____	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
_____	<input type="checkbox"/>	_____	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
_____	<input type="checkbox"/>	_____	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
_____	<input type="checkbox"/>	_____	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
_____	<input type="checkbox"/>	_____	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

*Relationship suspected means that a causal relationship between the treatment and an adverse event is at least a reasonable possibility, i.e., the relationship cannot be ruled out.

***Other Seriousness is defined as an event which is fatal or life-threatening, results in persistent or significant disability/incapacity, constitutes a congenital defect, is medically significant (i.e., jeopardizes the patient or may require medical or surgical intervention), or requires inpatient hospitalization or continuation of existing hospitalization