

TABLE 6B – QUALITY OF CARE INDICATORS

SECTION A: AGE CATEGORIES FOR PRENATAL PATIENTS (GRANTEES WHO PROVIDE PRENATAL CARE ONLY)				
DEMOGRAPHIC CHARACTERISTICS OF PRENATAL CARE PATIENTS				
AGE		NUMBER OF PATIENTS (a)		
1	LESS THAN 15 YEARS			
2	AGES 15-19			
3	AGES 20-24			
4	AGES 25-44			
5	AGES 45 AND OVER			
6	TOTAL PATIENTS (SUM LINES 1 – 5)			
SECTION B – TRIMESTER OF ENTRY INTO PRENATAL CARE				
TRIMESTER OF FIRST KNOWN VISIT FOR WOMEN RECEIVING PRENATAL CARE DURING REPORTING YEAR		Women Having First Visit with Grantee (a)	Women Having First Visit with Another Provider (b)	
7	First Trimester			
8	Second Trimester			
9	Third Trimester			
SECTION C – CHILDHOOD IMMUNIZATION				
CHILDHOOD IMMUNIZATION		TOTAL NUMBER OF PATIENTS WITH 3 RD BIRTHDAY DURING MEASUREMENT YEAR (a)	NUMBER CHARTS SAMPLED OR EHR TOTAL (b)	NUMBER OF PATIENTS IMMUNIZED (c)
10	MEASURE: Children who have received age appropriate vaccines who had their 3 rd birthday during measurement year (on or prior to 31 December)			
SECTION D – CERVICAL CANCER SCREENING				
PAP TESTS		TOTAL NUMBER OF FEMALE PATIENTS 24-64 YEARS OF AGE (a)	NUMBER CHARTS SAMPLED OR EHR TOTAL (b)	NUMBER OF PATIENTS TESTED (c)
11	MEASURE: Female patients aged 24-64 who received one or more Pap tests to screen for cervical cancer			
SECTION E – WEIGHT ASSESSMENT AND COUNSELING FOR CHILDREN AND ADOLESCENTS				
CHILD AND ADOLESCENT WEIGHT ASSESSMENT AND COUNSELING		TOTAL PATIENTS AGED 3 – 17 ON DECEMBER 31 (a)	NUMBER CHARTS SAMPLED OR EHR TOTAL (b)	NUMBER OF PATIENTS WITH COUNSELING AND BMI DOCUMENTED (c)
12	MEASURE: Children and adolescents aged 3 – through 17 during measurement year (on or prior to 31 December) with a BMI percentile, and counseling on nutrition and physical activity documented for the current year			

SECTION F – ADULT WEIGHT SCREENING AND FOLLOW-UP				
ADULT WEIGHT SCREENING AND FOLLOW-UP		TOTAL PATIENTS AGED 18 AND OVER (a)	NUMBER CHARTS SAMPLED OR EHR TOTAL (b)	NUMBER OF PATIENTS WITH BMI CHARTED AND FOLLOW-UP PLAN DOCUMENTED AS APPROPRIATE (c)
13	MEASURE: Patients aged 18 and over with (1) BMI charted and (2) follow-up plan documented if patients are overweight or underweight			
SECTION G – TOBACCO USE SCREENING AND CESSATION				
TOBACCO USE SCREENING AND CESSATION		TOTAL PATIENTS AGED 18 AND OVER (a)	NUMBER CHARTS SAMPLED OR EHR TOTAL (b)	NUMBER OF PATIENTS ASSESSED AND INTERVENED WITH AS APPROPRIATE (c)
14a	MEASURE: Patients screened for tobacco use AND who received cessation counseling intervention or medication if identified as a tobacco user one or more times in the measurement year or prior year			
SECTION H – ASTHMA PHARMACOLOGICAL THERAPY				
ASTHMA TREATMENT PLAN		TOTAL PATIENTS AGED 5 - 40 WITH PERSISTENT ASTHMA (a)	NUMBER CHARTS SAMPLED OR EHR TOTAL (b)	NUMBER OF PATIENTS WITH ACCEPTABLE PLAN (c)
16	MEASURE: Patients aged 5 through 40 diagnosed with persistent asthma who have an acceptable pharmacological treatment plan			
SECTION J – ISCHEMIC VASCULAR DISEASE (IVD): ASPIRIN OR ANTITHROMBOTIC THERAPY				
ASPIRIN OR OTHER ANTITHROMBOTIC THERAPY		TOTAL PATIENTS 18 AND OVER WITH IVD DIAGNOSIS OR AMI, CABG, OR PTCA PROCEDURE (a)	CHARTS SAMPLED OR EHR TOTAL (b)	NUMBER OF PATIENTS WITH ASPIRIN OR OTHER ANTITHROMBOTIC THERAPY (c)
18	MEASURE: Patients aged 18 and older with a diagnosis of IVD or AMI, CABG, or PTCA procedure with aspirin or another antithrombotic therapy			
SECTION K – COLORECTAL CANCER SCREENING				
COLORECTAL CANCER SCREENING		TOTAL PATIENTS 51 THROUGH 74 YEARS OF AGE (a)	CHARTS SAMPLED OR EHR TOTAL (b)	NUMBER OF PATIENTS WITH APPROPRIATE SCREENING FOR COLORECTAL CANCER (c)

19	MEASURE: Patients age 51 through 74 years of age during measurement year (on or prior to 31 December) with appropriate screening for colorectal cancer			
SECTION L – NEWLY IDENTIFIED HIV CASES WITH TIMELY FOLLOWUP				
NEW HIV CASES WITH TIMELY FOLLOWUP		PATIENTS FIRST DIAGNOSED WITH HIV (a)	CHARTS SAMPLED OR EHR TOTAL (b)	PATIENTS FIRST DIAGNOSED SEEN WITHIN 90 DAYS OF DIAGNOSIS (c)
20	Measure: Patients whose first ever HIV diagnosis was made by health center staff between October 1 and September 30 and who were seen for followup within 90 days of that first ever diagnosis. (See manual.)			
SECTION M – PATIENTS SCREENED FOR DEPRESSION AND FOLLOWED UP IF APPROPRIATE				
PATIENTS SCREENED FOR DEPRESSION AND FOLLOWED UP AS APPROPRIATE		TOTAL PATIENTS AGED 12 AND OVER (a)	NUMBER CHARTS SAMPLED OR EHR TOTAL (b)	NUMBER OF PATIENTS SCREENED AND FOLLOW-UP PLAN DOCUMENTED AS APPROPRIATE (c)
21	Measure: Patients aged 12 and over who were (1) screened for depression with a standardized tool and (2) had a follow-up plan documented if patients were considered depressed			