

**FDA warns OTC Constipation-Treating Sodium Phosphate Products may cause harm when exceeding the recommended dose**

Safety Announcement

Drug Safety Risk	Warning: using more than one dose in 24 hours of over-the-counter (OTC) sodium phosphate drugs to treat constipation can cause rare but serious harm to the kidneys and heart, and even death.
Facts about over-the-counter (OTC) sodium phosphate drug products	<ul style="list-style-type: none"> <li>• Marketed under the brand-name Fleet, and as store brands and generic products.</li> <li>• Used for the relief of occasional constipation.</li> <li>• Available as a solution for oral or rectal (enema) use.</li> <li>• Available as single-ingredient drug products, containing either sodium biphosphate or sodium phosphate, and as combination drug products containing both ingredients.</li> <li>• The recommended dose is a single dose given once a day for not more than 3 days.</li> </ul>
Basis for Drug Safety Risk	Side effects have resulted from taking more than the recommended dose of OTC sodium phosphate products. The side effects have resulted in serious adverse effects on organs, such as the kidneys and heart, and in some cases resulting in death. Negative side effects include severe dehydration, which is a severe loss of body water, and changes in the levels of serum electrolytes. The serum electrolytes are salts, minerals and other substances in the fluids of the body such as calcium and sodium. According to the reports, most cases of serious harm occurred with a single dose of sodium phosphate that was larger than recommended or with more than one dose in a day.
People at Higher Risk	Some individuals may be at higher risk for side effects when they take more than the recommended dose of OTC sodium phosphate. These individuals include young children; individuals older than 55; patients who are dehydrated; patients with kidney disease, bowel obstruction, or inflammation of the bowel; and patients who are using medications that may affect kidney function. These medications include diuretics or water pills; angiotensin converting enzyme inhibitors (ACEIs) and angiotensin receptor blockers (ARBs) used to treat high blood pressure; and nonsteroidal anti-inflammatory drugs (NSAIDs) such as aspirin, ibuprofen, and naproxen used to treat pain and fever.
Information for Consumers and Parents/Caregivers	<ul style="list-style-type: none"> <li>• <b>Always read and follow the directions on the Drug Facts labels</b> to find the correct dose and dosing frequency.</li> <li>• <b>Do not use more than one dose of these products in 24 hours.</b> Even if you or your children do not have a bowel movement after taking a single oral or rectal dose, do not use another dose within 24 hours. Contact a</li> </ul>

	<p>health care professional for advice.</p> <ul style="list-style-type: none"> <li>• Serious harm can occur with use of either the oral or rectal forms of OTC sodium phosphate.</li> <li>• Do not give these products rectally to children younger than 2 years.</li> <li>• Do not give these products by mouth to children 5 years and younger without first talking with a health care professional.</li> <li>• <b>Seek medical attention immediately if</b> you or your child experiences symptoms of kidney injury, and do not take another dose of the product. Symptoms of kidney injury include drowsiness; sluggishness; decreased amount of urine; or swelling of the ankles, feet, and legs.</li> <li>• Report side effects from OTC sodium phosphate drug products to the FDA MedWatch program, using the information in the "Contact FDA" box at the bottom of this page.</li> </ul>
<p>Information for Health Care Professionals</p>	<ul style="list-style-type: none"> <li>• The severity of adverse events is similar regardless of the route of sodium phosphate administration. Rectal forms of sodium phosphate drug products should not be considered safer than the oral forms.</li> <li>• Rectal sodium phosphate drug products should not be used at all in children younger than 2 years.</li> <li>• Use caution if recommending use of an oral OTC sodium phosphate drug product in children 5 years and younger.</li> <li>• Avoid exceeding the maximum recommended rectal or oral dose of sodium phosphate products for both children and adults.</li> <li>• Additional doses are not recommended within 24 hours for patients who do not have a bowel movement after taking an oral or rectal dose.</li> <li>• Avoid concomitant treatment with laxatives containing sodium phosphate.</li> <li>• Use caution when recommending these products to patients at potentially higher risk for product-related adverse events. These include those older than 55 years; patients with hypovolemia; or decreased intravascular volume; those who have baseline kidney disease, decreased bowel transit time, bowel obstruction, or active colitis; and those who are using medications that affect renal perfusion or function, such as diuretics, ACEIs, ARBs, or NSAIDs.</li> <li>• Advise patients to ensure they are adequately hydrated during product use. Assess serum electrolytes and renal function in patients who may be at higher risk for product-related adverse events, which includes those who have retained a rectal dose for more than 30 minutes, who are vomiting, or who may have signs of dehydration.</li> <li>• Report adverse events involving OTC sodium phosphate drug products to the FDA MedWatch program, using the information in the "Contact FDA" box at the bottom of this page.</li> </ul>
<p>Previous Communications on this Risk</p>	<p>FDA communicated previously about the risk of kidney injury with the use of oral sodium phosphate drug products at higher doses for bowel cleansing prior to colonoscopy or other procedures. These 2008 communications</p>

	<p>included an <a href="#">Information for Healthcare Professionals Sheet</a>, an <a href="#">FDA News Release</a>, and a <a href="#">Questions and Answers document</a>.</p>
<p>Data Summary</p>	<p>FDA reviewed the FDA Adverse Event Reporting System (FAERS) database from 1969 through 2012 and the medical literature from 1957 through August 2013 for cases describing serious adverse events associated with the oral or rectal use of over-the-counter (OTC) sodium phosphate drug products used to treat constipation. We identified 54 cases describing serious adverse events in 25 adults and in 29 children. Ten cases were reported to the FAERS database and 44 were published in the medical literature.<sup>1-39</sup> The age of the consumers ranged widely from 8 days to 97 years, but most cases involved older adults and children younger than 5 years. Nearly two-thirds of the adults (approximately 66%) and nearly half of the children (approximately 50%) in whom adverse events were reported had one or more of the following:</p> <ul style="list-style-type: none"> <li>• Dehydration, kidney disease, acute colitis, or delayed bowel emptying</li> <li>• Concomitant use of drugs that act on renal function, including diuretics, ACEIs, ARBs, and NSAIDs</li> </ul> <p>All reports of serious outcomes were characterized by dehydration and/or electrolyte disturbances with associated complications such as acute kidney injury and death. The severity of adverse events was similar regardless of whether the products were administered orally or rectally. The predominant electrolyte disturbances were hyperphosphatemia, hypocalcemia, and hypernatremia. Nearly half (12/25) of adult cases and 3% (1/29) of pediatric cases reported a fatal outcome. The remaining non-fatal cases were life-threatening in more than two-thirds of affected adults (more than 66%) and in all of the affected children (100%). These included severe deterioration in respiratory status, mental status, and heart function. Four adults required dialysis and two underwent surgery for suspected bowel perforation. Three children required dialysis, including two younger than 2 years. Two children developed abdominal distention significant enough to require surgery. One child younger than 2 years had residual neurological defects.</p> <p>In the 50 cases for which the administered dose was reported (27 pediatric, 23 adult), serious adverse events occurred in patients who took a dose in excess of the labeled amount, including 60% (16/27) of the pediatric cases and 70% (16/23) of the adult cases. Adults and pediatric patients who overdosed either received a single dose that contained a greater amount of sodium phosphate than recommended on the label, or they received the product more frequently than recommended on the label. The duration of use in the majority of the overdose cases was 1 to 2 days. Forty percent (11/27) of the pediatric cases for which the dose was reported occurred in young children for whom FDA has not proposed a safe and effective dose. Nine of these 11 cases (approximately 82%) were children younger</p>

	<p>than 2 years who received a rectal sodium phosphate product, and two were in children younger than 5 years who received an oral product. These children received doses comparable to those recommended on the label for use in adults or older children. Seven adults developed serious adverse events despite taking dosages that did not exceed recommendations on the label; however, all had one or more of the above conditions, potentially increasing their adverse event risk.</p>
What is FDA?	<p>FDA includes pharmacists, doctors, nurses, researchers and other health professionals. We also work with external experts. We protect public health by assuring drugs are safe and effective.</p>
References	<ol style="list-style-type: none"> <li>1. Moseley PK, Segar WE. Fluid and serum electrolyte disturbances as a complication of enemas in Hirschsprung's disease. <i>Am J Dis Child</i> 1968;115:714-8.</li> <li>2. Young JF, Brooke BN. Enema shock in Hirschsprung's disease. <i>Dis Colon Rectum</i> 1968;11:391-5.</li> <li>3. Chesney RW, Haughton PB. Tetany following phosphate enemas in chronic renal disease. <i>Am J Dis Child</i> 1974;127:584-6.</li> <li>4. Oxnard SC, O'Bell J, Grupe WE. Severe tetany in an azotemic child related to a sodium phosphate enema. <i>Pediatrics</i> 1974;53:105-6.</li> <li>5. Swerdlow DB, Labow S, D'Anna J. Tetany and enemas: report of a case. <i>Dis Colon Rectum</i> 1974;17:786-7.</li> <li>6. Zipser RD, Bischel MD, Abrams DE. Hypocalcemic tetany due to sodium phosphate ingestion in acute renal failure. <i>Nephron</i> 1975;14:378-81.</li> <li>7. Davis RF, Eichner JM, Bleyer WA, Okamoto G. Hypocalcemia, hyperphosphatemia, and dehydration following a single hypertonic phosphate enema. <i>J Pediatr</i> 1977;90:484-5.</li> <li>8. Loughnan P, Mullins GC. Brain damage following a hypertonic phosphate enema. <i>Am J Dis Child</i> 1977;131:1032.</li> <li>9. Forman J, Baluarte HJ, Gruskin AB. Hypokalemia after hypertonic phosphate enemas. <i>J Pediatr</i> 1979;94:149-51.</li> <li>10. Biberstein M, Parker BA. Enema-induced hyperphosphatemia. <i>Am J Med</i> 1985;79:645-6.</li> <li>11. Haskell LP. Hypocalcaemic tetany induced by hypertonic-phosphate</li> </ol>

enema. *Lancet* 1985;2:1433.

12. Spinrad S, Sztern M, Grosskopf Y, Graff E, Blum I. Treating constipation with phosphate enema: an unnecessary risk. *Isr J Med Sci* 1989;25:237-8.

13. Wason S, Tiller T, Cunha C. Severe hyperphosphatemia, hypocalcemia, acidosis, and shock in a 5-month-old child following the administration of an adult Fleet enema. *Ann Emerg Med* 1989;18:696-700.

14. Edmondson S, Almquist TD. Iatrogenic hypocalcemic tetany. *Ann Emerg Med* 1990;19:938-40.

15. McCabe M, Sibert JR, Routledge PA. Phosphate enemas in childhood: cause for concern. *BMJ* 1991;302:1074.

16. Korzets A, Dicker D, Chaimoff C, Zevin D. Life-threatening hyperphosphatemia and hypocalcemic tetany following the use of fleet enemas. *J Am Geriatr Soc* 1992;40:620-1.

17. Hunter MF, Ashton MR, Griffiths DM, Ilangovan P, Roberts JP, Walker V. Hyperphosphataemia after enemas in childhood: prevention and treatment. *Arch Dis Child* 1993;68:233-4.

18. Craig JC, Hodson EM, Martin HC. Phosphate enema poisoning in children. *Med J Aust* 1994;160:347-51.

19. Selva O'Callaghan A, San Jose A, Simeon CP, Arruche M, Galicia M, Benavente V, et al. Fatal hypocalcemia from sodium phosphate enemas. *J Am Geriatr Soc* 1995;43:1447-8.

20. Knobel B, Petchenko P. Hyperphosphatemic hypocalcemic coma caused by hypertonic sodium phosphate (fleet) enema intoxication. *J Clin Gastroenterol* 1996;23:217-9.

21. Helikson MA, Parham WA, Tobias JD. Hypocalcemia and hyperphosphatemia after phosphate enema use in a child. *J Pediatr Surg* 1997;32:1244-6.

22. Szabo A, Eitan A, Zveibil F, Weiss M, Waksman I, Makhoul N. Life-threatening electrolyte abnormalities induced by hypertonic phosphate enema intoxication. *Clinical Intensive Care* 1999;10:21-3.

23. Nir-Paz R, Cohen R, Haviv YS. Acute hyperphosphatemia caused by sodium phosphate enema in a patient with liver dysfunction and chronic

renal failure. *Ren Fail* 1999;21:541-4.

24. Ismail EA, Al-Mutairi G, Al-Anzy H. A fatal small dose of phosphate enema in a young child with no renal or gastrointestinal abnormality. *J Pediatr Gastroenterol Nutr* 2000;30:220-1.

25. Walton DM, Thomas DC, Aly HZ, Short BL. Morbid hypocalcemia associated with phosphate enema in a six-week-old infant. *Pediatrics* 2000;106:E37.

26. Melvin JD, Watts RG. Severe hypophosphatemia: a rare cause of intravascular hemolysis. *Am J Hematol* 2002;69:223-4.

27. Marraffa JM, Hui A, Stork CM. Severe hyperphosphatemia and hypocalcemia following the rectal administration of a phosphate-containing Fleet pediatric enema. *Pediatr Emerg Care* 2004;20:453-6.

28. Butani L. Life-threatening hyperphosphatemia and hypocalcemia from inappropriate use of Fleet enemas. *Clin Pediatr (Phila)* 2005;44:93.

29. Farah R. Fatal acute sodium phosphate enemas intoxication. *Acta Gastroenterol Belg* 2005;68:392-3.

30. Domico MB, Huynh V, Anand SK, Mink R. Severe hyperphosphatemia and hypocalcemic tetany after oral laxative administration in a 3-month-old infant. *Pediatrics* 2006;118:e1580-3.

31. Eckstein J, Savic S, Eugster T, Pargger H, Gürke L, Pechula M, et al. Extensive calcifications induced by hyperphosphatemia caused by phosphate-based enema in a patient after kidney transplantation. *Nephrol Dial Transplant* 2006;21:2013-6.

32. Hay E, Boksenbojm P, Esther P. Case report: Fleet enema in the renal patient. *Israeli Journal of Emergency Medicine* 2006;6:49-51.

33. Hebbar K, Fortenberry JD, Parks JS. Severe hypocalcemic tetany and respiratory failure in an infant given oral phosphate soda. *Pediatr Emerg Care* 2006;22:118-20.

34. Hsu HJ, Wu MS. Extreme hyperphosphatemia and hypocalcemic coma associated with phosphate enema. *Intern Med* 2008;47:643-6.

35. Biebl A, Grillenberger A, Schmitt K. Enema-induced severe hyperphosphatemia in children. *Eur J Pediatr* 2009;168:111-2.

36. Viel G, Cecchetto G, Fabbri LD, Furlan C, Ferrara SD, Montisci M. Forensic application of ESEM and XRF-EDS techniques to a fatal case of sodium phosphate enema intoxication. *Int J Legal Med* 2009;123:345-50.

37. Kostic D, Rodrigues AB, Leal A, Metran C, Nagaiassu M, Watanabe A, et al. Flow-through peritoneal dialysis in neonatal enema-induced hyperphosphatemia. *Pediatr Nephrol* 2010;25:2183-6.

38. Ladenhauf HN, Stundner O, Spreitzhofer F, Deluggi S. Severe hyperphosphatemia after administration of sodium-phosphate containing laxatives in children: case series and systematic review of literature. *Pediatr Surg Int* 2012;28:805-14.

39. Szoke D, Dolci A, Genderini A, Panteghini M. Fatal electrolyte abnormalities following enema administration. *Clin Chem* 2012;58:1515-8.