Household ID# \_\_\_\_\_

## **Instructions:**

If child (who is participating in this study) develops at least 3 of the following: **fever, stuffy/runny nose, cough, sore throat, body aches or tiredness**, for more than 24 hours --- please do the following:

1. Swab the nose and throat of the child using the directions we gave you when we dropped off the swabs.







- a. Refrigerate the swabs
- b. Call the study coordinator at ###.######



2. Complete the Illness Checklist (next page), and keep an Illness Log



Public reporting burden of this collection of information is estimated to average 5 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Information Collection Review Office, 1600 Clifton Road NE, MS D-74, Atlanta, Georgia 30333; ATTN: PRA (0920-0906)

## ILLNESS CHECKLIST

		Symptom Checklist				
		INSTRUCTIONS: Check box for all symptoms experienced. Check "none" if the symptom is absent.				
			severity rating (see footnote*)			
YES (this symptom developed)	NO (this symptom did not develop)	Symptoms	mild 1	moderate 2	severe 3	
General						
		Fever (o) temp, if known				
		chills				
		weakness/tired				
Lungs						
		coughing				
		wheezing				
		difficulty breathing				
		Stuffy/ runny nose				
		Sore throat				
		other:				
Arms, legs, back, neck						
		muscle aches				
		joint pain				
		headache				
		other:				

NOTE: To be given to study technician along with nose/throat swabs

Date first symptom (checked above) started: _	/	_/	_ (mm/	dd/yyyy)
Date that mother/primary caregiver swabbed ch	ild:	/	_/	(mm/dd/yyyy)

\*\*\*Nose and throat swabs should be done within 24-36 hours from the beginning of symptoms\*\*\*

\*\*\*Do not swab child's nose/throat after 5 days of the beginning of symptoms\*\*\*

## \*Rating

- 1. mild child notices a difference but still able to carry out usual everyday activities (play, school, hobbies)
- 2. moderate difference noticeable by others; some difficulty and loss in carrying out usual everyday activities (play, school, hobbies)
- 3. severe noticeable by others; unable to carry out usual everyday activities (play, school, hobbies)

## **ILLNESS LOG**



Did the child's asthma get worse during the illness?	Yes	No			
Did the child become so ill that he/she had to see the doctor?	Yes	No			
Did doctor prescribe Tamiflu or Relenza?	Yes	No			
Did doctor prescribe antibiotics?	Yes	No			
Did the child become so ill that he/she had to be admitted to a hospital for overnight care?					
	Yes	No			
Date when the child was well enough to do usual activities://		(mm/dd/yyyy)			