Zika Prevention Take a Community - Screen Shot

https://www.cdc.gov/zika/pdfs/protectcommunity-onepager.pdf



Mosquito Control: Do Your Part - Screen Shot

Public reporting burden of this collection of information is estimated to average 1.5 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Reports Clearance Officer; 1600 Clifton Road NE, MS D-74, Atlanta, Georgia 30333; Attn: OMB-PRA (0920-0572)

https://www.cdc.gov/zika/pdfs/aerial-spraying-community.pdf



MOSQUITO CONTROL: DO YOUR PART

Effective mosquito control requires an integrated approach that uses a combination of methods. Everyone should do their part to help control mosquitoes and protect pregnant women from Zika.

State or Local Government



- Apply insecticides to control mosquitoes. This may include truck, plane, or handheld spraying.
- Treat water with larvicides to kill larvae that hatch from eggs.
- Track mosquitoes and the viruses they carry.

Household

- Remove standing water in and around your home.
- Use, install, or repair window and door screens.
- Keep windows and doors shut.
- Use air conditioning when available.

Her

- Use insect repellent.
- Wear protective clothing.
- Use condoms with a partner who has been in an area with Zika.
- Talk to her doctor.



Community

- Remove standing water around your community.
- Clean up trash, containers, and tires that can hold water.
- Use larvicides in water that can't be removed and won't be used for drinking.



You

- · Wear insect repellent.
- Cover up with long-sleeved shirts and long pants.
- Treat clothes with permethrin or buy treated clothes and gear.
- Use condoms with a partner who has been in an area with Zika.



U.S. Department of Health and Human Services Centers for Disease Control and Prevention

CS269893-A October 25, 2016

Public reporting burden of this collection of information is estimated to average 1.5 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Reports Clearance Officer; 1600 Clifton Road NE, MS D-74, Atlanta, Georgia 30333; Attn: OMB-PRA (0920-0572)

Public reporting burden of this collection of information is estimated to average 1.5 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Reports Clearance Officer; 1600 Clifton Road NE, MS D-74, Atlanta, Georgia 30333; Attn: OMB-PRA (0920-0572)