Survey	Years Conducted	Sample size	Mode	Study Design	Primary purpose	HINTS communication core	Tobacco Message Exposure Source/Frequency/Reaction	Tobacco use phenotypes	Menthol	Product constituents	Risk perceptions about tobacco products	Attitudes and beliefs about product regulation	Tobacco product claims and labels
2014-2015 TUS-CPS	1992-1993, 1995-1996, 1998-1999, 2000, 2001-2002, 2003, 2006-2007, and 2010-11	~ 240,000	64% by telephone and 36% in person	Cross-sectional	NCI-sponsored survey of tobacco use administered as part of the US Census Bureau's Current Population Survey	No	No	Yes	Yes	No	No	No	No
2014 NHIS	Annually, since 1957	~ 35,000 households	In-person	Cross-sectional	NHIS data on a broad range of health topics are collected through personal household interviews. The U.S. Census Bureau is the data collection agent.	No	No	Yes	No	No	No	No	No
2013 BRFSS	Annually, since 1984	More than 500,000 adults are interviewed each year	Telephone	Cross-sectional	BRFSS collects data via telephone in all 50 states plus DC on health-related risk behaviors and events (including tobacco use), chronic health conditions, and use of preventive services.	No	No	Yes	No	No	No	No	No
2013-2014 ATS-CDC / NATS	33 CDC supported ATS's were conducted in 19 states from 2003-2007; 2009 - Present NATS (yearly)	~75,000	RDD design telephone survey (Final questionnaires were developed by each state)	Cross-sectional	NATS seeks to determine tobacco use prevalence and the factors promoting and impeding tobacco use among adults in a nationally representative sample of adults and will provide time-critical monitoring data needed to evaluate the effectiveness of new regulatory authorities given to FDA under the Family Smoking Prevention and Tobacco Control Act (the Tobacco Control Act)	No	Yes	Yes	Yes	Yes	Yes	No	No
2013 NSDUH	Periodically since 1972 and annually since 1991	Approximately 70,000 individuals, age 12 and older	In-person interviews with sample persons	Cross-sectional	The NSDUH provides data on prevalence and correlates of substance use, serious mental illness, related problems, and treatment in the civilian population aged 12 or older in the U.S.	No	No	Yes	Yes	No	No	No	No
2013 NYTS and 2014 NYTS	1999, 2000, 2002, 2004, 2006, 2009, 2011, 2012	18,406 in 2013	In school, self-administered survey via pencil	Cross-sectional	The NYTS provides prevalence estimates for selected tobacco use behaviors among middle school and high school students regarding tobacco.	No	Yes	Yes	Yes	No	Yes	No	No
2015 YRBS	Every odd year since 1991	~15,000	School-based, self-administered questionnaire	Cross-sectional	The Youth Risk Behavior Surveillance System (YRBSS) monitors priority health-risk behaviors and the prevalence of obesity and asthma among youth and young adults.	No	No	Yes	No	No	No	No	No
2012 MTF	Annually since 1975	~50,000	School-based, self-administered questionnaire with mailed follow up	Cross-sectional with annual follow-up mailed questionnaires among a sample of each graduating class.	MTF is an ongoing study of the behaviors, attitudes, and values of American secondary school students, college students, and young adults.	No	No	Yes	No	No	Yes	No	No
2011 YTS	Periodically since 1999 in participating states	~27,000, varies by year and state	School-based survey of students in grades 6 through 12.	Cross-sectional	The YTS is intended to enhance the capacity of agencies and organizations to design, implement, and evaluate tobacco prevention and control programs.	No	Yes	Yes	Yes	No	Yes	No	No
2013-2014 NHANES	A series of surveys since 1959; 1999-2000, 2001-02, 2003-04, 2005-06, 2007-08, 2009-10, 2011-12, 2013-14	~5,000	In-person inteviews and physician examinations	Cross-sectional	NHANES findings are used to determine the prevalence of major diseases and risk factors for diseases, assess nutritional status and its association with health promotion and disease prevention.	No	No	Yes	Yes	No	No	No	No
2013 Adult PATH	Baseline data collection began in September 2013 and followed annually for at least two additional data collection waves	~60,000	In-person	Longitudinal	PATH findings will be used to inform FDA's regulatory authority over tobacco products.	No	Yes	Yes	Yes	Yes	Yes	No	No
HINTS	2003, 2005, 2007, yearly from 2011-2014 (in progress)	~3,500	Mailed survey; additional RDD telephone sample in 2005	Cross-sectional	NCI-sponsored survey that collects nationally representative data routinely about adults' access to and use of health information in the broad public communication environment. The FDA Cycle will collect data relevant to the Family Smoking Prevention and Tobacco Control Act. Data will be used to inform FDA's tobacco prevention campaigns and tobacco control outreach and education.	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes

| | | |