

Form Approved  
OMB No. 0920-XXXX  
Exp. Date xx/xx/20xx

***Brief Questionnaire for Nursing Organization Memberships (Online)***

You are invited to participate in this survey for members of nursing organizations. The purpose of the survey is to gather information about nurses' practice behaviors related to FASD prevention and alcohol screening and brief intervention. Your participation is important in order to understand effects of the CDC FASD Project as well as to inform future activities of the Project.

This survey will take approximately 10 minutes to complete. Your participation is voluntary and responses will be kept secure. There are no costs associated with participating nor will you directly benefit from participation.

CDC estimates the average public reporting burden for this collection of information as 10 minutes per response, including the time for reviewing instructions, searching existing data/information sources, gathering and maintaining the data/information needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Information Collection Review Office, 1600 Clifton Road NE, MS D-74, Atlanta, Georgia 30333; ATTN: PRA (0920-XXXX).

How often do you talk to **your patients** about their alcohol use?

Never

Occasionally

About half the time

Frequently

Always

My practice has a protocol in place to screen all patients for their alcohol use.

Yes

No

Don't know

Not applicable

What does initial patient screening for alcohol use consist of in your practice?

Informal questions (e.g., Do you drink? How much?)

Formal screening questions/instruments (e.g., AUDIT, CAGE, CRAFFT)

Don't know

Who generally conducts the screen for alcohol use?

Nurse (including nurse practitioner)

Social worker

Behavioral health specialist

Psychologist

Physician

Medical assistant

Other

In your practice, is screening for alcohol use followed by some type of intervention?

No, there is no education or intervention done with the initial screening.

Yes, all patients are given educational materials/information on "safe" levels of alcohol and health risks associated with consuming too much alcohol.

Yes, patients who screen positive for risky alcohol use are asked follow-up questions and provided brief counseling.

Yes, patients who screen positive for risky alcohol use are asked follow-up questions and provided with additional resource (e.g., a list of treatment and/or counseling services in the community).

Who generally conducts intervention?

Physician

Physician's assistant

Psychologist

Social worker

Behavioral health specialist

Nurse (including nurse practitioner)

Other (please specify)

Please indicate the extent to which you disagree or agree with each of the following statements.

	<b>Strongly Disagree</b>	<b>Disagree</b>	<b>Neither Agree nor Disagree</b>	<b>Agree</b>	<b>Strongly Agree</b>
It is important to routinely screen all patients/clients for alcohol use.	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Screening a person for alcohol use confers a stigma to the person being screened.	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
It is important to screen all pregnant women for alcohol use.	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>



Please indicate the extent to which you disagree or agree with each of the following statements.

	<b>Strongly Disagree</b>	<b>Disagree</b>	<b>Neither Agree nor Disagree</b>	<b>Agree</b>	<b>Strongly Agree</b>
It is important to screen all women of reproductive age for alcohol use.	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
It is important to educate women of reproductive age, including those who are pregnant, about the effects of alcohol on a developing fetus	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
It is important to inquire about and document potential prenatal exposure for all pediatric patients	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Diagnosis of one of the FASDs may confer a stigma to a child and/or his or her family	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

On a scale from 1 to 5, where 1 means "not confident in my skills" and 5 means "totally confident in my skills", how confident are you in your skills to do the following? (Select one answer per row).

	<b>1. Not at all confident in my skills</b>	<b>2. Slightly confident in my skills</b>	<b>3. Moderately confident in my skills</b>	<b>4. Very confident in my skills</b>	<b>5. Totally confident in my skills</b>
Asking women, including pregnant women, about their alcohol use	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Having a conversation with patients/clients who indicate risky alcohol use	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Educating women of childbearing age, including those who are pregnant, about the effects of alcohol on a developing baby	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Conducting brief interventions for reducing alcohol use	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Utilizing resources to refer patients/clients who need formal treatment for alcohol abuse	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

	<b>1. Not at all confident in my skills</b>	<b>2. Slightly confident in my skills</b>	<b>3. Moderately confident in my skills</b>	<b>4. Very confident in my skills</b>	<b>5. Totally confident in my skills</b>
Inquiring about potential prenatal alcohol exposure for my patients/clients	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Identifying persons who may have one of the FASDs	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Diagnosing persons who may have one of the FASDs	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Referring patients/clients for diagnosis and/or treatment services for an FASD or alcohol use disorder	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Managing/coordinating the treatment and care of persons who have one of the FASDs	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

We thank you for your time spent taking this survey.  
Your response has been recorded.

Powered by Qualtrics