	Report Abuse	
Page 1 / 2		m = 100

Create your own, FREE ONLINE SURVEY

## Social Work & Family Physicians 6-Month Follow-up Survey

Form Approved

OMB No. 0920-XXXX

Exp. Date xx/xx/20xx

CDC estimates the average public reporting burden for this collection of information as 8 minutes per response, including the time for reviewing instructions, searching existing data/information sources, gathering and maintaining the data/information needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Information Collection Review Office, 1600 Clifton Road NE, MS D-74, Atlanta, Georgia 30333; ATTN: PRA (0920-XXXXX).

Instructions: Please answer the questions below. Your responses will be kept secure, and will be summarized only in aggregate with those of other respondents. Individual, identifiable responses will NOT be shared.

My practice has a protocol to screen all patients/clients for risky alcohol use.
○ Yes
○ No
O Don't know
<ul> <li>Does not apply to my practice setting</li> </ul>
IF YES TO Q1 CONTINUE, ELSE GO TO Q3.
2 What ages do you screen for risky alcohol use?
O to
<ul> <li>In your current position, do you provide services to women of childbearing age?</li> <li>Yes</li> <li>No</li> </ul>
IF NO, GO TO Q6

4 Fetal alcohol spectrum disord	ders (FASDs)	are:			
<ul> <li>Disorders a pregnant wor</li> </ul>	man experien	ces when sh	e drinks alco	phol	
O Disorders that affect the a	ability of a pre	gnant woma	n who drinks	s alcohol to go	full term
<ul> <li>Physical disroders that af</li> </ul>	fect a fetus w	hen a pregn	ant woman o	Irinks alcohol	
<ul> <li>The range of effects that alcohol</li> </ul>	can occur in a	an individual	who was ex	posed prenatal	ly to
5 What sources of information of	do you typical	lly consult for	rinformation	about FASDs?	?
○ Colleagues					
Media     ME/CELL Seminare					
CME/CEU Seminars	i a al atala a a a a				
Medical Journals/Electron	iic databases				
Other (Please specificy)					
6 How often do you talk to your parents/caregivers about their  Never  Occaisionally  About half the time  Frequently  Always  Not applicable  7 If you talk to your patients/clies  to	r alcohol use? nts, what age	s do you incl	ude?		now
important is each of the follow	- Ti		Madandali	Was Inc. 4	F
	Not important	Somewhat Important	Moderately Important	Very Important	Extremely Important
a. Discussing patient/clients' alcohol use could help reduce risky drinking.	0	0	0	0	0
b. I am too busy to have time to talk to patients/clients about their alcohol use.	0	О	0	0	0
c. Discussing their alcohol use could improve my patient/clients' health.	0	0	0	0	0
d. Helping patients/clients overcome problem drinking creates a stronger doctor-patient relationship.	0	0	0	0	0
	0	0	0	0	0

<ul> <li>e. I do not have adequate training to dea with patients/clients who are at-risk drinkers.</li> </ul>	al									
f. Discussing their alcohol use could improve my patients'/clients' work productivity.	C	)	(	0		0		0		0
g. Discussing alcohol issues is uncomfortable for me.	C	)	0		0		0			0
h. Discussing their alcohol use could improve patients'/clients' family and personal relationships	0		0		0			0		0
i. I will upset my patients/clients if I talk to them about their alcohol use.	° O		0			0		0		0
j. My talking with a patient/client about alcohol use is unlikely to make a difference in his/her drinking.	С		(	)		0		0		0
9 Rate the importance of providuse.	ling cou	nselir	ig to yo	ur pat	ients/	clients	on ris	ky alco	hol	
	1. Not at all important	2	3	4	5	6	7	8	9	10. Extremely important
a. Pregnant women.	0	0	0	0	0	0	0	0	0	0
b. Mothers after giving birth.	0	0	0	0	0	0	0	0	0	0
c. Women of childbearing age.	0	0	0	0	0	0	0	0	0	0
Rate your confidence in provuse.	iding co	unse	ling to y	our p	atient	s/clier	nts on r	isky alc	ohol	
	Not at all confident	2	3	4	i)	5	7	8	9	10. Extremely confident
a. Pregnant women.	0	0	0	С	)	0	0	0	0	0
b. Mothers after giving birth.	0	0	0	С	)	0	0	0	0	0
c. Women of childbearing age.	0	0	0	C	)	0	0	0	0	0
Rate the importance of discurage.	ssing th	e follo	owing to	opics v	with a	ll wom	en of o	childbea	aring	
	1. Not at all important	2	3	4	5	6	7	8	9	10. Extremely important
a. Contraception/family planning	0	0	0	0	0	0	0	0	0	0
b. Risky drinking	0	0	0	0	0	0	0	0	0	0

c. Risks of alcohol use during pregnancy	0	0	0	0	0	0	0	0	0	0
12 Rate the importance of discr	ussing th	ne foll	owing	topics	with w	omen	who a	re likel	y to	
conceive in the near future.	1. Not at all important	2	3	4	5	6	7	8	9	10. Extremel importan
d. Risky drinking	0	0	0	0	0	0	0	0	0	0
e. Risks of alcohol use during pregnancy	0	0	0	0	0	0	0	0	0	0
Rate your confidence in disc	cussing t	he fol	lowing	topics	with a	all wom	en of	childbe	earing	
	Not at     all     confident	2	3	4	5	6	7	8	9	10. Extremely confident
a. Contraception/family planning	0	0	0	0	0	0	0	0	0	0
b. Risky drinking	0	0	0	0	0	0	0	0	0	0
c. Risks of alcohol use during pregnancy	0	0	0	0	0	0	0	0	0	0
Rate your confidence in disci	ussing the state of the state o	ne foli	lowing 3	topics 4	with w	omen 6	likely 1	0 cond 8	eive 9	10. Extremely confident
d. Risky drinking	0	0	0	0	0	0	0	0	0	0
e. Risks of alcohol use during pregnancy	0	0	0	0	0	0	0	0	0	0
15 Which one of these is NOT c  12 ounces of beer  12 ounces of wine cooler  5 ounces of wine  3 ounces of vodka  All of the above are standar			tandar	d drink	?					
Risky drinking for non-pregnal standard drinks in a day or m  2, 6  3, 7  3, 8  4, 7										