



Attachment 9: References

References

Albert P. Longitudinal data analysis (repeated measures) in clinical trials. *Statistics in Medicine*. 1999; 18:1707-1732.

Bureau of Labor Statistics, U.S. Department of Labor. News release, August 18, 2015: Employment and Unemployment Among Youth – Summer 2015. <http://www.bls.gov/news.release/pdf/youth.pdf> , Accessed 2/09/16.

Bureau of Labor Statistics, U.S. Department of Labor. News release, January 22, 2016: Usual Weekly Earnings of Wage and Salary Workers Fourth Quarter, 2015. <http://www.bls.gov/news.release/pdf/wkyeng.pdf>, Accessed 2/09/16.

Buuren SV, Groothuis-Oudshoorn K: MICE: Multivariate imputation by chained equations in R. *J Stat Software*. 2011; 45:67.

Currie C Nic Gabhainn S, Godeau E, et al (eds.) Young people's health in context: International report from the HBSC 2001/02 survey. Health Policy for Children and Adolescents, No.4. 2004; WHO Regional Office for Europe, Copenhagen.

Currie C, Nic Gabhainn S, Godeau E, et al. (eds.) () Inequalities in young people's health: HBSC international report from the 2005/06 Survey. Health Policy for Children and Adolescents, No. 5. 2008; WHO Regional Office for Europe, Copenhagen, Denmark.

David, MC, Ware, RS. Meta-analysis of randomized controlled trials supports the use of incentives for inducing response to electronic health surveys. *Journal of Clinical Epidemiology*. 2014; 67:1210-1221.

Davis, E, Demby, H, Jenner, LW, Gregory, A, Broussard, M. Adapting an evidence-based model to retain adolescent study participants in longitudinal research. *Evaluation and Program Planning*. 2016; 54:102-111.

Drewnowski, KL, Harrell JS, Thompson D, et al. Recruitment and retention strategies and methods in the HEALTHY Study. *International Journal of Obesity*. 2009; 33: S21-S28.

Diggle P, Heagerty P, Liang KY, Zeger, S. *Analysis of longitudinal data*. Oxford University Press; 2002.

Expert Panel on Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents: Summary Report. *Pediatrics* 128: Supplement 5, 2011.

Federal Interagency Forum on Child and Family Statistics. *America's Children: Key National Indicators of Well-Being*. 2015; Washington, DC: U.S. Government Printing Office.

Goodman E, Daniels SR, Meigs JB, Dolan LM. Instability in the diagnosis of metabolic syndrome in adolescents. *Circulation*. 2007; 115:16-22.

Grunbaum JS, Kann L., Kinchen S, Ross J, Hawkins J, Lowry, R, Harris WA, McManus T, Chyen D, Colins J. Youth risk behavior surveillance- United States, 2003. *Morbidity and Mortality Weekly Report. Surveillance Summaries*. 2004; 53:1-96.

Hanna KM, Scott, LL, Schmidt, KK. Retention strategies in longitudinal studies with emerging adults. *Clinical Nurse Specialist*. 2014; 28:41-45.

Johnston, LD, O'Malley PM, Bachman JG, Schulenberg JE. *Monitoring the future national survey results on drug use 1975-2006: Volume 1, secondary school students*. NIH National Institute on Drug Abuse , 2007. NIH Publication No. 07-6205

Krieger N. Why epidemiologist cannot afford to ignore poverty. *Epidemiology*. 2007; 18:658-663.

Lipsky LM, Haynie DL, Liu D, Chaurasia A, Gee B, Li, K, Iannotti, RJ, Simons-Morton B. Trajectories of eating behaviors in a nationally representative cohort of U.S. adolescent during the transition to young adulthood. *International Journal of Behavioral Nutrition and Physical Activity*. 2015; 12:138. Doi 10.1186/s12966-015-0298-x

Little RJA. Pattern-mixture models for multivariate incomplete data. *Journal of the American Statistical Association*. 1993; 88:125-134.

Little RJA, Rubin DB: *Statistical Analysis with Missing Data*: Wiley, 2002.

Martinson BC, Lazovich D, Lando HA, Perry CL, McGovern PG, Boyle, RG. Effectiveness of monetary incentive for recruiting adolescents to an intervention trial to reduce smoking. *Preventive Medicine*. 2000; 31:706-13.

Morrison TC, Wahlgren D., Hovell MF, Zakarian J, Burkham-Kreitner S, Hofstetter CR, Slyment DJ, Keating K, Russos S, Jones JA. Tracking and follow-up of 16,915 adolescents: Minimizing attrition bias. *Controlled Clinical Trials*. 1997;18:383-396.

Muthen, Bengt. Latent variable mixture modeling. *New developments and techniques in structural equation modeling*. Mahwah, NJ: Lawrence Erlbaum Associates, Inc. 2001; 1-33.

NHANES: https://www.cdc.gov/Nchs/Nhanes/2013-2014/DEMO_H.htm#DMDMARTL

Ronbinson, KA, Dinglas, VD, Sukrithan, V, Yalamanchilli, R, Mendez-Tellez, PA, Dinnison_Himmelfarb, C, Needham, DM. Updated systematic review identifies substantial number of retention strategies: using more strategies retains more study participants. *Journal Of Clinical Epidemiology*. 2015; 68:1481-1487.

Schoeppe, S, Oliver, M, Badland, HM, Burke, M, Duncan, MJ. Recruitment and retention of children in behavioral health risk factor studies: REACH strategies. *International Journal of Behavioral Medicine*. 2014; 21:794-803.

Simons-Morton B, Haynie D, Li D, Chaurasia A, Li K, Hingson R.. The effect of residential, school, work status and social influence on the prevalence of alcohol use among emerging adults. *Journal of Studies on Alcohol and Drugs*. (in press)

Youth Risk Behavior Survey (YRBS): Centers for Disease Control and Prevention. Youth Risk Behavior Survey. Available at: www.cdc.gov/yrbs.

Wu MC, Carrol RJ. Estimation and comparison of changes in the presence of informative right censoring by modeling the censoring process. *Biometrics*. 1988;44:175-188.