

STD Treatment Guidelines Complete Long-Term Evaluation

Public reporting burden of this collection of information is estimated to average 5 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Reports Clearance Officer; 1600 Clifton Road NE, MS D-74, Atlanta, Georgia 30333; ATTN: PRA (0920-0995).

1. I am using what I learned in this training in my work.

strongly disagree strongly agree NA

2. If you have not used what you learned, please explain why not.

3. Did you make a change in your practice, worksite setting, or job as a result of this training?

- Yes
 No
 Not applicable to my job or patients
 I already use these practices

Other reason (please specify)

4. If you made a change, what change did you make?

5. As a result of this training, did you share what you learned with any of the following? (select all that apply)

- Supervisor
 Colleagues/co-workers
 Policy makers
 Community

Other (please specify)

6. Did any of these factors MAKE IT HARDER for you to apply the STD practices recommended in the training? (select all that apply)

- lack of time with patients
 more important patient concerns
 cost/lack of reimbursement
 policies where I work
 resistance to change by supervisor or colleagues
 lack of equipment or supplies
 no opportunity to apply practices
 nothing interfered

Other (please specify)

7. Did any of these factors HELP you apply the STD practices recommended in the training? (select all that apply)

- reimbursement or other financial incentive
 support of supervisor and/or colleagues
 standing orders
 reminder in chart
 convenient supplies
 posted patient instructions for obtaining specimens
 nothing specific helped

Other (please specify)

8. Do you use the CDC STD Treatment Guidelines to guide the care of your patients/clients?

- No, I am not aware of the Guidelines
- I am aware of the Guidelines but do not use them
- I use the Guidelines occasionally
- I use the Guidelines consistently
- I use another source to guide my STD care (please specify)

9. As a result of the information presented did you download the CDC STD Treatment Guidelines app?

- Yes
- No
- Not applicable
- I was already doing this

10. As a result of the information presented did you send a consult to the STD Clinical Consultation Network? www.stdccn.org

- Yes
- No
- Not applicable
- I was already doing this

11. As a result of the information presented did you recommend rescreening in 3 months following a gonorrhea, chlamydia or trichomonas diagnosis?

- Yes
- No
- Not applicable
- I was already doing this

12. As a result of the information presented did you use the STD Treatment Guidelines wall chart or pocket guide?

- Yes
- No
- Not applicable
- I was already doing this

13. Now, 3 months AFTER training, approximately what % of sexually active asymptomatic female patients under age 25 are you screening annually for chlamydia?

- 0%
- 1-25%
- 26-50%
- 51-75%
- 76-90%
- >91%
- NA

14. Now, 3 months AFTER training, approximately what % of sexually active asymptomatic female patients under age 25 are you screening annually for gonorrhea?

- 0%
- 1-25%
- 26-50%
- 51-75%
- 76-90%
- >91%
- NA

15. Now, 3 months AFTER training, approximately what % of your male patients who have sex with men are you screening annually for gonorrhea and chlamydia?

- 0%
- 1-25%
- 26-50%
- 51-75%
- 76-90%
- >91%
- NA

16. As a result of the information presented did you provide Expedited Partner Therapy (EPT) to heterosexual partners of those diagnosed with gonorrhea and/or chlamydia?

- Yes
- No
- Not applicable to my practice or job
- My practice/worksites is in the planning stages to offer EPT
- My practice/worksites already offers EPT
- EPT was not discussed

17. Now, 3 months AFTER training, approximately what % of your male patients who have sex with men are you screening at least once a year for syphilis?

- 0%
- 1-25%
- 26-50%
- 51-75%
- 76-90%
- >91%
- NA

18. Now, 3 months AFTER training, approximately what % of your patients recently diagnosed with syphilis are you testing for HIV?

- 0% 1-25% 26-50% 51-75% 76-90% >91% NA

19. What is the recommended treatment for a patient diagnosed with uncomplicated urethral, cervical, or rectal gonorrhea?

- Ceftriaxone 250 mg intramuscularly only
 Azithromycin 2 g orally in a single dose only
 Ceftriaxone 250 mg intramuscularly plus azithromycin 1 g orally in a single dose
 Cefixime 400 mg orally plus doxycycline 100 mg orally BID for 7 days

20. What is the recommended follow-up for a non-pregnant patient after diagnosis and treatment of chlamydia, gonorrhea, and/or trichomonas?

- A test of cure at 2 weeks, and repeat at 3 months
 A test of cure at 2 weeks, and repeat test at 12 months
 Repeat test in 3 months
 Repeat test om 12 months

21. What is recommended for STD screening of an HIV-negative man who reports oral sex (oral and penile exposure) and receptive anal sex with multiple male partners?

- Pharyngeal GC, rectal GC/CT, urethral GC/CT and syphilis every 3-6 months
 Pharyngeal GC, Rectal GC/CT, urethral GC/CT and syphilis every 12-24 months
 Urethral GC/CT and syphilis every 3-6 months, with pharyngeal GC and rectal GC/CT if symptoms are present
 Urethral GC/CT and syphilis every 12-24 months, with pharyngeal GC and rectal GC/CT if symptoms

THANK YOU!

OMB Control Number 0920-0995

Exp. Date 10/31/2016