

## STD Treatment Guidelines Short Long-Term Evaluation

Public reporting burden of this collection of information is estimated to average 3 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Reports Clearance Officer, 1600 Clifton Road NE, MS D-74, Atlanta, Georgia 30333; ATTN: PRA (0920-0995).

1. Did you incorporate any of the information presented into your patient care or practice setting?

- Yes
- No
- Not applicable to my practice or patients
- I was already performing the recommended practices
- Other (please specify)

2. As a result of the information presented did you download the CDC STD Treatment Guidelines app?

- Yes
- No
- Not applicable
- I was already doing this

3. As a result of the information presented did you increase the proportion of your sexually active asymptomatic female patients under age 25 screened for urogenital chlamydia and gonorrhea?

- Yes
- No
- Not applicable
- I was already doing this

4. As a result of the information presented did you increase the proportion of your male patients who have sex with men screened for syphilis, gonorrhea, and chlamydia at least annually?

- Yes
- No
- Not applicable
- I was already doing this

5. As a result of the information presented did you use dual antibiotic therapy to treat uncomplicated gonorrhea?

- Yes
- No
- Not applicable
- I was already doing this

6. As a result of the information presented did you send a consult to the STD Clinical Consultation Network? [www.stdccn.org](http://www.stdccn.org)

- Yes
- No
- Not applicable
- I was already doing this

7. As a result of the information presented did you recommend rescreening in 3 months following a gonorrhea, chlamydia or trichomonas diagnosis?

- Yes
- No
- Not applicable
- I was already doing this

8. As a result of the information presented did you use the STD Treatment Guidelines wall chart or pocket guide?

- Yes
- No
- Not applicable
- I was already doing this

9. As a result of the information presented did you share information about the STD Treatment Guidelines with others?

- Yes
- No
- Not applicable
- I was already doing this

10. As a result of the information presented did you provide Expedited Partner Therapy (EPT) to heterosexual partners of those diagnosed with gonorrhea and/or chlamydia?

- Yes
- No
- Not applicable to my practice or job
- My practice/worksites is in the planning stages to offer EPT
- My practice/worksites already offers EPT
- EPT was not discussed

11. Did any of these factors MAKE IT HARDER for you to apply the STD practices recommended in the training? (select all that apply)

- lack of time with patients
- more important patient concerns
- cost/lack of reimbursement
- policies where I work
- resistance to change by supervisor or colleagues
- lack of equipment or supplies
- no opportunity to apply practices
- nothing interfered
- Other (please specify)

12. Did any of these factors HELP you apply the STD practices recommended in the training? (select all that apply)

- reimbursement or other financial incentive
- support of supervisor and/or colleagues
- standing orders
- reminder in chart
- convenient supplies
- posted patient instructions for obtaining specimens
- nothing specific helped
- Other (please specify)

13. Do you use the CDC STD Treatment Guidelines to guide the care of your patients/clients?

- No, I am not aware of the Guidelines
- I am aware of the Guidelines but do not use them
- I use the Guidelines occasionally
- I use the Guidelines consistently
- I use another source to guide my STD care (please specify)

14. What is the recommended treatment for a patient diagnosed with uncomplicated urethral, cervical, or rectal gonorrhea?

- Ceftriaxone 250 mg intramuscularly only
- Azithromycin 2 g orally in a single dose only
- Ceftriaxone 250 mg intramuscularly plus azithromycin 1 g orally in a single dose
- Cefixime 400 mg orally plus doxycycline 100 mg orally BID for 7 days

15. What is the recommended follow-up for a non-pregnant patient after diagnosis and treatment of chlamydia, gonorrhea, and/or trichomonas?

- A test of cure at 2 weeks, and repeat at 3 months
- A test of cure at 2 weeks, and repeat test at 12 months
- Repeat test in 3 months
- Repeat test om 12 months

16. What is recommended for STD screening of an HIV-negative man who reports oral sex (oral and penile exposure) and receptive anal sex with multiple male partners?

- Pharyngeal GC, rectal GC/CT, urethral GC/CT and syphilis every 3-6 months
- Pharyngeal GC, Rectal GC/CT, urethral GC/CT and syphilis every 12-24 months
- Urethral GC/CT and syphilis every 3-6 months, with pharyngeal GC and rectal GC/CT if symptoms are present
- Urethral GC/CT and syphilis every 12-24 months, with pharyngeal GC and rectal GC/CT if symptoms

**THANK YOU!**

OMB Control Number 0920-0995

Exp. Date 10/31/2016