

CDC estimates the average public reporting burden for this collection of information as 3 minutes per survey, including the time for reviewing instructions, searching existing data/information sources, gathering and maintaining the data/information needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Information Collection Review Office, 1600 Clifton Road NE, MS D74, Atlanta, Georgia 30333; ATTN: PRA (0920XXXX).

OBGYN BI-MI Proficiency Rating Scale (PRS) - Provider Skills-Training Baseline

Instructions:

Please indicate how well you did each of the following during this simulated screening and brief intervention.

Use the following anchors for items 1-10 and please note that Item 11 has its own set of anchors below it.

1 = I did not do this.

2 = I attempted, but could improve on skill/technique for best practice.

3 = I performed this skill/technique at a level that is approaching acceptable.

4 = I did this well, with good technique.

5 = I did very well, with positive reception & engagement from the patient.

1. Ask for permission to talk about patient's substance use.	1	2	3	4	5
2. Assess quantity, frequency, & consequences of alcohol use.	1	2	3	4	5
3. Explain NIAAA low risk drinking guidelines (including "0 drinks for pregnant women" and associated health risks.	1	2	3	4	5
4. Advise the patient to quit or cut down on alcohol use.	1	2	3	4	5
5. Help her think about pros & cons of her alcohol use.	1	2	3	4	5
6. Ask how ready she is to make a change.	1	2	3	4	5
7. Help the patient make a plan or set a goal for decreasing use and/or discussing further.	1	2	3	4	5

8. Explore patient's own reasons for quitting or cutting down on alcohol use.	1	2	3	4	5
9. Work with the patient as a partner in addressing her alcohol use issues.	1	2	3	4	5
10. Support her autonomy and choice regarding substance use.	1	2	3	4	5

11. Because of this intervention, the patient's motivation to cut down or quit using alcohol, or at least to consider doing so, was increased.

1	2	3	4	
5				
Strongly	Disagree	Neutral	Agree	Strongly
Disagree				
Agree				