

This document contain screenshots for the **AAP Pre-Training Evaluation Survey**.

As noted in the request for approval of non-substantive changes, respondents to this survey see a subset of the questions depending on the training they take prior to receiving the survey. There are three trainings that respondents may take, leading to two “paths” through the survey. Screenshots for these paths are presented separately in this document, as follows:

1. Pre-Training Survey Path 1 (Training: Screening, Assessment, and Diagnosis or ND-PAE)¹
2. Pre-Training Survey Path 2 (Training: Treatment Across the Lifespan)

¹ Note that the Screening, Assessment, and Diagnosis survey screenshots are included as the example in this document, but the content of the ND-PAE survey is identical.]

AAP Pre-Training Evaluation Survey

Survey Path 1



Training: Screening, Assessment, and Diagnosis or ND-PAE¹

¹ Note that the Screening, Assessment, and Diagnosis survey screenshots are included as the example here, but the content of the ND-PAE survey is identical.

Peds DSW Pre-Training Evaluation Survey - Screening, Assessment and Diagnosis

You have unlocked your survey. When you have finished editing, please lock your survey again.

▼ Default Question Block
Block Options ▼

Form Approved
OMB No. 0920-1129
Exp. Date 08/31/2019

Thank you for your interest in fetal alcohol spectrum disorders (FASD). We would like to invite you to complete a pre-training evaluation survey. We appreciate your willingness to help us evaluate the effectiveness of the training and its impact on your practice as you address the prevention, identification, and treatment of FASD.




This survey will take approximately 3 minutes to complete. Your responses will be kept secure and no individually identifying information will be included. Risks to participating in this survey are minimal and include the risk of your information becoming known to individuals outside the AAP.

Your participation is voluntary. You may decline to answer any question and you have the right to stop the survey at any time.

Please submit questions to the project partners at PEHDIC@aap.org.

The average public reporting burden for this collection of information is estimated to average 3 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any aspect of this collection of information, including suggestions for reducing this burden to - CDC/ATSDR Reports Clearance Officer, 1600 Clifton Road NE, MS D-74, Atlanta, Georgia 30333 ATTN: PRA (0920-1129).

Page Break



Unique identifier to help us match pre and post-training responses:


First 2 letters of your mother's maiden name:

Month of your birthday:

Last 2 digits of your social security number:


■ State in which you practice. Use the drop down menu to find your State.



.....Page Break.....


■ PK5 Which of the following are the primary facial dysmorphic features associated with Fetal Alcohol Syndrome (FAS)? (Check all that apply)



- Wide inner canthal distance
- Short palpebral fissures
- Full lips
- Smooth philtrum
- Thin upper lip
- Flaring nares
- Don't know/unsure

.....Page Break.....


■ PK6 The diagnosis of "neurobehavioral disorder associated with prenatal alcohol exposure (ND-PAE)," as identified in the Diagnostic and Statistical Manual of Mental Disorders Fifth Edition (DSM-5):



- a. Requires recognition of neurocognitive impairment, self-regulation impairment, and deficits in adaptive functioning
- b. Can be diagnosed without knowledge of confirmed prenatal alcohol exposure
- c. Includes recognition of the 3 primary morphologic features of prenatal alcohol exposure
- d. Is the least common manifestation of prenatal alcohol exposure
- e. All of the above

.....Page Break.....

■ PK9 Which of the following approaches/care strategies is **not** applicable for children diagnosed with an FASD?



- a. Regularly scheduled follow-up in the medical home to anticipate/address needs across the lifespan.
- b. Refer child's case to therapist to provide all follow-up and lifelong monitoring because traditional behavioral therapies work best for children with an FASD.
- c. Evaluation by a psychologist to assess neurocognitive functioning, self-regulation, and adaptive functioning skills.
- d. Medication management for co-occurring conditions as needed to optimize care.
- e. All of the above are applicable approaches/care strategies for children with an FASD.

.....Page Break.....

P10 c, d

To what extent do you agree with the following statements? (Mark one response per row):

	Strongly Disagree	Disagree	Neither Agree nor Disagree	Agree	Strongly Agree
a. Diagnosis of one of the FASDs may confer a negative stigma to a child and/or his or her family	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Diagnosis of one of the FASDs only needs to be considered for certain populations	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

P11

In your opinion, how much alcohol is safe to drink during pregnancy?

Page Break

Q12 1, 2

Which of the following two statements below best corresponds with your personal viewpoint. Please check only ONE.

Occasional consumption of one standard alcoholic drink per day or less (i.e., 1.5 oz. hard liquor, 12 oz. of beer or 5 oz. of wine) during pregnancy is not harmful to the mother or the fetus.

Pregnant women or women who are trying to become pregnant should completely abstain from consuming alcohol.

Page Break

P14 b, e, l

To what extent do you agree with the following statements about alcohol consumption during pregnancy? (Mark one response per row)

	Strongly Disagree	Disagree	Neither Agree nor Disagree	Agree	Strongly Agree
a. Is more prevalent in women with higher income	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Is more prevalent in women with higher levels of education	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Does not vary between ethnic or racial groups	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Page Break

P15 a, b, c, d

How often do you do the following? (Mark one response per row)

	Never	Rarely	Sometimes	Usually	Always
a. Inquire routinely about prenatal exposure to alcohol	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Identify patient as someone who may have one of the FASDs	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Diagnose patient as someone who may have one of the FASDs	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. Refer patient for diagnosis and/or treatment services	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

P17 a, b, c

How confident are you in your skills to do the following? (Mark one response per row)

	Not at all Confident in my Skills	A Little Confident in my Skills	Moderately Confident in my Skills	Confident in my Skills	Completely Confident in my Skills
a. Inquire about potential prenatal alcohol exposure for pediatric patients	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Identify persons with possible FAS or other prenatal alcohol-related disorders	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Diagnose persons with possible FAS or other prenatal alcohol-related disorders	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Page Break

P18 a, b, c

How willing are you to do the following? (Mark one response per row)

	Not at all Willing	A little Willing	Moderately Willing	Confidently Willing	Completely Willing
a. Inquire about potential prenatal alcohol exposure for pediatric patients	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Identify persons with possible FAS or other prenatal alcohol-related disorders	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Diagnose persons with possible FAS or other prenatal alcohol-related disorders	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

P19 yes=1
no=2

Is stigma a barrier as you consider assessing for prenatal alcohol exposure in your clinical practice?

Yes

No

Page Break

P20 yes=1
no=2

During the past six months, did you diagnose any children with fetal alcohol syndrome (FAS) or one of the fetal alcohol spectrum disorders (FASDs)?

Yes

No

P20 schema

If yes, which diagnostic schema (if any) did you use to support your diagnosis: (Mark all that apply)

- Institute of Medicine criteria
- American Academy of Pediatrics algorithm and/or toolkit
- Seattle 4-Digit Diagnostic Code (University of Washington)
- Diagnostic and Statistical Manual of Mental Disorders (DSM-5)
- Other schema (please specify)

I did not use any particular schema

P21

During the past six months, did you refer any children for FASD assessment?

- Yes
- No

Page Break

P22

Please feel free to comment on your response to any of the questions in this survey.

Thank you for taking the time to answer these questions!

[Add Block](#)



AAP Pre-Training Evaluation Survey


Survey Path 2

Training: Treatment Across the Lifespan

Treatment Across the Lifespan: Pre-Training Evaluation Survey

This survey is currently LOCKED to prevent invalidation of collected responses! Please [unlock](#) your survey to make changes.

▼ Default Question Block
Block Options ▼



Thank you for your interest in fetal alcohol spectrum disorders (FASD). We would like to invite you to complete a pre-training evaluation survey. We appreciate your willingness to help us evaluate the effectiveness of the training and its impact on your practice as you address the prevention, identification, and treatment of FASD.

This survey will take approximately 3 minutes to complete. Your responses will be kept secure and no individually identifying information will be included. Risks to participating in this survey are minimal and include the risk of your information becoming known to individuals outside the AAP.


Your participation is voluntary. You may decline to answer any question and you have the right to stop the survey at any time.

Please submit questions to the project partners at PEHDIC@aap.org.


The public reporting burden of this collection of information is estimated to average 3 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any aspect of this collection of information, including suggestions for reducing this burden to - CDC/ATSDR Reports Clearance Officer; 1600 Clifton Road NE, MS D-74, Atlanta, Georgia 30333 ATTN: PRA (0920-1129).

Form Approved
OMB No. 0920-1129
Exp. Date 08/31/2019


-----Page Break-----




Unique identifier to help us match pre and post-training responses:



First 2 letters of your mother's maiden name:





Month of your birthday:




Last 2 digits of your social security number:

-----Page Break-----


State in which you practice. Use the drop down menu to find your State.



Page Break

PK9 Which of the following approaches/care strategies is **not** applicable for children diagnosed with an FASD? (Check all that apply)

 Regularly scheduled follow-up in the medical home to anticipate/address needs across the lifespan.

Refer child's case to therapist to provide all follow-up and lifelong monitoring because traditional behavioral therapies work best for children with an FASD.


Evaluation by a psychologist to assess neurocognitive functioning, self-regulation, and adaptive functioning skills.

Medication management for co-occurring conditions as needed to optimize care.

All of the above are applicable approaches/care strategies for children with an FASD.


Page Break

P10 c To what extent do you agree with the following statement?

	Strongly Disagree	Disagree	Neither Agree nor Disagree	Agree	Strongly Agree
 a. Diagnosis of one of the FASDs may confer a negative stigma to a child and/or his or her family	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>



Page Break



P15 e How often do you do the following?

	Never	Rarely	Sometimes	Usually	Always
 a. Manage/coordinate the treatment of patient	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>


Page Break

P17 d, e How confident are you in your skills to do the following? (Mark one response per row)



	Not at all Confident in my Skills	A Little Confident in my Skills	Moderately Confident in my Skills	Confident in my Skills	Completely Confident in my Skills
 a. Utilize resources to refer patients for diagnosis and/or treatment for FAS(D)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
 b. Manage/coordinate the treatment of persons with FASDs	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>


Page Break

 How willing are you to do the following? (Mark one response per row)


P18 d, e


	Not at all Willing	A little Willing	Moderately Willing	Completely Willing
  a. Utilize resources to refer patients for diagnosis and/or treatment for FAS(D)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Manage/coordinate the treatment of persons with FASDs	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>


Page Break

 Please feel free to comment on your response to any of the questions in this survey.

P22



 Thank you for taking the time to answer these questions!



[Add Block](#)

