

	(aff	ix label here)		
Patient ID Number	Site	Sub-site	Sequential ID	

SEARCH Low Blood Sugar Survey – Parent version

1. <u>Behavior</u>: Below is a list of things people with diabetes sometimes DO IN ORDER TO AVOID LOW BLOOD SUGAR. Read each item carefully. Circle one of the numbers that best describes YOUR ACTIVITY.

The response categories are:

1 = Never 2 = Rarely 3 = Sometimes 4 = Often 5 = Very Often

		Never	Rarely	Some- times	Often	Very Often
1.	Have your child eat large snacks at bedtime	1	2	3	4	5
2.	Avoid having your child being alone when his/her sugar is likely to be low	1	2	3	4	5
3.	Allow blood sugar to be a little high to be on the safe side	1	. 2	3	4	5
4.	Keep child's sugar higher when he/she will be alone for a while	1	2	3	4	5
5.	Have child eat something as soon as he/she feels the first sign of low blood sugar	1	2	3	4	5
6.	Reduce my child's insulin when I think his/her sugar is too low	1	2	3	4	5
7.	Keep my child's blood sugar higher when he/she plans to be away from me for a while	1	2	3	4	5
8.	Have your child carry fast-acting sugar	1	2	3	4	5
9.	Have child avoid a lot of exercise when I think his/her sugar is low	1	2	3	4	5
10.	Check my child's sugar often when he/she plans to go on an outing	1	2	3	4	5

SEARCH 3 Cohort Study – Low Blood Sugar Survey – Parent Version – 11-01-10

Page 1 of 3

(For Parents) 2. <u>Worry</u>: Below is a list of concerns parents of children with diabetes sometimes have. Read each item carefully. Circle one of the numbers that best describes HOW OFTEN YOU WORRY ABOUT EACH ITEM.

The response categories are:

1 = Never 2 = Rarely 3 = Sometimes 4 = Often 5 = Very Often

	Never	Rarely	Some- times	Often	Very Often
11. Child not recognizing/realizing he/she is having a reaction.	1	2	3	4	5
12. Child not having food, fruit, or juice with him/her	1	2	3	4	5
13. Child feeling dizzy or passing out in public	1	2	3	4	5
14. Child having a reaction while asleep	1	2	3	4	5
15. Child embarrassing self or friends/family in a social situation	1	2	3	4	5
16. Child having a reaction while alone	1	2	3	4	5
17. Child appearing "stupid" or clumsy	1	2	3	4	5
18. Child losing control	1	2	3	4	5
19. No one being around to help child during a reaction	1	2	3	4	5
20. Child making a mistake or having an accident at school	1	2	3	4	5
21. Child getting a bad evaluation at school because of something that happens when his/her sugar is low.	1	2	3	4	5
22. Child having seizures or convulsions	1	2	3	4	5
23. Child developing long term complications from frequent low blood sugar	1	2	3	4	5
24. Child feeling light-headed or faint	1	2	3	4	5
25. Child having an insulin reaction	1	2	3	4	5

This is the end of the questionnaire. Thank you for answering these questions.

FOR STUDY USE ONLY							
Date Completed	Month	Day	Year	Completed By			
Date Reviewed				Reviewer Code			
Date Entered	Month Month	Day Day	Year Year	Data Entry Code			