

Azalea Prevention Curriculum PRE-TEST

	Agree	Neutral	Disagree
I would willingly accept someone who has been treated for substance use as a close friend.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
I feel that entering treatment for substance use is a sign of personal failure.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
I think less of a person who has been in treatment for substance use.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
I would treat someone who has been treated for substance use just as I would treat anyone else.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
I believe that someone who has been treated for substance use is just as trustworthy as the average citizen.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
I believe that a person who has been treated for substance use is just as intelligent as the average person.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

For the Statements below, please mark whether you Agree, Disagree, or are Neutral.

For the Questions below, please mark whether you believe the answer is True or False.

	True	False
Substance abuse and substance misuse are interchangeable terms.	<input type="checkbox"/>	<input type="checkbox"/>
Substance use disorder is a diagnosable medical condition.	<input type="checkbox"/>	<input type="checkbox"/>
Opioids refers to both street drugs like Heroin and prescription drugs like Vicodin.	<input type="checkbox"/>	<input type="checkbox"/>
Withdrawal symptoms from opioids is limited to physical pain such as tremors and cramping.	<input type="checkbox"/>	<input type="checkbox"/>
The infant mortality rate in Northeast Florida is less than the State of Florida average.	<input type="checkbox"/>	<input type="checkbox"/>
	True	False
Since 2013, the infant mortality rate for minorities has been double or more than the infant mortality rate of whites.	<input type="checkbox"/>	<input type="checkbox"/>
A majority of infant deaths are caused by preventable factors, such as a baby facing down when sleeping, bed sharing, and unsafe sleep environments.	<input type="checkbox"/>	<input type="checkbox"/>
Neonatal Abstinence Syndrome is a result of crack/cocaine use during pregnancy.	<input type="checkbox"/>	<input type="checkbox"/>
Infants with Neonatal Abstinence Syndrome may have lifelong developmental, behavioral and cognitive issues.	<input type="checkbox"/>	<input type="checkbox"/>
Among substance users, 25% of pregnancies are unintended.	<input type="checkbox"/>	<input type="checkbox"/>

	Agree	Neutral	Disagree
I would willingly accept someone who has been treated for substance use as a close friend.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
I feel that entering treatment for substance use is a sign of personal failure.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
I think less of a person who has been in treatment for substance use.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
I would treat someone who has been treated for substance use just as I would treat anyone else.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
I believe that someone who has been treated for substance use is just as trustworthy as the average citizen.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
I believe that a person who has been treated for substance use is just as intelligent as the average person.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Azalea Prevention Curriculum POST-TEST

For the Statements below, please mark whether you Agree, Disagree, or are Neutral.

For the Questions below, please mark whether you believe the answer is True or False.

	True	False
Substance abuse and substance misuse are interchangeable terms.	<input type="checkbox"/>	<input type="checkbox"/>
Substance use disorder is a diagnosable medical condition.	<input type="checkbox"/>	<input type="checkbox"/>
Opioids refers to both street drugs like Heroin and prescription drugs like Vicodin.	<input type="checkbox"/>	<input type="checkbox"/>
Withdrawal symptoms from opioids is limited to physical pain such as tremors and cramping.	<input type="checkbox"/>	<input type="checkbox"/>
The infant mortality rate in Northeast Florida is less than the State of Florida average.	<input type="checkbox"/>	<input type="checkbox"/>
	True	False
Since 2013, the infant mortality rate for minorities has been double or more than the infant mortality rate of whites.	<input type="checkbox"/>	<input type="checkbox"/>
A majority of infant deaths are caused by preventable factors, such as a baby facing down when sleeping, bed sharing, and unsafe sleep environments.	<input type="checkbox"/>	<input type="checkbox"/>
Neonatal Abstinence Syndrome is a result of crack/cocaine use during pregnancy.	<input type="checkbox"/>	<input type="checkbox"/>
Infants with Neonatal Abstinence Syndrome may have lifelong developmental, behavioral and cognitive issues.	<input type="checkbox"/>	<input type="checkbox"/>
Among substance users, 25% of pregnancies are unintended.	<input type="checkbox"/>	<input type="checkbox"/>