

Opioid Prescribing Presentation Pre-Assessment

Start of Block: Default Question Block

Introduction: The purpose of this study is to research best practices for pain education curricula, develop an evidence-based curriculum focusing on improving opiate prescribing for women, and create core competencies for opioid prescription education for the School of Osteopathic Medicine of Rowan University. In order to continuously improve upon opioid education, this questionnaire was developed to measure provider knowledge and attitudes. **This is not a performance based test**, it is simply to gauge practitioner impressions of identifying where further education should be directed. Please answer honestly to the best of your ability.

Instructions: Please answer the following questions #1-10 to the best of your ability.

1. Which of the following are the most appropriate pharmacologic treatments for opioid dependence?

- Acamprosate.
 - Disulfiram.
 - Buprenorphine.
 - Naloxone.
-

According to the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0990-0379 . The time required to complete this information collection is estimated to average 5 minutes per response, including the time to review instructions, search existing data resources, gather the data needed, to review and complete the information collection. If you have comments concerning the accuracy of the time estimate(s) or suggestions for improving this form, please write to: U.S. Department of Health & Human Services, OS/OCIO/PRA, 200 Independence Ave., S.W., Suite 336-E, Washington D.C. 20201, Attention: PRA Reports Clearance Officer

2. Which of the following is the most appropriate approach to maintaining sobriety for opioid addiction?

- Abstinence-based approach with counseling.
 - Opioid replacement therapy with counseling (i.e. Methadone or Suboxone).
 - Behavior modification therapy.
 - Motivational interviewing based counseling.
-

3. Buprenorphine works most appropriately as a(n):

- Full opioid agonist.
 - Partial opioid agonist.
 - Opioid antagonist.
 - Combination opioid agonist/antagonist.
-

4. Naltrexone can be administered by the following method:

- Intravenous (IV).
 - Intramuscular (IM).
 - Intranasally.
 - All of the above.
-

5. According to the FDA, for which of the following conditions are extended release/long-acting (ER/LA) opioids indicated?

- Severe consistent pain at the end of life.
 - Episodic migraine pain.
 - Cancer breakthrough pain.
 - None of the above.
-

6. Which of the following are most likely to be true of female patients?

- Females have lower utilization of healthcare and physician visits.
 - Females have a higher mortality rate from opioid overdose compared to men.
 - Females experience and report pain more frequently compared to men.
 - All of the above.
-

7. Please read the following scenario and answer to the best of your ability.

Two female patients present with dull or achy pain in the lower back. Upon assessment, there appears to be no definitive cause.

The first patient, a non-Hispanic white middle-class woman, is being treated with benzodiazepines and extended release oxycodone for her chronic back pain. She has private insurance and complains that the prescribed medication is not enough. The second patient, a non-Hispanic black low-income woman with Medicaid, is being treated with metformin for her diabetes, but her primary care physician has declined to treat her with long-term opioid medication.

Both patients want opioid treatment. You continue the same dose for the first patient and start the second patient on low-dose, long acting opioids. Assuming both live in Camden County,

which patient is most at risk for engaging in heroin use?

- The first patient.
 - Both patients.
 - The second patient.
 - Neither patient.
-

8. When providing obstetric care to women, which of the following approaches to care is most appropriate?

- It is best practice in obstetric care to focus drug screenings based on factors such as poor adherence to prenatal care and history of prior adverse pregnancy outcome.
 - It is recommended for the patient to undergo medically supervised withdrawal to avoid any risks to the fetus when treating pregnant women with opioid use disorder.
 - Pregnant women with an opioid use disorder are prone to high relapse rates and it is preferable to treat patients with opioid agonist pharmacotherapy.
 - Providers should only use Screening Brief Intervention and Referral to Treatment (SBIRT) when a pregnant female patient is suspected of opiate use disorder.
-

9. When providing clinical care to breastfeeding women, the most appropriate recommendation should be:

- To discourage women from breastfeeding if taking an opioid agonist.
 - To encourage women to breastfeed if taking an opioid agonist.
 - To provide pros & cons of breastfeeding while on opioid agonists.
 - To assess for risk of relapse while on opioid agonists.
-

10. Before prescribing opioids for non-pregnant female patients, the most appropriate next step for health care providers is to:

- Ensure that there is no family medical history of opioid dependence or abuse.
- Discuss the risks and benefits of opioid use and check vital signs for patient' pain intensity.
- Take a thorough history of substance use and use formal screening tools with patients at risk.
- Discuss family planning and duration of opioid use effects on the body with reproductive-aged female patients.

Instructions: Please indicate your level of agreement with the statements listed.

Currently in my practice...

	Strongly Disagree	Disagree	Neither Agree nor Disagree	Agree	Strongly Agree
I trust that most of my patients with chronic pain are able to provide an accurate self-assessment of their pain.	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I trust that available pain scales provide reliable assessment of pain.	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Female patients are less likely to be compliant in pain management than male patients.	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I look forward to treating chronic pain patients in my practice.	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
It is very difficult to feel satisfied with the treatment outcomes of most chronic pain patients.	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
More men than women who receive opiates subsequently become addicts.	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Prescribing opioids is a high risk clinical activity for a primary care physician.



Page Break

Instructions: Please indicate your level of agreement with the statements listed.

Currently in my practice...

	Strongly Disagree	Disagree	Neither Agree nor Disagree	Agree	Strongly Agree
I understand general strategies on how to implement patient prescriber agreements.	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I able to educate my patient about proper storage and disposal of extended-release/long-acting (ER/LA) opioids.	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I counsel about the risk of overdose.	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I counsel about the particular risks associated with female patients when prescribing opioid analgesics.	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I explain to my patient the methods I use to monitor opioid misuse (urine drug tests and/or pill counts).	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

I understand how to read a urine drug screen.

I understand how to obtain information on state and federal requirements for prescribing opioids.

I know how to calculate conversion doses of commonly used opioids.

Page Break

Instructions: Please indicate your level of agreement with the statements listed in #1-10.

Currently in my practice I...

	Strongly Disagree	Disagree	Neither Agree nor Disagree	Agree	Strongly Agree
I am comfortable in responding to family calls about my patients' possible misuse of opioids.	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I feel prepared to diagnose addiction.	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I am confident that I am personally skilled in treating chronic pain.	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I am familiar with general strategies for providers prescribing opioid medications that help decrease the risk of misuse.	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I know how to refer a patient with an addiction.	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

I know how to distinguish requests for increased medication for untreated pain versus requests for increased medication because of an underlying addiction disorder.



Page Break