Form Approved OMB No. 0920-0879 Expiration Date 01/31/2021

Page 1

2020 Carbapenem Resistance Veterinary Diagnostic Laboratory Survey

The purpose of this survey is to assess current activities and technologies used to detect carbapenem resistance in Enterobacteriaceae, Pseudomonas, and Acinetobacter from dogs and cats and to determine how information regarding carbapenem-resistance is shared with laboratory clients, the veterinary and scientific community, human and animal health officials, public health, and other interested parties.

This survey is being administered through the American Association of Veterinary Laboratory Diagnosticians (AAVLD) in collaboration with the Centers for Disease Control and Prevention (CDC). It contains a series of questions regarding laboratory activities and practices related to detection, carbapenem susceptibility testing, carbapenemase mechanism testing, data sharing capabilities, and emerging technologies for detecting carbapenem resistance. Use of trade names and commercial sources is for identification only and does not imply endorsement. Participation is completely voluntary.

The estimated time to complete this survey is (estimated time from pilot testing to be input here). Your survey answers will be sent to a link at REDCap where data will be stored in a password protected electronic format. The information gathered in this survey may be published; data will be published in aggregate form without identifying any individual laboratory in order to protect the privacy of your laboratory and maintain confidentiality. A summary of the results from this survey will be provided to your laboratory via email through AAVLD.

We appreciate your participation in this survey. Please complete the survey by (date six weeks from initial email to be input here). If you have any questions or concerns regarding the survey, please contact Michelle Waltenburg [nvr6@cdc.gov].

Laboratory Name:	
State:	
Title of person completing the survey:	((director, administrator, supervisor))

CDC estimates the average public reporting burden for this collection of information as 20 minutes, including the time for reviewing instructions, searching existing data/information sources, gathering and maintaining the data/information needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing burden to CDC/ATSDR Information Collection Review Office, 1600 Clifton Road NE, MS D-74, Atlanta, Georgia 30333; ATTN: PRA (0920-0879).

09/15/2020 6:21pm

Section 1: The first set of questions ask about laboratory testing scope.
Please check the appropriate response or if needed write answers or comments in the spaces provided.
1. From which of the following sources does your laboratory receive clinical samples (mark all that apply)?
 ☐ Academic veterinary practices ☐ Private veterinary practices ☐ Other veterinary diagnostic laboratories ☐ Other (please specify below):
Specify other:
2. What proportion of samples received by your laboratory are from dogs and cats?
 ○ Less than 10% ○ Between 10% and < 25% ○ Between 25% and < 50% ○ Between 50% and < 75% ○ 75% or more

Section 2: The second set of questions ask about bacterial identification and antimicrobial
susceptibility testing (AST).
Please check the appropriate response or if needed write answers or comments in the spaces
provided.
3. Which method(s) does your laboratory currently use for Gram-negative species identification (mark all that apply)?
☐ Conventional biochemical tube tests
Commercially available test strips (e.g., API® strips)
☐ Automated bacterial ID systems (e.g., Vitek®, BD Phoenix (TM), Sensititre (TM), MicroScan) ☐ MALDI-TOF system
☐ Sequencing or other molecular based methods
Other (please specify below):
Specify other:
4. Does your laboratory conduct antimicrobial susceptibility testing (AST) on any Gram-negative isolates?
○ Yes ○ No
The following attachment contains information on carbapenem breakpoints by laboratory testing guidelines. Please consult the information as needed to answer Question 5.
consult the information as needed to answer Question 5.
[Attachment: "Laboratory testing guidelines.pdf"]
5. What laboratory testing standards does your laboratory use (select all that apply)?
☐ CLSI M100
CLSI VET08
UetCAST/EUCAST Other (places specify helew):
☐ Other (please specify below):
Specify other:



Any comments you wish to share if you use multiple standards for identification and AST:

Section 3: The following set of questions ask about Enterobacteriaceae from dogs and cats specifically. For the purpose of this survey, Enterobacteriaceae includes, but is not limited to, the following species: E. coli, Salmonella spp., Klebsiella spp., Enterobacter spp., Citrobacter spp., Providencia spp., Proteus spp., and Morganella spp.

Please check the appropriate response or if needed write answers or comments in the spaces provided.
6. Does your laboratory conduct AST on any Enterobacteriaceae isolates?
○ Yes ○ No
7. Please indicate which method your laboratory uses for primary susceptibility testing of Enterobacteriaceae from dogs and cats:
 Kirby-Bauer disk diffusion Automated testing instrument (e.g., Vitek®, BD Phoenix (TM), Sensititre (TM), MicroScan) Other broth microdilution method (please specify below): Agar dilution method Gradient diffusion method (e.g., Etest®) Other or use several methods (please specify or describe below):
Specify other broth microdilution method:
Specify other or describe several methods used:
8. Please indicate which method(s) your laboratory uses for secondary, supplemental, or confirmatory testing of Enterobacteriaceae from dogs and cats (if performed):
 ☐ Kirby-Bauer disk diffusion ☐ Automated testing instrument (e.g., Vitek®, BD Phoenix (TM), Sensititre (TM), MicroScan) ☐ Other broth microdilution method (please specify below): ☐ Agar dilution method ☐ Gradient diffusion method (e.g., Etest®) ☐ Other method (please specify below): ☐ Not performed
Specify other broth microdilution method:
Specify other:
9. Does your laboratory currently conduct AST on Enterobacteriaceae for clinical management of individual patients?

REDCap°

9a. Which isolates undergo AST for clinical decision making?
 Most Enterobacteriaceae isolates received routinely undergo AST. Only Enterobacteriaceae isolates from suspected infections undergo AST. Only Enterobacteriaceae isolates that are requested by submitter/clinician undergo AST. Other Enterobacteriaceae isolates (please specify below)
Specify other:
10. Does your laboratory currently conduct AST on Enterobacteriaceae for any monitoring/surveillance programs?
○ Yes ○ No
10a. Please select the organism(s) included in surveillance AST:
□ E. coli, commensal □ Pathogenic E. coli (e.g., STEC, EHEC, AEEC) □ Salmonella □ Klebsiella □ Enterobacter □ Citrobacter □ Providencia □ Proteus □ Morganella □ Other (please specify below):
Specify other organism(s):
11. Does your laboratory currently conduct AST on Enterobacteriaceae for any research projects?
○ Yes ○ No
11a. Please select the organism(s) included in research project AST:
□ E. coli, commensal □ Pathogenic E. coli (e.g., STEC, EHEC, AEEC) □ Salmonella □ Klebsiella □ Enterobacter □ Citrobacter □ Providencia □ Proteus □ Morganella □ Other (please specify below):
Specify other organism(s):
12. For Enterobacteriaceae, if preliminary AST suggests multidrug-resistance (e.g., resistance was detected to 2 or more classes of antibiotics), does your laboratory routinely do any additional AST or mechanism of resistance testing (e.g., ESBL, carbapenemase-production) to investigate further? Orea No At request of client

13. For Enterobacteriaceae, does your laboratory's primary AST panel for dogs and cats include one or more third-generation cephalosporins (e.g., cefpodoxime, ceftiofur, cefovecin, ceftriaxone)?
○ Yes ○ No
14. Are any third-generation cephalosporins (e.g., cefpodoxime, ceftiofur, cefovecin, ceftriaxone) included in any secondary, supplemental, or confirmatory AST for Enterobacteriaceae?
○ Yes ○ No
15. For Enterobacteriaceae, does your laboratory's primary AST panel for dogs and cats include one or more carbapenems?
○ Yes ○ No
15a. Please select the carbapenem(s) included:
☐ Imipenem ☐ Meropenem ☐ Doripenem ☐ Ertapenem ☐ Other carbapenem (please specify below):
Specify other carbapenem: ———————————————————————————————————
16. Are any carbapenems included in any secondary, supplemental, or confirmatory AST for Enterobacteriaceae?
○ Yes ○ No
16a. Please select the carbapenem(s) included:
☐ Imipenem ☐ Meropenem ☐ Doripenem ☐ Ertapenem ☐ Other carbapenem (please specify below):
Specify other carbapenem: ———————————————————————————————————
Specify other carbapenem: 17. Does your laboratory provide MIC interpretations for carbapenems in Enterobacteriaceae?
<u></u>
17. Does your laboratory provide MIC interpretations for carbapenems in Enterobacteriaceae?
17. Does your laboratory provide MIC interpretations for carbapenems in Enterobacteriaceae? Yes No Sometimes (please specify below):
17. Does your laboratory provide MIC interpretations for carbapenems in Enterobacteriaceae? O Yes O No O Sometimes (please specify below): Specify:



Section 4: The following set of questions ask about Pseudomonas aeruginosa from dogs and cats specifically. Please check the appropriate response or if needed write answers or comments in the spaces provided. 19. Does your laboratory conduct AST on any Pseudomonas aeruginosa isolates? 20. Please indicate which method your laboratory uses for primary susceptibility testing of Pseudomonas aeruginosa from dogs and cats: Kirby-Bauer disk diffusion Automated testing instrument (e.g., Vitek®, BD Phoenix (TM), Sensititre (TM), MicroScan) Other broth microdilution method (please specify below): Agar dilution method Gradient diffusion method (e.g., Etest®) Other or use several methods (please specify or describe below): Specify other broth microdilution method: Specify other or describe several methods used: 21. Please indicate which method(s) are used for secondary, supplemental, or confirmatory testing of Pseudomonas aeruginosa from dogs and cats (if performed): ☐ Kirby-Bauer disk diffusion ☐ Automated testing instrument (e.g., Vitek®, BD Phoenix (TM), Sensititre (TM), MicroScan) Other broth microdilution method (please specify below): ☐ Agar dilution method Gradient diffusion method (e.g., Etest®) Other method (please specify below): ☐ Not performed Specify other broth microdilution method: Specify other method: 22. Does your laboratory currently conduct AST on Pseudomonas aeruginosa for clinical management of individual patients? 22a. Which isolates undergo AST for clinical decision making? Most Pseudomonas aeruginosa isolates received routinely undergo AST. Only Pseudomonas aeruginosa isolates from suspected infections undergo AST. Only Pseudomonas aeruginosa isolates that are requested by submitter/clinician undergo AST. Other Pseudomonas aeruginosa isolates (please specify below):



Specify other:

23. Does your laboratory currently conduct AST on Pseudomonas aeruginosa for any monitoring/surveillance programs?
○ Yes ○ No
24. Does your laboratory currently conduct AST on Pseudomonas aeruginosa for any research projects?
○ Yes ○ No
25. For Pseudomonas aeruginosa, if preliminary AST suggested multidrug-resistance (e.g., resistance was detected to 2 or more classes of antibiotics), does your laboratory routinely do any additional AST or mechanism of resistance testing (e.g., carbapenemase-production, PCR) to investigate further?
○ Yes ○ No ○ At request of client
26. For Pseudomonas aeruginosa, does your laboratory's primary AST panel for dogs and cats include one or more third-generation cephalosporins (e.g., cefpodoxime, ceftiofur, cefovecin, ceftriaxone)?
○ Yes ○ No
27. Are any third-generation cephalosporins (e.g., cefpodoxime, ceftiofur, cefovecin, ceftriaxone) included in any secondary, supplemental, or confirmatory AST for Pseudomonas aeruginosa?
○ Yes ○ No
28. For Pseudomonas aeruginosa, does your laboratory's primary AST panel for dogs and cats include one or more carbapenems?
○ Yes ○ No
28a. Please select which carbapenem(s) are included:
☐ Imipenem ☐ Meropenem
☐ Doripenem ☐ Other carbapenem (please specify below):
Specify other:
29. Are any carbapenems included in any secondary, supplemental, or confirmatory AST for Pseudomonas aeruginosa?
○ Yes ○ No
29a. Please select which carbapenem(s) are included:
☐ Imipenem ☐ Meropenem ☐ Doripenem
☐ Other carbapenem (please specify below):
Specify other:

Page 9

30. Does your laboratory provide MIC interpretations for carbapenems in Pseudomonas aeruginosa?
○ Yes ○ No ○ Sometimes (please specify below):
Specify:
31. Does your laboratory provide zone diameter interpretations for carbapenems in Pseudomonas aeruginosa?
○ Yes ○ No ○ Sometimes (please specify below):
Specify:

Section 5: The following set of questions ask about Acinetobacter baumannii complex from dogs and cats specifically.

ease check the appropriate response or if needed write answers or comments in the space	25
ovided.	

provided.
32. Does your laboratory conduct AST on any Acinetobacter baumannii complex isolates?
○ Yes ○ No
33. Please indicate which method your laboratory uses for primary susceptibility testing of Acinetobacter baumannii complex from dogs and cats:
 ○ Kirby-Bauer disk diffusion ○ Automated testing instrument (e.g., Vitek®, BD Phoenix (TM), Sensititre (TM), MicroScan) ○ Other broth microdilution method (please specify below): ○ Agar dilution method ○ Gradient diffusion method (e.g., Etest®) ○ Other or use several methods (please specify or describe below):
Specify other broth microdilution method: ———————————————————————————————————
Specify other or describe several methods used:
34. Please indicate which method(s) are used for secondary, supplemental, or confirmatory testing of Acinetobacter baumannii complex from dogs and cats (if performed):
 ☐ Kirby-Bauer disk diffusion ☐ Automated testing instrument (e.g., Vitek®, BD Phoenix (TM), Sensititre (TM), MicroScan) ☐ Other broth microdilution method (please specify below): ☐ Agar dilution method ☐ Gradient diffusion method (e.g., Etest®) ☐ Other (please specify below): ☐ Not performed
Specify other broth microdilution method:
Specify other:
35. Does your laboratory currently conduct AST on Acinetobacter baumannii complex for clinical management of individual patients?
○ Yes ○ No
35a. Which isolates undergo AST for clinical decision making?
 Most Acinetobacter baumannii complex isolates received routinely undergo AST. Only Acinetobacter baumannii complex isolates from suspected infections undergo AST. Only Acinetobacter baumannii complex isolates that are requested by submitter/clinician undergo AST. Other Acinetobacter baumannii complex isolates (please specify below):
Specify other:



36. Does your laboratory currently conduct AST on Acinetobacter baumannii complex for any monitoring/surveillance programs?
○ Yes ○ No
37. Does your laboratory currently conduct AST on Acinetobacter baumannii complex for any research projects?
○ Yes ○ No
38. For Acinetobacter baumannii complex, if preliminary AST suggested multidrug-resistance (e.g., resistance was detected to 2 or more classes of antibiotics), does your laboratory routinely do any additional AST or mechanism of resistance testing (e.g., carbapenemase-production, PCR) to investigate further?
○ Yes ○ No ○ At request of client
39. For Acinetobacter baumannii complex, does your laboratory's primary AST panel for dogs and cats include one or more third-generation cephalosporins (e.g., cefpodoxime, ceftiofur, cefovecin, ceftriaxone)?
○ Yes ○ No
40. Are any third-generation cephalosporins (e.g., cefpodoxime, ceftiofur, cefovecin, ceftriaxone) included in any secondary, supplemental, or confirmatory AST for Acinetobacter baumannii complex?
○ Yes ○ No
41. For Acinetobacter baumannii complex, does your laboratory's primary AST panel for dogs and cats include one or more carbapenems?
○ Yes ○ No
41a. Please select the carbapenem(s) included:
☐ Imipenem ☐ Meropenem ☐ Doripenem
Other carbapenem (please specify below):
Specify other:
42. Are any carbapenems included in any secondary, supplemental, or confirmatory AST for Acinetobacter baumannii complex?
○ Yes ○ No
42a. Please select which carbapenem(s) are included:
☐ Imipenem ☐ Meropenem ☐ Doripenem
☐ Other carbapenem (please specify below):
Specify other:



43. Does your laboratory provide MIC interpretations for carbapenems in Acinetobacter baumannii complex?	
○ Yes○ No○ Sometimes (please specify below):	
Specify:	
44. Does your laboratory provide zone diameter interpretations for carbapenems in Acinetobacter baumannii complex?	
○ Yes○ No○ Sometimes (please specify below):	
Specify:	
You answered that your laboratory does not perform AST on any Enterobacteriaceae, Pseudomonas aeruginosa, or Acinetobacter baumannii complex isolates. Is this true?	
○ Yes ○ No	

Please revisit question(s) 6, 19, and/or 32 to identify which Enterobacteriaceae, Pseudomonas, or Acinetobacter baumannii complex isolates your laboratory performs AST on.

REDCap⁰

Section 6: The following questions ask about what is done when carbapenem resistance from dogs and cats is detected in any Gram-negative organism.

Please check the appropriate response or if needed write answers or comments in the spaces provided.

45. If carbapenem resistance was detected in any isolate from a dog or cat, which of the following action(s) would your laboratory initiate (mark all that apply)?
 Notify clinician at submitting facility Notify biosecurity officer or infection prevention staff at submitting facility Notify the state agriculture department Notify the state veterinarian Notify other public health official/department Perform additional laboratory testing (e.g., mCIM, CarbaNP, Modified Hodge Test [MHT], whole genome sequencing) Other action (please specify below): None of the above
Specify other action:
46. If your laboratory identified a carbapenem-resistant isolate from a dog or cat, would your laboratory perform any tests to identify carbapenemase production or carbapenemase genes?
○ Yes ○ No
46a. For which organism(s) (mark all that apply)?
 ☐ Carbapenem-resistant Enterobacteriaceae ☐ Carbapenem-resistant Pseudomonas aeruginosa ☐ Carbapenem-resistant Acinetobacter baumannii complex ☐ Other (please specify below):
Specify other:
46b. What mechanism testing would your laboratory perform (mark all that apply)?
 □ Phenotypic testing □ Commercial molecular test (e.g., Cepheid Xpert® Carba-R, BioFire® FilmArray®, VERIGENE®) □ PCR for carbapenemase genes □ Whole genome sequencing
46c. What phenotypic testing would your laboratory perform for carbapenem-resistant Enterobacteriaceae?
 Modified carbapenem-inactivation method (mCIM) or carbapenem-inactivation method (CIM) □ Rapid commercial test (e.g., RAPIDEX® CarbaNP) □ Modified Hodge Test (MHT) □ Other (please specify below):
What other phenotypic testing is performed for carbapenem-resistant Enterobacteriaceae?



46d. What phenotypic testing would your laboratory perform for carbapenem-resistant Pseudomonas aeruginosa?
 Modified carbapenem-inactivation method (mCIM) or carbapenem-inactivation method (CIM) □ Rapid commercial test (e.g., RAPIDEX® CarbaNP) □ Modified Hodge Test (MHT) □ Other (please specify below):
What other phenotypic testing is performed for carbapenem-resistant Pseudomonas aeruginosa?
46e. What phenotypic testing would your laboratory perform for carbapenem-resistant Acinetobacter baumannii complex?
 Modified carbapenem-inactivation method (mCIM) or carbapenem-inactivation method (CIM) □ Rapid commercial test (e.g., RAPIDEX® CarbaNP) □ Modified Hodge Test (MHT) □ Other (please specify below):
What other phenotypic testing is performed for carbapenem-resistant Acinetobacter baumannii complex?
46f. Please indicate PCRs performed to detect which carbapenemase gene(s):
 □ blaKPC □ blaNDM □ blaOXA-48-like □ blaVIM □ blaIMP □ Other (please specify below):
Specify other carbapenemase gene(s):



Section 7: The following set of questions ask about laboratory test results from dogs and cats. Please reference the standards your laboratory uses (e.g., CLSI M100, CLSI VET08, VetCAST/EUCAST) to define carbapenem resistance when answering the following questions.

Please check the appropriate response or if needed write answers or comments in the spaces provided.

The following attachment contains information on carbapenem breakpoints for various laboratory t	testing guidelines.
Please consult this information as needed to answer questions in Section 7.	

[Attachment: "Laboratory testing guidelines.pdf"]
47. Did your laboratory identify any carbapenem-resistant isolate(s) from a dog or cat in 2019 (Jan-Dec)?
48a. Approximately how many Enterobacteriaceae isolates from dogs and cats were tested for antibiotic susceptibility in 2019 (Jan-Dec)?
○ 0-10 ○ 11-50 ○ 51-100 ○ 101-200 ○ >201 ○ Prefer not to answer
48b. Among the Enterobacteriaceae isolates from dogs and cats that your laboratory performed antibiotic susceptibility, how many were resistant to a carbapenem?
\bigcirc Zero \bigcirc 1-5 \bigcirc 6-10 \bigcirc 11-15 \bigcirc >16 \bigcirc Prefer not to answer
48c. Among the Enterobacteriaceae isolates from dogs and cats that your laboratory performed antibiotic susceptibility, how many were resistant to a third-generation cephalosporin?
○ 0-5 ○ 6-25 ○ 26-50 ○ 51-100 ○ >100 ○ Prefer not to answer
49a. Approximately how many Pseudomonas aeruginosa isolates from dogs and cats were tested for antibiotic susceptibility in 2019 (Jan-Dec)?
○ 0-10 ○ 11-50 ○ 51-100 ○ 101-200 ○ >201 ○ Prefer not to answer
49b. Among the Pseudomonas aeruginosa isolates from dogs and cats that your laboratory performed antibiotic susceptibility, how many were resistant to a carbapenem?
49c. Among the Pseudomonas aeruginosa isolates from dogs and cats that your laboratory performed antibiotic susceptibility, how many were resistant to a third-generation cephalosporin?
\bigcirc Zero \bigcirc 1-5 \bigcirc 6-10 \bigcirc 11-15 \bigcirc >16 \bigcirc Prefer not to answer
50a. Approximately how many Acinetobacter baumannii complex isolates from dogs and cats were tested for antibiotic susceptibility in 2019 (Jan-Dec)?
\bigcirc Zero \bigcirc 1-5 \bigcirc 6-10 \bigcirc 11-15 \bigcirc >16 \bigcirc Prefer not to answer

₹EDCap°

09/15/2020 6:21pm

50b. Among the Acinetobacter baumannii complex isolates from dogs and cats that your laboratory performed antibiotic susceptibility, how many were resistant to a carbapenem?
\bigcirc Zero \bigcirc 1-5 \bigcirc 6-10 \bigcirc 11-15 \bigcirc >16 \bigcirc Prefer not to answer
50c. Among the Acinetobacter baumannii complex isolates from dogs and cats that your laboratory performed antibiotic susceptibility, how many were resistant to a third-generation cephalosporin?
\bigcirc Zero \bigcirc 1-5 \bigcirc 6-10 \bigcirc 11-15 \bigcirc >16 \bigcirc Prefer not to answer
51. Does your laboratory save a viable culture of carbapenem-resistant isolates?
○ Yes ○ No
51a. Which isolates are routinely saved (select all that apply)?
☐ Clinical sterile site ☐ Clinical non-sterile site ☐ Surveillance or research specimens ☐ Other (please specify below):
Specify other:
51b. How long are carbapenem-resistant isolates saved?
52. To your knowledge, are there any public health reporting mandates that require your laboratory to report the identification of any carbapenem-resistant organisms to public health?
○ Yes (please specify below):○ No
52a. Please describe the public health reporting mandates:
((e.g., what organisms and/or resistance mechanisms, specimen sources, animals, and/or situations are covered by the mandate?))
53. May we contact your laboratory via AAVLD with questions about your survey responses?
○ Yes ○ No
54. Any additional comments you wish to share: