

NIAAA Website Survey for Website Satisfaction Rating

OMB#: 0925-0648 Exp. Date: 05/31/2021

The screenshot shows the NIAAA website page titled "Alcohol Facts and Statistics". The page includes a navigation menu with categories like "Alcohol & Your Health", "Publications & Multimedia", "Research", "Research Training", "Grants Funding", "News & Events", and "About NIAAA". A sidebar on the left lists "In this Section" with links to "Alcohol Facts & Statistics", "What Is A Standard Drink?", and "Drinking Levels Defined". The main content area is titled "Alcohol Facts and Statistics" and contains several sections with bullet points:

- Alcohol Use in the United States:**
 - Prevalence of Drinking:** According to the 2015 National Survey on Drug Use and Health (NSDUH), 86.4 percent of people ages 16 or older reported that they drank alcohol at some point in their lifetime. 56.0 percent reported that they drank in the past year. 56.0 percent reported that they drank in the past month. 1.0 percent reported that they engaged in binge drinking in the past month. 7.0 percent reported that they engaged in heavy alcohol use in the past month.² (See "Definitions" box for definitions of binge drinking and heavy alcohol use.)
 - Prevalence of Binge Drinking and Heavy Alcohol Use:** In 2015, 26.9 percent of people reported that they engaged in binge drinking in the past month. 7.0 percent reported that they engaged in heavy alcohol use in the past month.² (See "Definitions" box for definitions of binge drinking and heavy alcohol use.)
- Alcohol Use Disorder (AUD) in the United States:**
 - Adults (ages 18+):** According to the 2015 NSDUH, 15.1 million adults ages 18 and older in this age group⁴ had AUD. This includes 9.8 million men⁵ (8.4 percent of men in this age group) and 5.3 million women⁶ (4.2 percent of women in this age group).
 - About 6.7 percent of adults who had AUD in the past year received treatment. This includes 4.1 percent of men and 5.4 percent of women with AUD in this age group.⁷
 - Youth (ages 12-17):** According to the 2015 NSDUH, an estimated 623,000 adolescents ages 12-17 in this age group⁸ had AUD. This number includes 298,000 males⁹ (2.3 percent of males in this age group) and 325,000 females⁹ (2.7 percent of females in this age group).
 - About 5.2 percent of youth who had AUD in the past year received treatment. This includes 3.1 percent of males and 5.3 percent of females with AUD in this age group.⁹
- Alcohol-Related Deaths:**
 - An estimated 88,000⁸ people (approximately 62,000 men and 26,000 women⁸) die from alcohol-related causes annually, making alcohol the third leading preventable cause of death in the United States. Tobacco is the first, and the second is poor diet and physical inactivity.⁹
 - In 2014, alcohol-impaired driving fatalities accounted for 9,967 deaths (31 percent of overall road fatalities).¹⁰
- Economic Burden:**
 - In 2010, alcohol misuse cost the United States \$249.0 billion.¹¹
 - Three-quarters of the total cost of alcohol misuse is related to binge drinking.¹¹
- Global Burden:**
 - In 2012, 3.3 million deaths, or 5.9 percent of all global deaths (7.6 percent for men and 4.1 percent for women), were attributable to alcohol consumption.¹²
 - In 2014, the World Health Organization reported that alcohol contributed to more than 200 diseases and injury-related health conditions, most notably DSM-IV alcohol dependence (see sidebar), liver cirrhosis, cancers, and injuries.¹³ In 2012, 5.1 percent of the burden of disease and injury worldwide (139 million disability-adjusted life-years) was attributable to alcohol consumption.¹²
 - Globally, alcohol misuse was the fifth leading risk factor for premature death and disability in 2010. Among people between the ages of 15 and 49, it is the first.¹⁴ In the age group 20-39 years, approximately 25 percent of the total deaths are alcohol attributable.¹⁵

A red callout box on the right side of the page contains the following text: "From here, a user can choose to answer the first survey question. If the user answers the first question, subsequent questions may appear on the same page in the same place (no more than 6 questions). A user can also click the 'x' to make the survey go away, or they can click on the hyperlinked 'Form Approved OMB#' text to visit our privacy page, which will include the full OMB burden statement."

At the bottom right of the page, there is a feedback form titled "Was this page helpful?" with "Yes" and "No" buttons. Below the buttons, it says "Form Approved OMB# 0925-0648 Exp. Date: 05/31/2021".

<https://www.niaaa.nih.gov/alcohol-health/overview-alcohol-consumption/alcohol-facts-and-statistics>

Surveys on any NIAAA websites will link to NIAAA website privacy page (<https://www.niaaa.nih.gov/privacy>), which will be updated with the following text when we start running the survey:

Survey Disclaimer

OMB # and expiration date:

Public reporting burden for this collection of information is estimated to average five minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to, a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to: NIH, Project Clearance Branch, 6705 Rockledge Drive, MSC 7974, Bethesda, MD 20892-7974, ATTN: PRA (0925-0648). Do not return the completed form to this address.