

Form Approved OMB No. 0920-0666 Exp. Date: 11/30/2019 www.cdc.gov/nhsn

Urinary Tract Infection (UTI) for LTCF

Page 1 of 4		*required for saving				
*Facility ID:		Event #:				
*Resident ID:		*Social Security #:				
Medicare number (or comparable railroad insurance number):						
Resident Name, Last: *Gender: M F Other	First:	Middle: *Date of Birth: / /				
		Race (specify):				
*Resident type: Short-stay	∠ Long-stay					
*Date of First Admission to Facility:		*Date of Current Admission to Facility: _ / _ /				
*Event Type: UTI *Date of Event:/_/ *Resident Care Location:						
*Primary Resident Service Type: (ch	neck one)					
\square Long-term general nursing	\square Long-term dementia	☐ Long-term psychiatric				
Skilled nursing/Short-term reha						
*Has resident been transferred from	an acute care facility to your fa	acility in the past 4 weeks? \square Yes \square No				
If Yes, date of last transfer from a	acute care to your facility:/_	<u></u>				
If Yes, did the resident have an ir	ndwelling urinary catheter at the	e time of transfer to your facility?				
*Indwelling Urinary Catheter status a	at time of event onset (check o	ne):				
		\square Not in place				
If indwelling urinary catheter sta		last 2 calendar days:				
Site where indwelling uring Inserted (check one):	ary catheter $\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \$	cy \square Acute care hospital \square Other \square Unknown				
` ` ` ` ` ` ` ` ` ` ` ` ` ` ` ` ` ` ` `	catheter Insertion://_	•				
If indwelling urinary catheter not in place, was another urinary device type present at the time of event onset? \square Yes \square No						
in mawering armary eatherer ne		device type present at the time of event offset: Yes \(\square\$ No				
If Yes, other device type:		om (males only)				
If Yes, other device type: Event Details	Suprapubic Cond					
If Yes, other device type: Event Details *Specify Criteria Used: (check all that	Suprapubic Cond					
If Yes, other device type: Event Details *Specify Criteria Used: (check all that	Suprapubic Conde	om (males only) Intermittent straight catheter Laboratory & Diagnostic Testing				
If Yes, other device type: Event Details *Specify Criteria Used: (check all the Signs & S	Suprapubic Conde	Laboratory & Diagnostic Testing Specimen collected from clean catch voided urine and a positive culture with no more than 2 species of				
If Yes, other device type: Event Details *Specify Criteria Used: (check all that Signs & Si	Suprapubic Condensate apply) Symptoms 8°C (>100°F), or > 37.2°C (>99) se of >1.1°C (>2°F) over basel	Laboratory & Diagnostic Testing Specimen collected from clean catch voided urine and a positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of				
If Yes, other device type: Event Details *Specify Criteria Used: (check all that Signs & Si	Suprapubic Condensate apply) Symptoms 8°C (>100°F), or > 37.2°C (>99) se of >1.1°C (>2°F) over basel onset hypotension	Description (males only) ☐ Intermittent straight catheter Laboratory & Diagnostic Testing				
If Yes, other device type: Event Details *Specify Criteria Used: (check all that Signs & Si	Suprapubic Conde	Laboratory & Diagnostic Testing Specimen collected from clean catch voided urine and a positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of ≥ 10 ⁵ CFU/ml Specimen collected from in/out straight catheter and a positive culture with any number of microorganisms, at				
If Yes, other device type: Event Details *Specify Criteria Used: (check all that Signs & Si	Suprapubic Conde	Laboratory & Diagnostic Testing Specimen collected from clean catch voided urine and a positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of ≥ 10 ⁵ CFU/ml Specimen collected from in/out straight catheter and a				
If Yes, other device type: Event Details *Specify Criteria Used: (check all that Signs & Si	Suprapubic Conde	Laboratory & Diagnostic Testing Specimen collected from clean catch voided urine and a positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of ≥ 10 ⁵ CFU/ml Specimen collected from in/out straight catheter and a positive culture with any number of microorganisms, at least one of which is a bacterium of ≥ 10 ² CFU/ml				
If Yes, other device type: Event Details *Specify Criteria Used: (check all that Signs & Si	Suprapubic Condent apply) Symptoms 8°C (>100°F), or > 37.2°C (>99) se of >1.1°C (>2°F) over basel onset hypotension decline less of the testes, epididymis, or lent drainage at catheter insertion	Laboratory & Diagnostic Testing Specimen collected from clean catch voided urine and a positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of ≥ 10 ⁵ CFU/ml Specimen collected from in/out straight catheter and a positive culture with any number of microorganisms, at least one of which is a bacterium of ≥ 10 ² CFU/ml Specimen collected from indwelling catheter and a				
If Yes, other device type: Event Details *Specify Criteria Used: (check all that Signs & Si	Suprapubic Conde	Laboratory & Diagnostic Testing Specimen collected from clean catch voided urine and a positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of ≥ 10 ⁵ CFU/ml Specimen collected from in/out straight catheter and a positive culture with any number of microorganisms, at least one of which is a bacterium of ≥ 10 ² CFU/ml				
If Yes, other device type: Event Details *Specify Criteria Used: (check all that Signs & Si	Suprapubic Condent apply) Symptoms 8°C (>100°F), or > 37.2°C (>99) se of >1.1°C (>2°F) over basel onset hypotension decline less of the testes, epididymis, or lent drainage at catheter insertion	Laboratory & Diagnostic Testing Specimen collected from clean catch voided urine and a positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of ≥ 10 ⁵ CFU/ml Specimen collected from in/out straight catheter and a positive culture with any number of microorganisms, at least one of which is a bacterium of ≥ 10 ² CFU/ml Specimen collected from indwelling catheter and a positive culture with any number of microorganisms, at least one of which is a bacterium of ≥ 10 ⁵ CFU/ml Specimen collected from indwelling catheter and a positive culture with any number of microorganisms, at least one of which is a bacterium of ≥ 10 ⁵ CFU/ml Leukocytosis (> 14,000 cells/mm³), or Left shift (> 6% or				
If Yes, other device type: Event Details *Specify Criteria Used: (check all that Signs & Si	Suprapubic Conde	Laboratory & Diagnostic Testing Specimen collected from clean catch voided urine and a positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of ≥ 10 ⁵ CFU/ml Specimen collected from in/out straight catheter and a positive culture with any number of microorganisms, at least one of which is a bacterium of ≥ 10 ² CFU/ml Specimen collected from indwelling catheter and a positive culture with any number of microorganisms, at least one of which is a bacterium of ≥ 10 ⁵ CFU/ml				
If Yes, other device type: Event Details *Specify Criteria Used: (check all that Signs & Si	Suprapubic Conde	Laboratory & Diagnostic Testing Specimen collected from clean catch voided urine and a positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of ≥ 10 ⁵ CFU/ml Specimen collected from in/out straight catheter and a positive culture with any number of microorganisms, at least one of which is a bacterium of ≥ 10 ² CFU/ml Specimen collected from indwelling catheter and a positive culture with any number of microorganisms, at least one of which is a bacterium of ≥ 10 ⁵ CFU/ml Specimen collected from indwelling catheter and a positive culture with any number of microorganisms, at least one of which is a bacterium of ≥ 10 ⁵ CFU/ml Leukocytosis (> 14,000 cells/mm³), or Left shift (> 6% or				
If Yes, other device type: Event Details *Specify Criteria Used: (check all that Signs & Si	Suprapubic Conde	Laboratory & Diagnostic Testing Specimen collected from clean catch voided urine and a positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of ≥ 10 ⁵ CFU/ml Specimen collected from in/out straight catheter and a positive culture with any number of microorganisms, at least one of which is a bacterium of ≥ 10 ² CFU/ml Specimen collected from indwelling catheter and a positive culture with any number of microorganisms, at least one of which is a bacterium of ≥ 10 ⁵ CFU/ml Specimen collected from indwelling catheter and a positive culture with any number of microorganisms, at least one of which is a bacterium of ≥ 10 ⁵ CFU/ml Leukocytosis (> 14,000 cells/mm³), or Left shift (> 6% or 1,500 bands/mm³) Positive blood culture with 1 matching organism in urine				
If Yes, other device type: Event Details *Specify Criteria Used: (check all that Signs & Si	Suprapubic Conde	Laboratory & Diagnostic Testing Specimen collected from clean catch voided urine and a positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of ≥ 10⁵ CFU/ml Specimen collected from in/out straight catheter and a positive culture with any number of microorganisms, at least one of which is a bacterium of ≥ 10² CFU/ml Specimen collected from indwelling catheter and a positive culture with any number of microorganisms, at least one of which is a bacterium of ≥ 10² CFU/ml Specimen collected from indwelling catheter and a positive culture with any number of microorganisms, at least one of which is a bacterium of ≥ 10⁵ CFU/ml Leukocytosis (> 14,000 cells/mm³), or Left shift (> 6% or 1,500 bands/mm³) Positive blood culture with 1 matching organism in urine culture SUTI) Asymptomatic Bacteremic UTI (ABUTI)				
If Yes, other device type: Event Details *Specify Criteria Used: (check all that Signs & Si	Suprapubic Conde	Laboratory & Diagnostic Testing Specimen collected from clean catch voided urine and a positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of ≥ 10 ⁵ CFU/ml Specimen collected from in/out straight catheter and a positive culture with any number of microorganisms, at least one of which is a bacterium of ≥ 10 ² CFU/ml Specimen collected from indwelling catheter and a positive culture with any number of microorganisms, at least one of which is a bacterium of ≥ 10 ⁵ CFU/ml Specimen collected from indwelling catheter and a positive culture with any number of microorganisms, at least one of which is a bacterium of ≥ 10 ⁵ CFU/ml Leukocytosis (> 14,000 cells/mm³), or Left shift (> 6% or 1,500 bands/mm³) Positive blood culture with 1 matching organism in urine culture				
If Yes, other device type: Event Details *Specify Criteria Used: (check all that Signs & Si	Suprapubic Conde	Laboratory & Diagnostic Testing Specimen collected from clean catch voided urine and a positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of ≥ 10⁵ CFU/ml Specimen collected from in/out straight catheter and a positive culture with any number of microorganisms, at least one of which is a bacterium of ≥ 10² CFU/ml Specimen collected from indwelling catheter and a positive culture with any number of microorganisms, at least one of which is a bacterium of ≥ 10² CFU/ml Specimen collected from indwelling catheter and a positive culture with any number of microorganisms, at least one of which is a bacterium of ≥ 10⁵ CFU/ml Leukocytosis (> 14,000 cells/mm³), or Left shift (> 6% or 1,500 bands/mm³) Positive blood culture with 1 matching organism in urine culture SUTI) Asymptomatic Bacteremic UTI (ABUTI)				

Assurance of Confidentiality: The voluntarily provided information obtained in this surveillance system that would permit identification of any individual or institution is collected with a guarantee that it will be held in strict confidence, will be used only for the purposes stated, and will not otherwise be disclosed or released without the consent of the individual, or the institution in accordance with Sections 304, 306 and 308(d) of the Public Health Service Act (42 USC 242b, 242k, and 242m(d)).

242m(d)). Public reporting burden of this collection of information is estimated to average 35 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC, Reports Clearance Officer, 1600 Clifton Rd., MS D-74, Atlanta, GA 30333, ATTN: PRA (0920-0666).



Urinary Tract Infection (UTI) for LTCF

Page 2 of 4

Pathogen #	Gram-positive Organisms									
	Staphylococcus coagulase-negative (specify species if available):		VANC SIRN							
	Enterococcu	Enterococcus faecium				LNZ SIRN	VANC SIRN			
	Enterococcu	Enterococcus faecalis								
	Enterococcus spp. (Only those not identified to the species level)									
	Staphylococcus aureus	CIPRO/LEVO/MOXI SIRN		CLIND SIRN	DAPTO S NS N	DOXY/MINO SIRN	ERYTH SIRN	GENT SIRN	LNZ SRN	
	ota, o a o	OX/CEFOX SIRN	X/METH	RIF SIRN	TETRA SIRN	TIG S NS N	TMZ SIRN	VANC SIRN		
Pathogen #	Gram-negative C	Organisms	•							
	Acinetobacter (specify species)	AMK SIRN	AMPSUL SIRN	AZT SIRN			CIPRO/L SIRN	CIPRO/LEVO SIRN TETRA/DOXY/MINO SIRN		
		GENT SIRN	IMI SIRN							
		TMZ SIRN	TOBRA SIRN							
	Escherichia coli	AMK SIRN	AMP SIRN	AMPSUL SIRN	/AMXCLV	AZT SIRN	CEFAZ SIRN	CEFEP S I/S-DD R N	CEFOT/CEFTRX SIRN	
		CEFTAZ SIRN	CEFUR SIRN	CEFOX/C SIRN	CEFOX/CETET SIRN		CIPRO/LEVO/MOXI SIRN		COL/PB [†] S R N	
		ERTA SIRN	GENT SIRN	IMI SIRN	MERO/DO SIRN	RI	PIPTAZ SIRN	TETRA/DOXY/I	MINO	
		TIG SIRN	TMZ SIRN	TOBRA SIRN						
	Enterobacter (specify species)	AMK SIRN	AMP SIRN	AMPSUL SIRN	/AMXCLV	AZT SIRN	CEFAZ SIRN	CEFEP S I/S-DD R N	CEFOT/CEFTRX SIRN	
		CEFTAZ SIRN	CEFUR SIRN	CEFOX/C SIRN	CEFOX/CETET CIPRO/LEV		O/MOXI	OXI COL/PB [†] S R N		
		ERTA SIRN	GENT SIRN	IMI SIRN	MERO/DO SIRN	RI	PIPTAZ SIRN	TETRA/DOXY/I	MINO	
		TIG SIRN	TMZ SIRN	TOBRA SIRN						
	Klebsiella pneumonia	AMK SIRN	AMP SIRN	AMPSUL SIRN	/AMXCLV	AZT SIRN	CEFAZ SIRN	CEFEP S I/S-DD R N	CEFOT/CEFTRX SIRN	
	, Klebsiella	CEFTAZ SIRN	CEFUR SIRN	CEFOX/C SIRN	ETET	CIPRO/LEVO SIRN	CIPRO/LEVO/MOXI COL/PB [†] SIRN SRN			
	oxytoca	ERTA SIRN	GENT SIRN	IMI SIRN	MERO/DORI		RI PIPTAZ TETRA/DOXY/MINO SIRN SIRN		MINO	
		TIG SIRN	TMZ SIRN	TOBRA SIRN						



Urinary Tract Infection (UTI) for LTCF

Page 3 of 4

Pathogen #	Gram-negative Organisms (continued)									
	Pseudomonas aeruginosa	AMK SIRN	AZT SIRN	CEFEP SIRN	CEFTAZ SIRN		CIPRO/LEVO SIRN	COL/PE SIRN	GENT SIRN	J
		IMI SIRN	MERO/DO SIRN	ORI	PIP/PIPT SIRN	ΓAZ	TOBRA SIRN			
Pathogen #	Fungal Organis	sms								
	Candida (specify species if available)	ANID SIRN	CASPO S NS N	FLUCO S S-DD R N		FLUCY SIRN	ITRA S S-DD R N	MICA S NS N	VORI S S-DI	O R N
Pathogen #	Other Organism	ns								
	Organism 1 (specify)	Drug 1 S I R N	Drug 2 S I R N	Drug 3 S I R N	Drug 4 S I R N	Drug 5 S I R N	Drug 6 S I R N	Drug 7 S I R N	Drug 8 S I R N	Drug 9 S I R N
	Organism 1 (specify)	Drug 1 S I R N	Drug 2 S I R N	Drug 3 S I R N	Drug 4 S I R N	Drug 5 S I R N	Drug 6 S I R N	Drug 7 S I R N	Drug 8 S I R N	Drug 9 S I R N
	Organism 1 (specify)	Drug 1 S I R N	Drug 2 S I R N	Drug 3 S I R N	Drug 4 S I R N	Drug 5 S I R N	Drug 6 S I R N	Drug 7 S I R N	Drug 8 S I R N	Drug 9 S I R N

Result Codes

 $S = Susceptible \ I = Intermediate \ R = Resistant \ NS = Non-susceptible \ S-DD = Susceptible-dose dependent \ N = Not tested$ § GENTHL results: S = Susceptible/Synergistic and R = Resistant/Not Synergistic

Drug Codes:

AMK = amikacin	CEFTRX = ceftriaxone	FLUCY = flucytosine	OX = oxacillin
AMP = ampicillin	CEFUR= cefuroxime	GENT = gentamicin	PB = polymyxin B
AMPSUL = ampicillin/sulbactam	CETET= cefotetan	GENTHL = gentamicin –high level test	PIP = piperacillin
AMXCLV = amoxicillin/clavulanic acid	CIPRO = ciprofloxacin	IMI = imipenem	PIPTAZ = piperacillin/tazobactam
ANID = anidulafungin	CLIND = clindamycin	ITRA = itraconazole	RIF = rifampin
AZT = aztreonam	COL = colistin	LEVO = levofloxacin	TETRA = tetracycline
CASPO = caspofungin	DAPTO = daptomycin	LNZ = linezolid	TIG = tigecycline
CEFAZ= cefazolin	DORI = doripenem	MERO = meropenem	TMZ = trimethoprim/sulfamethoxazole
CEFEP = cefepime	DOXY = doxycycline	METH = methicillin	TOBRA = tobramycin
CEFOT = cefotaxime	ERTA = ertapenem	MICA = micafungin	VANC = vancomycin
CEFOX= cefoxitin	ERYTH = erythromycin	MINO = minocycline	VORI = voriconazole
CEFTAZ = ceftazidime	FLUCO = fluconazole	MOXI = moxifloxacin	

[†] Clinical breakpoints have not been set by FDA or CLSI, Sensitive and Resistant designations should be based upon epidemiological cutoffs of Sensitive MIC ≤ 2 and Resistant MIC ≥ 4



Form Approved OMB No. 0920-0666 Exp. Date: 11/30/2019 www.cdc.gov/nhsn

Urinary Tract Infection (UTI) for LTCF

Page 4 of 4

Custom Fields			
Label		Label	
Ed.So.	1 1		, ,
	/		
			
Comments			