

National HIV Surveillance System (NHSS)

Attachment 5.

HIV Surveillance Report

HIV Surveillance **Report**

Volume 26

Diagnoses of HIV Infection in the United States and Dependent Areas, 2014

National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention
Division of HIV/AIDS Prevention



The *HIV Surveillance Report* is published annually by the Division of HIV/AIDS Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, Georgia.

Data are presented for diagnoses of HIV infection reported to CDC through July 2015.

The *HIV Surveillance Report* is not copyrighted and may be used and reproduced without permission. Citation of the source is, however, appreciated.

Suggested citation

Centers for Disease Control and Prevention. <i>HIV Surveillance Report, 2014</i> ; vol. 26. http://www.cdc.gov/hiv/library/reports/surveillance/ . Published November 2015. Accessed [date].

On the Web: <http://www.cdc.gov/hiv/library/reports/surveillance/>

Confidential information, referrals, and educational material on HIV infection

CDC-INFO

1-800-232-4636 (in English, en Español)

1-888-232-6348 (TTY)

<http://www.cdc.gov/cdc-info/requestform.html>

Acknowledgments

Publication of this report would not have been possible without the contributions of the state and territorial health departments and the HIV surveillance programs that provided surveillance data to CDC.

This report was prepared by the following CDC staff and contractors: Zanetta Gant, Kristen Hess, Anna Satcher Johnson, Xiaohong Hu, Jianmin Li, Baohua Wu, Xiuchan Guo, Hong Zhu, Yuan Liu, Marie Morgan, Michael Friend, Mi Chen, Sridevi Wilmore, John Gerstle, Shelley Ren, Jason Price, Irene Hall, Timothy Green, Kimberly Crenshaw, the HIV Incidence and Case Surveillance Branch, the Quantitative Sciences and Data Management Branch, and the Prevention Communication Branch.

Contents

Commentary	5
Technical Notes	12
Tables	
Section 1 Diagnoses of HIV Infection and Diagnoses of Infection Classified as Stage 3 (AIDS)	
1a Diagnoses of HIV infection, by year of diagnosis and selected characteristics, 2010–2014—United States	18
1b Diagnoses of HIV infection, by year of diagnosis and selected characteristics, 2010–2014—United States and 6 dependent areas	20
2a Stage 3 (AIDS), by year of diagnosis and selected characteristics, 2010–2014 and cumulative—United States	22
2b Stage 3 (AIDS), by year of diagnosis and selected characteristics, 2010–2014 and cumulative—United States and 6 dependent areas	24
3a Diagnoses of HIV infection, by race/ethnicity and selected characteristics, 2014—United States	26
3b Diagnoses of HIV infection, by race/ethnicity and selected characteristics, 2014—United States and 6 dependent areas	28
4a Stage 3 (AIDS), by race/ethnicity and selected characteristics, 2014—United States	30
4b Stage 3 (AIDS), by race/ethnicity and selected characteristics, 2014—United States and 6 dependent areas	32
5a Diagnoses of HIV infection, by transmission category, race/ethnicity, and selected characteristics, 2014—United States	34
5b Diagnoses of HIV infection, by transmission category, race/ethnicity, and selected characteristics, 2014—United States and 6 dependent areas	42
6a Stage 3 (AIDS), by transmission category, race/ethnicity, and selected characteristics, 2014—United States	50
6b Stage 3 (AIDS), by transmission category, race/ethnicity, and selected characteristics, 2014—United States and 6 dependent areas	58
7a Diagnoses of HIV infection among children aged <13 years, by race/ethnicity, 2010–2014—United States	66
7b Diagnoses of HIV infection among children aged <13 years, by race/ethnicity, 2010–2014—United States and 6 dependent areas	66
8a Stage 3 (AIDS) among children aged <13 years, by race/ethnicity, 2010–2014 and cumulative—United States	67
8b Stage 3 (AIDS) among children aged <13 years, by race/ethnicity, 2010–2014 and cumulative—United States and 6 dependent areas	67
9 Stage 3 (AIDS) among children aged <13 years, by year of diagnosis, 1992–2014—United States and 6 dependent areas	68
10 Diagnoses of HIV infection among adult and adolescent Hispanics/Latinos, by transmission category and place of birth, 2014—United States and 6 dependent areas	69
11 Stage 3 (AIDS) among adult and adolescent Hispanics/Latinos, by transmission category and place of birth, 2014—United States and 6 dependent areas	70
Section 2 Deaths and Survival after a Diagnosis of HIV Infection or Stage 3 (AIDS) Classification	
12a Deaths of persons with diagnosed HIV infection, by year of death and selected characteristics, 2010–2013—United States	71
12b Deaths of persons with diagnosed HIV infection, by year of death and selected characteristics, 2010–2013—United States and 6 dependent areas	73
13a Deaths of persons with diagnosed HIV infection ever classified as stage 3 (AIDS), by year of death and selected characteristics, 2010–2013 and cumulative—United States	75
13b Deaths of persons with diagnosed HIV infection ever classified as stage 3 (AIDS), by year of death and selected characteristics, 2010–2013 and cumulative—United States and 6 dependent areas	77

14a	Deaths of persons with diagnosed HIV infection, by race/ethnicity and selected characteristics, 2013— United States	79
14b	Deaths of persons with diagnosed HIV infection, by race/ethnicity and selected characteristics, 2013— United States and 6 dependent areas	81
15a	Deaths of persons with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and selected characteristics, 2013—United States	83
15b	Deaths of persons with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and selected characteristics, 2013—United States and 6 dependent areas	85
16a	Survival for more than 12, 24, and 36 months after a diagnosis of HIV infection during 2005–2010, by selected characteristics—United States	87
16b	Survival for more than 12, 24, and 36 months after a diagnosis of HIV infection during 2005–2010, by selected characteristics—United States and 6 dependent areas	88
17a	Survival for more than 12, 24, and 36 months after a stage 3 (AIDS) classification during 2005–2010, by selected characteristics—United States	89
17b	Survival for more than 12, 24, and 36 months after a stage 3 (AIDS) classification during 2005–2010, by selected characteristics—United States and 6 dependent areas	90
Section 3 Persons Living with Diagnosed HIV Infection or with Infection Ever Classified as Stage 3 (AIDS) (Prevalence)		
18a	Persons living with diagnosed HIV infection, by year and selected characteristics, 2010–2013—United States	91
18b	Persons living with diagnosed HIV infection, by year and selected characteristics, 2010–2013—United States and 6 dependent areas	93
19a	Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by year and selected characteristics, 2010–2013—United States	95
19b	Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by year and selected characteristics, 2010–2013—United States and 6 dependent areas	97
20a	Persons living with diagnosed HIV infection, by race/ethnicity and selected characteristics, year-end 2013— United States	99
20b	Persons living with diagnosed HIV infection, by race/ethnicity and selected characteristics, year-end 2013— United States and 6 dependent areas	101
21a	Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and selected characteristics, year-end 2013—United States	103
21b	Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and selected characteristics, year-end 2013—United States and 6 dependent areas	105
Section 4 Data for States and Metropolitan Statistical Areas		
22	Diagnoses of HIV infection, by area of residence, 2014—United States and 6 dependent areas	107
23	Stage 3 (AIDS), by area of residence, 2014 and cumulative—United States and 6 dependent areas	108
24	Adults and adolescents living with diagnosed HIV infection, by race/ethnicity and area of residence, year-end 2013—United States and 6 dependent areas	110
25	Adults and adolescents living with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and area of residence, year-end 2013—United States and 6 dependent areas	112
26	Diagnoses of HIV infection, 2014, and persons living with diagnosed HIV infection (prevalence), year-end 2013, by metropolitan statistical area of residence—United States and Puerto Rico	114
27	Stage 3 (AIDS), 2014 and cumulative, and persons living with diagnosed HIV infection ever classified as stage 3 (AIDS) (prevalence), year-end 2013, by metropolitan statistical area of residence—United States and Puerto Rico	118
Web Addresses for Reports of State and Local HIV Surveillance		123

The Centers for Disease Control and Prevention (CDC) collects, analyzes, and disseminates surveillance data on HIV infection; these data are one of the nation's primary sources of information on HIV in the United States. The annual surveillance report, published by the Division of HIV/AIDS Prevention (DHAP), summarizes information about diagnosed HIV infection in the United States and dependent areas. HIV surveillance data are used by CDC's public health partners in other federal agencies, health departments, nonprofit organizations, academic institutions, and the general public to help focus prevention efforts, plan services, allocate resources, develop policy, and monitor trends in HIV infection.

Estimated numbers and rates of diagnoses of HIV infection during 2010–2014 are included from all 50 states, the District of Columbia, and 6 U.S. dependent areas (American Samoa, Guam, the Northern Mariana Islands, Puerto Rico, the Republic of Palau, and the U.S. Virgin Islands).

During 2010–2014, the annual estimated number and rate of diagnoses of HIV infection in the United States remained stable. Estimated numbers and rates of diagnoses of HIV infection increased in some subgroups and decreased in others. Variations in trends among groups are expected and may be due to differences in testing behaviors, targeted HIV testing initiatives, or changes in the numbers of new HIV infections in some subgroups.

REPORT CHANGES

In response to requests for expanded data from our partners, this report now includes tables that provide additional stratifications by transmission category and race/ethnicity. Tables 5a/b and 6a/b display data on diagnoses of HIV infection and stage 3 (AIDS), respectively, by transmission category and selected characteristics for each racial/ethnic group. Tables 14a/b display data on deaths of persons with diagnosed HIV infection, by race/ethnicity and selected characteristics; Tables 15a/b display data on deaths of persons with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and selected characteristics.

DEFINITIONS AND DATA SPECIFICATIONS

In 2014, the HIV surveillance case definition was revised to adapt to changes in diagnostic criteria used by laboratories and clinicians. The laboratory criteria for defining a confirmed case of HIV infection were changed to accommodate multitest algorithms that do not include previously required tests (e.g., Western blot). New to the case definition is the inclusion of criteria for differentiating HIV-1 and HIV-2 infections and for recognizing early HIV infection (stage 0), during which viral loads may be high enough and CD4 T-lymphocyte counts low enough to be confused with stage 3 (AIDS). In addition, the revised definition consolidates the staging systems for adults/adolescents and children, simplifies surveillance criteria for opportunistic illnesses (OIs) indicative of stage 3, and incorporates revisions of clinical criteria (i.e., medical record documentation) for reporting diagnoses without laboratory evidence. Because retroactive implementation of some features (e.g., the new staging system) of the 2014 case definition would be impractical, the following criteria were used to classify cases in this report:

- Cases diagnosed before 2014 were classified according to the 2008 HIV case definition.
- Cases diagnosed in 2014 were classified according to the 2014 HIV case definition.

The new CD4-based criteria for stage 3 (AIDS) among children resulted in a large increase in the number of stage 3 (AIDS) classifications among pediatric cases diagnosed during 2014. Before implementation of the 2014 revised case definition, an OI diagnosis was required for a pediatric case to meet the criteria for stage 3 classification.

The term *diagnosis of HIV infection* is defined as a diagnosis of HIV infection regardless of the stage of disease (stage 0, 1, 2, 3 [AIDS], or unknown) and refers to all persons with a diagnosis of HIV infection. The term *HIV infection, stage 3 (AIDS)*, and its condensed version—*stage 3 (AIDS)*—refer specifically to persons with diagnosed HIV whose infection was classified as stage 3 (AIDS) during a given year (for diagnoses) or whose infection has ever been classified as stage 3 (AIDS) (for prevalence and deaths). For more

information on the stages of HIV infection, please review the Technical Notes.

The data on diagnoses of HIV infection reflect the date of diagnosis (diagnosed by December 31, 2014; reported to CDC as of July 31, 2015), not the date of report to CDC. Similarly, the data specific to stage 3 (AIDS) reflect the first date the criteria for stage 3 (AIDS) were met. HIV diagnoses do not necessarily represent new infections (incidence): some infections may be classified as recent; others may be classified as longstanding. Because of reporting delays, the actual number of cases diagnosed in a given year may be higher than the numbers of diagnoses of HIV infection (unadjusted) presented for recent years; however, fluctuations in the numbers of diagnoses for a calendar year typically subside after 2 to 3 years of reporting. Where indicated, counts of diagnoses and deaths have been statistically adjusted (estimated) to correct for delays in reporting, but not for incomplete reporting.

HIGHLIGHTS OF ANALYSES

The statements in this section, unless otherwise indicated, are based on estimated (statistically adjusted) data. All rates are per 100,000 population. The standard used for reporting trends in numbers and rates is based on an increase or a decrease of 5% or more during the specified time frame (e.g., when comparing 2010 and 2014). Exceptions to this standard were made where numbers were small or rates were based on small numbers.

Please read all table titles and footnotes carefully to ensure a complete understanding of the displayed data.

Diagnoses of HIV Infection and Diagnoses of Infection Classified as Stage 3 (AIDS)

Diagnoses of HIV infection

From 2010 through 2014, the annual estimated number and the rate of diagnoses of HIV infection in the United States remained stable (Table 1a). In 2014, the estimated rate was 13.8.

- **Age group:** From 2010 through 2014, the rate for persons aged 25–29 years increased. The rates for children (aged less than 13 years) and persons aged 13–14, 15–19, 35–39, 40–44, 45–49, 50–54, 55–59, and 60–64 years decreased. The rates for persons aged 20–24, 30–34, and 65 years and older remained stable. In 2014, the highest rate was for persons aged 25–29 years

(35.8), followed by the rate for persons aged 20–24 years (34.3).

- **Race/ethnicity:** From 2010 through 2014, the rates for American Indians/Alaska Natives and Asians increased. The rates for blacks/African Americans, Native Hawaiians/other Pacific Islanders, and persons of multiple races decreased. The rates for Hispanics/Latinos and whites remained stable. In 2014, the rates were 49.4 for blacks/African Americans, 18.4 for Hispanics/Latinos, 15.4 for persons of multiple races, 10.6 for Native Hawaiians/other Pacific Islanders, 9.5 for American Indians/Alaska Natives, 6.2 for Asians, and 6.1 for whites. Please use caution when interpreting data for Native Hawaiians/other Pacific Islanders: the numbers are small.
- **Sex:** From 2010 through 2014, the rate for female adults and adolescents decreased; the rate for males remained stable. In 2014, males accounted for 81% of all diagnoses of HIV infection among adults and adolescents. The rate for male adults and adolescents was 27.4, and the rate for females was 6.1.
- **Transmission category:** From 2010 through 2014, among male adults and adolescents, the annual number of diagnosed HIV infections attributed to male-to-male sexual contact increased. The number of infections attributed to injection drug use, male-to-male sexual contact *and* injection drug use, or heterosexual contact decreased. Among female adults and adolescents, the number of infections attributed to injection drug use or heterosexual contact decreased. In 2014, among male and female adults and adolescents, the diagnosed infections attributed to male-to-male sexual contact (70%, including male-to-male sexual contact *and* injection drug use) and those attributed to heterosexual contact (24%) accounted for approximately 94% of diagnosed HIV infections in the United States.
- **Region:** From 2010 through 2014, the rates of diagnoses of HIV infection in the Northeast and the South decreased; the rates in the Midwest and the West remained stable. In 2014, rates were 18.5 in the South, 14.2 in the Northeast, 11.2 in the West, and 8.2 in the Midwest.

Stage 3 (AIDS)

From 2010 through 2014, the annual estimated number and the estimated rate of infections classified as stage 3 (AIDS) in the United States decreased (Table 2a). In 2014, the estimated rate of infections classified as stage 3 (AIDS) was 6.6.

- **Age group:** From 2010 through 2014, the rates for all age groups except children (aged less than 13 years) decreased. In 2014, the highest rate was for persons aged 45–49 years (13.7), followed by the rate for persons aged 40–44 years (12.9). Because of the change in case definition, the trend among children (aged less than 13 years) was not assessed.
- **Race/ethnicity:** From 2010 through 2014, rates for all race/ethnicity groups decreased. In 2014, the rates were 25.4 for blacks/African Americans, 12.6 for persons of multiple races, 7.7 for Hispanics/Latinos, 4.0 for American Indians/Alaska Natives, 3.5 for Native Hawaiians/other Pacific Islanders, 2.7 for whites, and 2.1 for Asians. Please use caution when interpreting data for Native Hawaiians/other Pacific Islanders: the numbers are small.
- **Sex:** From 2010 through 2014, the rates for male and female adults and adolescents decreased. In 2014, males accounted for 75% of all infections classified as stage 3 (AIDS) among adults and adolescents. The 2014 rate for males was 12.0; the 2014 rate for females was 3.8.
- **Transmission category:** From 2010 through 2014, the annual number of infections classified as stage 3 (AIDS) among male adults and adolescents with HIV infection attributed to male-to-male sexual contact, injection drug use, male-to-male sexual contact *and* injection drug use, or heterosexual contact decreased. The number of infections classified as stage 3 (AIDS) among female adults and adolescents with HIV infection attributed to injection drug use or heterosexual contact decreased.
- **Region:** From 2010 through 2014, the rates of infections classified as stage 3 (AIDS) in the 4 regions of the United States decreased. In 2014, rates were 9.2 in the South, 6.8 in the Northeast, 4.7 in the West, and 3.9 in the Midwest.

Deaths

Deaths of persons with diagnosed HIV infection

From 2010 through 2013 in the United States, the annual estimated number and rate of deaths of persons with diagnosed HIV infection decreased (Table 12a). In 2013, the estimated rate of deaths of persons with diagnosed HIV infection was 5.1. Deaths of persons with diagnosed HIV infection may be due to any cause (i.e., may or may not be related to HIV infection).

- **Age group:** From 2010 through 2013, the rate of deaths for persons aged 65 years and older increased. The rates of deaths for persons aged 15–19, 20–24, 30–34, 35–39, 40–44, 45–49, and 50–54 years decreased. The rates for children (aged less than 13 years at death) and persons aged 13–14, 25–29, 55–59, and 60–64 years remained stable.
- **Race/ethnicity:** From 2010 through 2013, the rates of deaths for all race/ethnicity groups decreased: American Indians/Alaska Natives, Asians, blacks/African Americans, Hispanics/Latinos, Native Hawaiians/other Pacific Islanders, whites, and persons of multiple races. In 2013, the highest rate of deaths was for blacks/African Americans: 19.4. Please use caution when interpreting data for Native Hawaiians/other Pacific Islanders: the numbers are small.
- **Sex and transmission category:** From 2010 through 2013, the rates of deaths for male and female adults and adolescents decreased. The number of deaths among males with diagnosed HIV infection attributed to perinatal transmission increased. The number of deaths among males with infection attributed to injection drug use, male-to-male sexual contact *and* injection drug use, or heterosexual contact decreased. The number of deaths among males with diagnosed HIV infection attributed to male-to-male sexual contact remained stable. The number of deaths among females with infection attributed to injection drug use or heterosexual contact decreased. The number of deaths among female adults and adolescents with infection attributed to perinatal transmission remained stable. Please use caution when interpreting trend data for persons with infection attributed to perinatal transmission: the numbers are small.

- **Region:** From 2010 through 2013, the rates of deaths in the 4 regions of the United States decreased. In 2013, the rate was 7.7 in the Northeast, 6.4 in the South, 3.3 in the West, and 2.8 in the Midwest.

Deaths of persons with infection ever classified as stage 3 (AIDS)

From 2010 through 2013 in the United States, the annual estimated number and rate of deaths of persons with infection ever classified as stage 3 (AIDS) decreased (Table 13a). In 2013, the estimated rate of deaths of persons with stage 3 (AIDS) was 4.1.

Deaths of persons with stage 3 (AIDS) may be due to any cause.

- **Age group:** From 2010 through 2013, the rate of deaths among persons aged 65 years and older increased. Rates of deaths among persons aged 15–19, 20–24, 30–34, 35–39, 40–44, 45–49, and 50–54 years decreased. Rates among children (aged less than 13 years) and among persons aged 13–14, 25–29, 55–59, and 60–64 years remained stable.
- **Race/ethnicity:** From 2010 through 2013, the rates of death for American Indians/Alaska Natives, blacks/African Americans, Hispanics/Latinos, Native Hawaiians/other Pacific Islanders, and persons of multiple races decreased. The rates of deaths for Asians and whites remained stable. The highest death rate was for blacks/African Americans: 15.3. Please use caution when interpreting data for Native Hawaiians/other Pacific Islanders: the numbers are small.
- **Sex and transmission category:** From 2010 through 2013, the overall rates for male and female adults and adolescents decreased. The number of deaths among males with stage 3 (AIDS) whose HIV infection was attributed to perinatal transmission increased. The number of deaths among males with infection attributed to injection drug use, male-to-male sexual contact *and* injection drug use, or heterosexual contact decreased. The number of deaths among males with infection attributed to male-to-male sexual contact remained stable. Among female adults and adolescents, the number of deaths with infection attributed to injection drug use or heterosexual contact decreased. The number of

deaths among females with infection attributed to perinatal transmission remained stable. Please use caution when interpreting data for persons with infection attributed to perinatal transmission: the numbers are small.

- **Region:** From 2010 through 2013, the rates of deaths in the 4 regions of the United States decreased. In 2013, the rate was 6.1 in the Northeast, 5.1 in the South, 2.7 in the West, and 2.2 in the Midwest.

Survival

Survival after a diagnosis of HIV infection

Unadjusted data on diagnoses of HIV infection made during 2005–2010 in the United States were used to describe the proportions of persons surviving 12, 24, and 36 months after a diagnosis of HIV infection (Table 16a).

- **Year of diagnosis:** Survival increased with the year of diagnosis for diagnoses made during 2005–2010, although year-to-year differences were small.
- **Age group:** Survival decreased as age at diagnosis increased, particularly among persons aged 50 years and older at time of diagnosis.
- **Race/ethnicity:** Survival was greatest among Native Hawaiian/other Pacific Islanders, followed by Asians, Hispanics/Latinos, whites, and persons of multiple races. Survival was lowest among American Indians/Alaska Natives and blacks/African Americans.
- **Transmission category:** Survival was greatest among children regardless of transmission category; survival was slightly lower among children with diagnosed HIV infection attributed to perinatal exposure. Among adults and adolescents, survival was greatest among males with diagnosed HIV infection attributed to male-to-male sexual contact, females with infection attributed to heterosexual contact, and males with infection attributed to male-to-male sexual contact *and* injection drug use. Survival was intermediate among male adults and adolescents with infection attributed to heterosexual contact. Survival was lowest among male and female adults and adolescents with infection attributed to injection drug use.

- **Region:** Survival was greatest among persons residing in the Midwest and the West, followed by those in the Northeast and the South.

Survival after stage 3 (AIDS) classification

Unadjusted data from diagnosed infections classified as stage 3 (AIDS) during 2005–2010 in the United States were used to describe persons surviving 12, 24, and 36 months after stage 3 (AIDS) classification (Table 17a).

- **Year of diagnosis:** In general, survival increased with the year of diagnosis for diagnoses made during 2005–2010, although year-to-year differences were small.
- **Age group:** In general, survival decreased as age at diagnosis increased, particularly among persons aged 35 years and older. Survival was greatest for persons aged 13–14 years and persons aged 15–19 years at the time of stage 3 (AIDS) classification.
- **Race/ethnicity:** Survival was greatest among Native Hawaiians/other Pacific Islanders, followed by Asians, Hispanics/Latinos, persons of multiple races, whites, blacks/African Americans, and American Indians/Alaska Natives.
- **Transmission category:** Survival was greatest among children with infection attributed to perinatal transmission, followed by male adults and adolescents with infection attributed to male-to-male sexual contact. Survival was intermediate among male adults and adolescents with infection attributed to male-to-male sexual contact *and* injection drug use and among male and female adults and adolescents with infection attributed to heterosexual contact. Survival was lowest among male and female adults and adolescents with infection attributed to injection drug use.
- **Region:** Survival was greatest among persons residing in the West, followed by those in the Northeast and the Midwest. Survival was lowest in the South.

Prevalence: Persons Living with Diagnosed HIV Infection and Persons Living with Infection Ever Classified as Stage 3 (AIDS)

Prevalence of diagnosed HIV infection

From 2010 through 2013, the estimated number of persons in the United States living with diagnosed HIV infection increased (Table 18a). At the end of 2013, an estimated 933,941 persons in the United States were living with diagnosed HIV infection. The estimated prevalence rate of diagnosed HIV infection was 295.1.

- **Age group:** At the end of 2013, persons aged 45–49 and persons aged 50–54 years made up the largest percentage of persons living with diagnosed HIV (17% each). The highest rate (754.3) was among persons aged 45–49 years, followed by those aged 50–54 (717.2) and those aged 40–44 (565.7). From 2010 through 2013, the largest percentage increase in rates (41%) was among persons aged 65 years and older (from 84.0 in 2010 to 118.5 in 2013).
- **Race/ethnicity:** At the end of 2013, the highest rate (1,018.1) and the largest percentage (43%) were those among blacks/African Americans. Among the remaining race/ethnicity groups, the rates were 523.5 for persons of multiple races, 350.8 for Hispanics/Latinos, 162.8 for Native Hawaiians/other Pacific Islanders, 150.3 for whites, 123.2 for American Indians/Alaska Natives, and 70.9 for Asians.
- **Sex and transmission category:** At the end of 2013, the estimated rates were 547.4 for male adults and adolescents, 167.7 for female adults and adolescents, and 4.6 for children (aged less than 13 years at year-end 2013). Among a total of 705,165 males (aged 13 years and older at year-end 2013), 70% of infections were attributed to male-to-male sexual contact, 11% to injection drug use, 10% to heterosexual contact, 7% to male-to-male sexual contact *and* injection drug use, and 1% to perinatal transmission. Among a total of 226,361 females (aged 13 years and older at year-end 2013), 74% of infections were attributed to heterosexual contact, 23% to injection drug use, and 2% to perinatal transmission. Among a total of 2,415 children aged less than 13 years at year-end 2013, 83% of infections were attributed to perinatal transmission.

- **Region:** At year-end 2013, the prevalence rates were 420.5 in the Northeast, 343.6 in the South, 241.2 in the West, and 165.3 in the Midwest.

- **Region:** At year-end 2013, the prevalence rates were 247.9 in the Northeast, 182.3 in the South, 140.0 in the West, and 84.8 in the Midwest.

Prevalence of stage 3 (AIDS)

From 2010 through 2013, the estimated number of persons living with infection ever classified as stage 3 (AIDS) increased (Table 19a). At the end of 2013, an estimated 516,401 persons in the United States were living with infection ever classified as stage 3 (AIDS). The estimated prevalence rate of infections classified as stage 3 (AIDS) was 163.2.

- **Age group:** At the end of 2013, the highest rate (456.8) and the largest percentage (20%) were those among persons aged 50–54 years, followed by persons aged 45–49 (454.6 and 19%, respectively). From 2010 through 2013, the largest percentage increase in rates (41%) was among persons aged 65 years and older (from 56.7 in 2010 to 79.7 in 2013).
- **Race/ethnicity:** At the end of 2013, the highest rate (554.8) and the largest percentage (42%) were those among blacks/African Americans. Among the remaining race/ethnicity groups, the rates were 318.5 for persons of multiple races, 205.5 for Hispanics/Latinos, 88.1 for Native Hawaiians/other Pacific Islanders, 80.9 for whites, 64.7 for American Indians/Alaska Natives, and 36.9 for Asians.
- **Sex and transmission category:** At the end of 2013, the estimated rates were 306.0 for male adults and adolescents, 90.4 for female adults and adolescents, and 0.5 for children (aged less than 13 years). Among a total of 394,124 males (aged 13 years and older at year-end 2013), 65% of infections were attributed to male-to-male sexual contact, 14% to injection drug use, 12% to heterosexual contact, 8% to male-to-male sexual contact *and* injection drug use, and 1% to perinatal transmission. Among a total of 122,005 females (aged 13 years and older at year-end 2013), 71% of infections were attributed to heterosexual contact, 26% to injection drug use, and 2% to perinatal transmission. Among a total of 272 children aged less than 13 years at year-end 2013, 94% of infections were attributed to perinatal transmission.

ADDITIONAL RESOURCES

The following were prepared by using HIV surveillance data:

- *MMWR* articles (selected): <http://www.cdc.gov/hiv/library/reports/mmwr.html>
- Other surveillance reports: <http://www.cdc.gov/hiv/library/reports/surveillance/>
- Public-use slides updated annually: <http://www.cdc.gov/hiv/library/slidesets/>
- NCHHSTP Atlas [interactive tool for accessing HIV/AIDS, STD, TB, and hepatitis data]: <http://www.cdc.gov/nchhstp/atlas/>

SUGGESTED READINGS

- CDC. Establishing a holistic framework to reduce inequities in HIV, viral hepatitis, STDs, and tuberculosis in the United States. <http://stacks.cdc.gov/view/cdc/11585>. Published October 2010. Accessed October 15, 2015.
- CDC. Estimated HIV incidence in the United States, 2007–2010. *HIV Surveillance Supplemental Report* 2012;17(No. 4). <http://go.usa.gov/p8P4>. Published December 2012. Accessed October 15, 2015.
- CDC. HIV surveillance—United States, 1981–2008. *MMWR* 2011;60(21):689–693.
- CDC. Monitoring selected national HIV prevention and care objectives by using HIV surveillance data—United States and 6 dependent areas—2013. *HIV Surveillance Supplemental Report* 2015;20(No. 2). <http://www.cdc.gov/hiv/library/reports/surveillance/>. Published July 2015. Accessed October 15, 2015.
- CDC. Revised surveillance case definition for HIV infection—United States, 2014. *MMWR* 2014;63(RR-03):1–10.
- CDC. Revised surveillance case definitions for HIV infection among adults, adolescents, and children aged <18 months and for HIV infection and AIDS among children aged 18 months to <13 years—United States, 2008. *MMWR* 2008;57(RR-10):1–12.

- CDC. Terms, definitions, and calculations used in CDC HIV surveillance publications. <http://www.cdc.gov/hiv/statistics/recommendations/terms.html>. Published September 2014. Accessed October 15, 2015.
- CDC. Vital Signs: HIV diagnosis, care, and treatment among persons living with HIV—United States, 2011. *MMWR* 2014;63(47):1113–1117.
- CDC. Vital Signs: HIV infection, testing, and risk behaviors among youths—United States. *MMWR* 2012;61(47):971–976.
- CDC. Vital Signs: HIV prevention through care and treatment—United States. *MMWR* 2011;60(47):1618–1623.
- CDC. Vital Signs: HIV testing and diagnosis among adults—United States. *MMWR* 2010;59(47):1550–1555.
- Cohen SM, Gray KM, Ocfemia MC, Johnson AS, Hall HI. The status of the National HIV Surveillance System, United States, 2013. *Public Health Rep* 2014;129(4):335–341.
- Hall HI, Song R, Rhodes P, et al. Estimation of HIV incidence in the United States. *JAMA* 2008;300(5):520–529.
- Johnson AS, Hall HI, Hu X, Lansky A, Holtgrave DR, Mermin J. Trends in diagnoses of HIV infection in the United States, 2002–2011 [Research Letter]. *JAMA* 2014;312(4):432–434. doi:10.1001/jama.2014.8534.
- Prejean J, Song R, Hernandez A, et al. Estimated HIV incidence in the United States, 2006–2009. *PLoS One* 2011;6(8):e17502. doi:10.1371/journal.pone.0017502.

SURVEILLANCE OF HIV INFECTION

This report includes case report data from 50 states, the District of Columbia, and 6 U.S. dependent areas (American Samoa, Guam, the Northern Mariana Islands, Puerto Rico, the Republic of Palau, and the U.S. Virgin Islands) in which laws or regulations require confidential reporting to the jurisdiction (not to CDC), by name, for adults, adolescents, and children with confirmed diagnoses of HIV infection. After the removal of personally identifiable information, data from these reports were submitted to CDC. Although AIDS cases have been reported to CDC since 1981, the implementation of HIV infection reporting has differed from state to state. All states, the District of Columbia, and 6 U.S. dependent areas had fully implemented name-based HIV infection reporting by April 2008.

Please use caution when interpreting data on diagnoses of HIV infection. HIV surveillance reports may not be representative of all persons with HIV because not all infected persons have been (1) tested or (2) tested at a time when the infection could be detected and diagnosed. Also, some states offer anonymous HIV testing; the results of anonymous tests are not reported to the confidential name-based HIV registries of state and local health departments. Therefore, reports of confidential test results may not represent all persons who tested positive for HIV infection. In addition, testing patterns are influenced by many factors, including the extent to which testing is routinely offered to specific groups and the availability of, and access to, medical care and testing services. The data presented in this report provide a minimum estimate of persons for whom HIV infection has been diagnosed and reported to the surveillance system. Finally, because surveillance practices among jurisdictions differ, the reporting and the updating of a person's clinical and vital status differ among states. The completeness of reporting of HIV infection is estimated at more than 80% [1].

STAGES OF HIV INFECTION

Both the 2008 and 2014 HIV case definitions were used to classify HIV infection among adults and adolescents and among children [2, 3].

The 2008 case definition was used to classify cases diagnosed from the beginning of the epidemic through 2013. For adults and adolescents, this definition incorporates an HIV infection staging system that includes AIDS (HIV infection, stage 3). The 2008 stages of HIV infection are defined as follows:

- **HIV infection, stage 1:** No AIDS-defining condition and either CD4 count of ≥ 500 cells/ μ L or CD4 percentage of total lymphocytes of ≥ 29 .
- **HIV infection, stage 2:** No AIDS-defining condition and either CD4 count of 200–499 cells/ μ L or CD4 percentage of total lymphocytes of 14–28.
- **HIV infection, stage 3 (AIDS):** Documentation of an AIDS-defining condition or either a CD4 count of < 200 cells/ μ L or CD4 percentage of total lymphocytes of < 14 . Documentation of an AIDS-defining condition supersedes a CD4 count or percentage that would not, by itself, be the basis for a stage 3 (AIDS) classification.
- **HIV infection, stage unknown:** No reported information on AIDS-defining conditions and no information available on CD4 count or percentage.

The 2014 case definition was used to classify cases diagnosed in 2014. It is similar to the 2008 case definition except for the following: (1) inclusion of criteria for stage 0, (2) the inclusion of CD4 testing criteria for stage 3 in children, and (3) changes in the cutoffs for CD4 percentage of total lymphocytes used for classification of stages 1 and 2 in persons aged 6 years and older. The stages of HIV infection in the 2014 case definition are based on age-specific CD4+ T-lymphocyte counts or percentages of total lymphocytes and are defined as follows:

- **HIV infection, stage 0:** First positive HIV test result within 6 months after a negative HIV test result. The stage remains stage 0 until 6 months after the first positive test result. After 6 months, the stage may be reclassified as 1, 2, 3, or unknown if based on a CD4+ T-lymphocyte

result or the diagnosis of an OI. The diagnosis of an AIDS-defining condition or a low CD4 test result before the 6 months have elapsed does not change the stage from stage 0 to stage 3.

- **HIV infection, stages 1, 2, and 3:** Documentation of an AIDS-defining condition (excluding stage 0 as described above) is stage 3. Otherwise, the stage is determined by the lowest CD4 test result [3, p 7].
- **HIV infection, stage unknown:** No reported information on AIDS-defining conditions and no information available on CD4 count or percentage.

The new CD4-based criteria for stage 3 (AIDS) among children resulted in an increase in the number of annual stage 3 (AIDS) classifications among pediatric cases diagnosed during 2014. Before implementation of the 2014 revised case definition, an OI diagnosis was required for a pediatric case to meet the criteria for stage 3 classification.

In this report, the term *diagnosis of HIV infection* is defined as a diagnosis of HIV infection regardless of the stage of disease (stage 0, 1, 2, 3 [AIDS], or unknown) and refers to all persons with a diagnosis of HIV infection. The term HIV infection, *stage 3 (AIDS)*, and its condensed version—*stage 3 (AIDS)*—refer specifically to persons with diagnosed HIV whose infection was classified as stage 3 (AIDS) during a given year (for diagnoses) or whose infection has ever been classified as stage 3 (AIDS) (for prevalence and deaths).

TABULATION AND PRESENTATION OF DATA

The data in this report include information received by CDC through July 31, 2015. The data are organized into 4 sections.

- Section 1 (Tables 1a/b–11): numbers (statistically adjusted [estimated] and unadjusted) and rates (adjusted) of diagnoses of HIV infection (any stage) and diagnoses of infection classified as stage 3 (AIDS)
- Section 2 (Tables 12a/b–17a/b): numbers (statistically adjusted and unadjusted) and rates (adjusted) of deaths and proportions (unadjusted) of survival among persons with diagnosed HIV infection (any stage) and persons with infection ever classified as stage 3 (AIDS)

- Section 3 (Tables 18a/b–21a/b): numbers (statistically adjusted and unadjusted) and rates (adjusted) of prevalence (i.e., data on persons living with diagnosed HIV infection [any stage] and data on persons living with infection ever classified as stage 3 [AIDS])
- Section 4 (Tables 22–27): numbers (statistically adjusted and unadjusted) and rates (adjusted) of diagnoses and prevalence of HIV infection (any stage) and diagnoses and prevalence of infection classified as stage 3 (AIDS), presented by state and metropolitan statistical area (MSA)

Many tables in the 2014 report present data in 2 formats. Tables in the first format—labeled “a”—exclude data from the dependent areas (American Samoa, Guam, the Northern Mariana Islands, Puerto Rico, the Republic of Palau, and the U.S. Virgin Islands). Tables in the second format—labeled “b”—include data from the dependent areas.

Please use caution when interpreting reported numbers less than 12, as well as estimated numbers and accompanying rates and trends based on these numbers: the underlying relative standard errors are greater than 30%, so the numbers are considered unreliable.

Diagnoses of HIV Infection and of Infection Classified as Stage 3 (AIDS)

This report is based on diagnoses of HIV infection that the 50 states, the District of Columbia, and 6 U.S. dependent areas reported to CDC as of July 31, 2015. Data from all areas are included in tables displaying numbers (unadjusted), estimated numbers, and estimated rates of diagnoses of HIV infection and of diagnoses of infection classified as stage 3 (AIDS), by selected characteristics, area of residence, and MSA (Tables 1a/b–11, 16a/b, 17a/b, 22, 23, 26, and 27).

Deaths

Persons reported to the National HIV Surveillance System are assumed alive unless their deaths have been reported to CDC. Death data include deaths of persons with diagnosed HIV infection (Tables 12a/b and 14a/b) or with infection ever classified as stage 3 (AIDS) (Tables 13a/b and 15a/b) regardless of the cause of death. Because of delays in the reporting of deaths, 4 years (2010–2013) of death data are displayed. The exclusion of data from the most recent

year allowed at least 18 months for deaths to be reported to CDC. The estimated numbers and rates of deaths resulted from statistical adjustment for delays in reporting. Please use caution when interpreting trend data on the estimated numbers of deaths: the estimates for the most recent year are most subject to uncertainty.

Survival Analyses

The Kaplan-Meier method was used to estimate the probability of survival for 12, 24, and 36 months after a diagnosis of HIV infection (Tables 16a/b) or after a diagnosis of infection classified as stage 3 (AIDS) (Tables 17a/b). To allow at least 3 years from the time of diagnosis to deaths occurring through December 31, 2013 (reported to CDC by July 31, 2015), table data were limited to persons whose diagnosis was made during 2005–2010. Data for each HIV reporting area were included in the survival tables beginning with the first full calendar year after implementation of code-based or name-based HIV infection reporting.

Prevalence: Persons Living with Diagnosed HIV Infection and Persons Living with Infection Ever Classified as Stage 3 (AIDS)

Because of delays in the reporting of deaths, 4 years of prevalence data are displayed. The data reflect persons living with diagnosed HIV infection, regardless of stage of disease, (Tables 18a/b, 20a/b, 24, and 26) and persons living with infection ever classified as stage 3 (AIDS) (Tables 19a/b, 21a/b, 25, and 27) at the end of a given year during 2010–2013. The exclusion of data from the most recent year allowed at least 18 months for deaths to be reported and for these deaths to be factored into calculations of prevalence.

Age

The designation “adults and adolescents” refers to persons aged 13 years and older; the designation “children” refers to persons aged less than 13 years. For presentations of data on persons living with diagnosed HIV infection (Tables 18a/b, 20a/b, and 24) and presentations of data on persons living with infection classified as stage 3 (AIDS) (Tables 19a/b, 21a/b, and 25), the age-group assignment (e.g., 20–24 years) or the age designation (e.g., “adults and adolescents”) is based on the person’s age as of December 31 of the

specified year. For presentations of data on deaths of persons with diagnosed HIV infection (Tables 12a/b and 14a/b) and deaths of persons with infection classified as stage 3 (AIDS) (Tables 13a/b and 15a/b), age-group assignment is determined by the person’s age at the time of death. For all other tables, the age designation or the specific age-group assignment is based on the person’s age at the time of diagnosis (HIV infection or infection classified as stage 3 [AIDS]).

Age-group designations for transmission categories in tables displaying data on prevalence and on deaths are based on a person’s age at the end of the specified year or age at death, respectively. That is, the transmission categories among “adults and adolescents” in these tables include persons who were children at the time of diagnosis but who were aged 13 years and older during the specified year or at death.

Sex

Sex designations in this report are based on a person’s sex at birth. *Transgender* is an umbrella term that is used to identify persons whose sex assigned at birth does not match current gender identity or expression. Data for transgender persons are not presented in this report because information on gender identity (a person’s internal understanding of his or her gender or the gender with which a person identifies) is not consistently collected or documented in the data sources used by HIV reporting jurisdictions. HIV surveillance personnel collect this information, when available, from sources such as case report forms submitted by health care or HIV testing providers and medical records, or by matching with other databases (e.g., Ryan White program data) in the health department. In May 2012, CDC issued guidance to state and local programs on methods for collecting data on transgender persons and working with transgender-specific data. However, characterization of HIV infection among transgender persons may require supplemental data from special studies.

Race and Ethnicity

In the *Federal Register* [4] for October 30, 1997, the Office of Management and Budget (OMB) announced the Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity. Implementation by January 1, 2003, was mandated. At a minimum, data on the following race categories should be collected:

- American Indian or Alaska Native
- Asian
- black or African American
- Native Hawaiian or other Pacific Islander
- white

Additionally, systems must be able to retain information when multiple race categories are reported. In addition to data on race, data on 2 categories of ethnicity should be collected:

- Hispanic or Latino
- not Hispanic or Latino

The Asian or Pacific Islander category displayed in annual surveillance reports published prior to the 2007 surveillance report was split into 2 categories:

(1) Asian and (2) Native Hawaiian or other Pacific Islander. The Asian category (in tables where foot-noted) includes the cases in Asians/Pacific Islanders (referred to as legacy cases) that were reported before the implementation of the new race categories in 2003 (e.g., cases of HIV infection that were diagnosed and reported to CDC before 2003 but that progressed to AIDS after 2003) and a small percentage of cases that were reported after 2003 but that were reported according to the old race category (Asian/Pacific Islander). In tables of diagnoses of HIV infection during 2010–2014, the Asian category does not include Asian/Pacific Islander cases because these cases were diagnosed after 2003 and were reported to CDC in accordance with OMB’s Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity [4].

This report also presents data for persons for whom multiple race categories are reported. In this report, persons categorized by race were not Hispanic or Latino. The number of persons reported in each race category may, however, include persons whose ethnicity was not reported.

Geographic Designations

Geographic designations reflect the area of residence at the time of diagnosis of HIV infection or at the time of stage 3 (AIDS) classification. The 4 regions of residence used in this report are defined by the U.S. Census Bureau as follows:

Northeast: Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, and Vermont

Midwest: Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin

South: Alabama, Arkansas, Delaware, District of Columbia, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, and West Virginia

West: Alaska, Arizona, California, Colorado, Hawaii, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington, and Wyoming

U.S. dependent areas: American Samoa, Guam, the Northern Mariana Islands, Puerto Rico, the Republic of Palau, and the U.S. Virgin Islands

Metropolitan Statistical Areas

In the *Federal Register* for June 28, 2010, the OMB published revised standards for defining MSAs in federal statistical activities [5]. These standards, which provided for the identification of MSAs in the United States and Puerto Rico, replaced the 2000 standards. The adoption of the new standards was effective as of June 28, 2010. On February 28, 2013, the OMB announced new MSA delineations based on the new standards and Census 2010 data [6]. Tables 26 (data on diagnosed HIV infection) and 27 (data on stage 3 [AIDS]) present numbers and rates of diagnoses and of prevalence, by MSA, for areas with populations of 500,000 or more. The MSAs listed in these tables were defined according to the OMB’s most recent update (February 2013) of statistical areas [6].

Transmission Categories

Transmission category is the term for the classification of cases that summarizes a person’s possible HIV risk factors; the summary classification results from selecting, from the presumed hierarchical order of probability, the 1 (single) risk factor most likely to have been responsible for transmission. For surveillance purposes, a diagnosis of HIV infection is counted only once in the hierarchy of transmission categories. Persons with more than 1 reported risk factor for HIV infection are classified in the transmission category listed first in the hierarchy. The exception is men who had sexual contact with other men

and injected drugs; this group makes up a separate transmission category.

Persons whose transmission category is classified as male-to-male sexual contact include men who have ever had sexual contact with other men (i.e., homosexual contact) and men who have ever had sexual contact with both men and women (i.e., bisexual contact). Persons whose transmission category is classified as heterosexual contact are persons who have ever had heterosexual contact with a person known to have, or to be at high risk for, HIV infection (e.g., an injection drug user). The heterosexual contact category excludes men who have ever had sexual contact with both men and women.

Tables of prevalence data and tables of death data include a separate transmission category of adults and adolescents with HIV infection attributed to perinatal transmission. The category includes persons infected through perinatal transmission but aged 13 years and older during the specified year or at death.

Cases of HIV infection reported without a risk factor listed in the hierarchy of transmission categories are classified as “no identified risk (NIR)” [7]. Cases classified as NIR include cases that are being followed up by local health department staff; cases in persons whose risk-factor information is missing because they died, declined to be interviewed, or were lost to follow-up; and cases in persons who were interviewed or for whom other follow-up information was available but for whom no risk factor was identified.

Because a substantial proportion of cases of HIV infection are reported to CDC without an identified risk factor, multiple imputation is used to assign a transmission category [8]. Multiple imputation is a statistical approach in which each missing transmission category is replaced with a set of plausible values that represent the uncertainty about the true, but missing, value [9]. The plausible values are analyzed by using standard procedures, and the results from these analyses are then combined to produce the final results. In tables displaying transmission categories, multiple imputation was used for adults and adolescents, but not for children (because the number of cases in children is small, missing transmission categories were not imputed).

Reporting Delays

Reporting delays (time between diagnosis or death and the reporting of diagnosis or death to CDC) may differ among demographic and geographic categories; for some, delays in reporting have been as long as several years. The statistical adjustment of the data on diagnoses and deaths is based on estimates of reporting-delay distributions, which are calculated by using a modified semiparametric life-table statistical procedure. This procedure takes into account differences in reporting delays due to sex, race/ethnicity, HIV transmission categories, geographic area (i.e., reporting city, state, or dependent area; region of residence), the size of the MSA, and the type of facility where the diagnosis was made or death occurred [10]. The statistical adjustment of data on diagnoses takes into consideration the information reported from the state or dependent area where the diagnosis was made; the statistical adjustment of data on deaths takes into consideration the information from the state where the death occurred. No adjustments are made for incomplete reporting.

Rates

Rates per 100,000 population were calculated for (1) the numbers of diagnoses of HIV infection and the numbers of infections classified as stage 3 (AIDS), (2) the numbers of deaths of persons with diagnosed HIV infection and deaths of persons with infection ever classified as stage 3 (AIDS), and (3) the numbers of persons living with diagnosed HIV infection and persons living with infection ever classified as stage 3 (AIDS).

The population denominators used to compute the rates for the 50 states, the District of Columbia, and Puerto Rico were based on the Vintage 2014 postcensal estimates file (for years 2010–2014) from the U.S. Census Bureau [11]. The population denominators for American Samoa, Guam, the Northern Mariana Islands, the Republic of Palau, and the U.S. Virgin Islands were based on estimates and projections from the U.S. Census Bureau’s International Data Base [12]. Each rate was calculated by dividing the estimated total number of diagnoses (or deaths or prevalence) for the calendar year by the population for that calendar year and then multiplying the result by 100,000. The denominators used for calculating the rates specific to age, sex, and race/ethnicity were

computed by applying the appropriate vintage estimates for age, sex, and race/ethnicity for the 50 states and the District of Columbia [11]. The same method was used to calculate the denominators for Puerto Rico, with the exception of race/ethnicity estimates; these data are not available for Puerto Rico (see next paragraph). For the other 5 U.S. dependent areas, estimates from the U.S. Census Bureau's International Data Base were used for age- and sex-specific population denominators [12].

CDC currently does not provide subpopulation rates by race/ethnicity for the 6 U.S. dependent areas because the U.S. Census Bureau does not collect information from all dependent areas. Rates for transmission categories are not provided in this report because of the absence of denominator data from the U.S. Census Bureau, the source of data used for calculating all rates in this report.

REFERENCES

1. Hall HI, Song R, Gerstle JE III, Lee LM; on behalf of HIV/AIDS Reporting System Evaluation Group. Assessing the completeness of reporting of human immunodeficiency virus diagnoses in 2002–2003: capture-recapture methods. *Am J Epidemiol* 2006;164(4):391–397.
2. CDC. Revised surveillance case definitions for HIV infection among adults, adolescents, and children aged <18 months and for HIV infection and AIDS among children aged 18 months to <13 years—United States, 2008. *MMWR* 2008;57(RR-10):1–12.
3. CDC. Revised surveillance case definition for HIV infection—United States, 2014. *MMWR* 2014;63(RR-03):1–10.
4. Office of Management and Budget. Revisions to the standards for the classification of federal data on race and ethnicity. *Federal Register* 1997;62:58781–58790. <http://go.usa.gov/vSdR>. Accessed October 15, 2015.
5. Office of Management and Budget. 2010 Standards for defining metropolitan and micropolitan statistical areas. *Federal Register* 2010;75(123):37246–39052. <http://go.usa.gov/vSpG>. Accessed October 15, 2015.
6. Office of Management and Budget. Revised definitions of metropolitan statistical areas, new definitions of micropolitan statistical areas and combined statistical areas, and guidance on uses of the delineations of these areas. OMB Bulletin 13-01. <http://go.usa.gov/vSyB>. Published February 28, 2013. Accessed October 15, 2015.
7. CDC. Current Trends: Heterosexually acquired AIDS—United States, 1993. *MMWR* 1994;43(9):158, Fig 3.
8. Harrison KM, Kajese T, Hall HI, Song R. Risk factor redistribution of the national HIV/AIDS surveillance data: an alternative approach. *Public Health Rep* 2008;123(5):618–627.
9. Rubin DB. *Multiple Imputation for Nonresponse in Surveys*. New York: John Wiley & Sons Inc; 1987.
10. Song R, Green TA. An improved approach to accounting for reporting delay in case surveillance systems. *JP J Biostat* 2012;7(1):1–14.
11. U.S. Census Bureau. Population estimates [entire data set]. <http://www.census.gov/popest/data/>. Updated July 1, 2014. Accessed October 15, 2015.
12. U.S. Census Bureau. International Data Base. <http://go.usa.gov/vSUj>. Updated December 2013. Accessed October 15, 2015.

Table 1a. Diagnoses of HIV infection, by year of diagnosis and selected characteristics, 2010–2014—United States

	2010			2011			2012			2013			2014		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Age at diagnosis (yr)															
<13	232	238	0.4	192	198	0.4	240	250	0.5	180	191	0.4	159	174	0.3
13–14	42	43	0.5	43	45	0.5	44	46	0.6	40	43	0.5	33	35	0.4
15–19	2,071	2,118	9.6	2,002	2,068	9.5	1,875	1,964	9.2	1,689	1,792	8.5	1,664	1,828	8.7
20–24	7,079	7,245	33.4	7,069	7,311	33.0	7,157	7,489	33.1	7,035	7,483	32.7	7,144	7,868	34.3
25–29	6,366	6,520	30.8	6,346	6,563	30.8	6,470	6,777	31.7	6,711	7,151	33.1	7,114	7,870	35.8
30–34	5,504	5,639	28.1	5,272	5,455	26.6	5,472	5,729	27.4	5,235	5,574	26.2	5,449	6,026	28.0
35–39	5,046	5,171	25.8	4,463	4,622	23.6	4,176	4,374	22.4	4,031	4,288	21.8	4,212	4,662	23.4
40–44	5,230	5,361	25.6	4,800	4,971	23.6	4,433	4,646	22.1	3,997	4,257	20.4	3,799	4,196	20.4
45–49	4,851	4,972	22.0	4,595	4,758	21.5	4,317	4,527	20.8	3,995	4,268	20.1	3,647	4,021	19.3
50–54	3,510	3,602	16.1	3,364	3,487	15.4	3,219	3,377	14.9	3,024	3,235	14.3	2,928	3,242	14.4
55–59	2,077	2,132	10.8	1,999	2,072	10.2	1,923	2,019	9.7	2,047	2,184	10.3	1,949	2,166	10.1
60–64	1,064	1,091	6.4	1,072	1,111	6.2	1,052	1,106	6.2	1,098	1,170	6.5	960	1,069	5.8
≥65	790	810	2.0	816	848	2.0	819	861	2.0	867	930	2.1	819	914	2.0
Race/ethnicity															
American Indian/Alaska Native	174	177	7.8	159	163	7.1	187	193	8.4	178	186	8.0	208	222	9.5
Asian	708	727	4.9	774	802	5.3	809	848	5.4	809	859	5.3	941	1,046	6.2
Black/African American	20,461	20,987	55.2	19,345	20,064	52.3	18,632	19,581	50.5	17,993	19,252	49.2	17,592	19,540	49.4
Hispanic/Latino ^b	9,072	9,291	18.3	8,919	9,230	17.8	8,954	9,372	17.7	8,829	9,386	17.3	9,227	10,201	18.4
Native Hawaiian/Other Pacific Islander	57	58	11.7	56	58	11.4	58	60	11.6	54	56	10.6	53	58	10.6
White	11,864	12,135	6.1	11,376	11,738	5.9	11,259	11,752	5.9	10,914	11,581	5.9	10,967	12,025	6.1
Multiple races	1,526	1,565	27.7	1,404	1,456	25.0	1,298	1,358	22.6	1,172	1,246	20.1	889	982	15.4
Transmission category															
Male adult or adolescent															
Male-to-male sexual contact	21,834	27,031	—	21,828	27,001	—	21,758	27,588	—	21,594	27,642	—	21,566	29,418	—
Injection drug use	1,307	2,115	—	1,080	1,819	—	916	1,642	—	874	1,575	—	809	1,590	—
Male-to-male sexual contact and injection drug use	1,238	1,562	—	1,100	1,393	—	1,036	1,342	—	924	1,216	—	870	1,217	—
Heterosexual contact ^c	2,891	4,111	—	2,739	3,883	—	2,500	3,617	—	2,398	3,545	—	2,049	3,285	—
Other ^d	6,771	52	—	6,245	50	—	6,491	69	—	6,163	57	—	6,891	60	—
Subtotal	34,041	34,871	27.9	32,992	34,146	27.0	32,701	34,259	26.9	31,953	34,034	26.4	32,185	35,571	27.4

Table 1a. Diagnoses of HIV infection, by year of diagnosis and selected characteristics, 2010–2014—United States (cont)

	2010			2011			2012			2013			2014		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Female adult or adolescent															
Injection drug use	803	1,455	—	672	1,284	—	617	1,178	—	559	1,073	—	495	1,045	—
Heterosexual contact ^c	4,740	8,340	—	4,307	7,833	—	3,905	7,439	—	3,782	7,213	—	3,282	7,242	—
Other ^d	4,046	36	—	3,870	49	—	3,734	39	—	3,475	55	—	3,756	41	—
Subtotal	9,589	9,831	7.5	8,849	9,166	6.9	8,256	8,656	6.5	7,816	8,340	6.2	7,533	8,328	6.1
Child (<13 yrs at diagnosis)															
Perinatal	181	185	—	142	147	—	168	175	—	119	127	—	115	127	—
Other ^e	51	53	—	50	51	—	72	75	—	61	64	—	44	48	—
Subtotal	232	238	0.4	192	198	0.4	240	250	0.5	180	191	0.4	159	174	0.3
Region of residence															
Northeast	8,381	8,597	15.5	7,800	8,087	14.5	7,646	8,039	14.4	7,236	7,750	13.8	7,137	7,953	14.2
Midwest	5,554	5,664	8.5	5,424	5,580	8.3	5,507	5,717	8.5	5,376	5,654	8.4	5,099	5,529	8.2
South	21,997	22,550	19.6	21,316	22,079	19.0	20,469	21,480	18.3	20,131	21,508	18.1	20,065	22,196	18.5
West	7,930	8,129	11.3	7,493	7,764	10.7	7,575	7,929	10.8	7,206	7,654	10.3	7,576	8,395	11.2
Total^f	43,862	44,940	14.5	42,033	43,510	14.0	41,197	43,165	13.7	39,949	42,566	13.4	39,877	44,073	13.8

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^b Hispanics/Latinos can be of any race.

^c Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^d Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 1b. Diagnoses of HIV infection, by year of diagnosis and selected characteristics, 2010–2014—United States and 6 dependent areas

	2010			2011			2012			2013			2014		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Age at diagnosis (yr)															
<13	236	242	0.5	194	200	0.4	240	250	0.5	180	191	0.4	160	176	0.3
13–14	42	43	0.5	44	46	0.5	45	47	0.6	41	44	0.5	33	35	0.4
15–19	2,100	2,148	9.6	2,023	2,090	9.5	1,894	1,984	9.2	1,707	1,811	8.4	1,686	1,854	8.7
20–24	7,155	7,324	33.3	7,147	7,393	32.9	7,263	7,603	33.2	7,123	7,578	32.7	7,245	7,983	34.4
25–29	6,462	6,620	30.9	6,443	6,665	30.9	6,566	6,880	31.8	6,791	7,238	33.1	7,195	7,963	35.8
30–34	5,616	5,755	28.3	5,361	5,549	26.7	5,571	5,835	27.5	5,322	5,669	26.3	5,507	6,093	28.0
35–39	5,144	5,273	25.9	4,547	4,711	23.7	4,243	4,445	22.5	4,098	4,362	21.9	4,288	4,750	23.5
40–44	5,351	5,487	25.9	4,903	5,080	23.8	4,529	4,749	22.3	4,065	4,331	20.5	3,878	4,288	20.6
45–49	4,964	5,090	22.2	4,701	4,870	21.7	4,415	4,632	21.1	4,078	4,358	20.3	3,716	4,101	19.4
50–54	3,580	3,674	16.2	3,452	3,580	15.7	3,292	3,455	15.1	3,104	3,323	14.5	2,973	3,294	14.4
55–59	2,123	2,179	10.9	2,047	2,123	10.3	1,960	2,059	9.8	2,092	2,234	10.4	1,991	2,215	10.2
60–64	1,100	1,129	6.6	1,097	1,138	6.3	1,075	1,131	6.3	1,134	1,210	6.6	986	1,099	5.8
≥65	819	840	2.0	841	874	2.1	857	902	2.1	893	959	2.1	835	933	2.0
Race/ethnicity															
American Indian/Alaska Native	174	177	—	159	163	—	187	193	—	178	186	—	208	222	—
Asian	709	728	—	774	802	—	809	848	—	809	859	—	942	1,047	—
Black/African American	20,475	21,001	—	19,361	20,081	—	18,636	19,585	—	18,006	19,266	—	17,607	19,557	—
Hispanic/Latino ^b	9,882	10,133	—	9,668	10,020	—	9,700	10,171	—	9,492	10,110	—	9,822	10,887	—
Native Hawaiian/Other Pacific Islander	60	61	—	57	59	—	59	61	—	54	56	—	53	58	—
White	11,866	12,137	—	11,377	11,739	—	11,260	11,753	—	10,917	11,584	—	10,971	12,029	—
Multiple races	1,526	1,565	—	1,404	1,456	—	1,299	1,360	—	1,172	1,246	—	890	983	—
Transmission category															
Male adult or adolescent															
Male-to-male sexual contact	22,110	27,335	—	22,085	27,291	—	22,046	27,909	—	21,868	27,956	—	21,842	29,771	—
Injection drug use	1,429	2,257	—	1,178	1,940	—	999	1,742	—	947	1,667	—	859	1,672	—
Male-to-male sexual contact and injection drug use	1,264	1,592	—	1,116	1,412	—	1,054	1,363	—	939	1,232	—	873	1,222	—
Heterosexual contact ^c	3,032	4,268	—	2,875	4,038	—	2,635	3,769	—	2,514	3,680	—	2,144	3,412	—
Other ^d	6,815	52	—	6,294	51	—	6,521	69	—	6,196	57	—	6,959	61	—
Subtotal	34,650	35,504	28.0	33,548	34,733	27.2	33,255	34,853	27.0	32,464	34,592	26.5	32,677	36,138	27.5

Table 1b. Diagnoses of HIV infection, by year of diagnosis and selected characteristics, 2010–2014—United States and 6 dependent areas (cont)

	2010			2011			2012			2013			2014		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Female adult or adolescent															
Injection drug use	824	1,479	—	704	1,319	—	638	1,204	—	585	1,105	—	507	1,060	—
Heterosexual contact ^c	4,918	8,541	—	4,472	8,018	—	4,061	7,626	—	3,892	7,363	—	3,376	7,369	—
Other ^d	4,064	37	—	3,882	49	—	3,756	39	—	3,507	56	—	3,773	42	—
Subtotal	9,806	10,057	7.6	9,058	9,386	7.0	8,455	8,869	6.5	7,984	8,524	6.2	7,656	8,471	6.1
Child (<13 yrs at diagnosis)															
Perinatal	184	188	—	143	148	—	168	175	—	119	127	—	116	128	—
Other ^e	52	54	—	51	52	—	72	75	—	61	64	—	44	48	—
Subtotal	236	242	0.5	194	200	0.4	240	250	0.5	180	191	0.4	160	176	0.3
Region of residence															
Northeast	8,381	8,597	15.5	7,800	8,087	14.5	7,646	8,039	14.4	7,236	7,750	13.8	7,137	7,953	14.2
Midwest	5,554	5,664	8.5	5,424	5,580	8.3	5,507	5,717	8.5	5,376	5,654	8.4	5,099	5,529	8.2
South	21,997	22,550	19.6	21,316	22,079	19.0	20,469	21,480	18.3	20,131	21,508	18.1	20,065	22,196	18.5
West	7,930	8,129	11.3	7,493	7,764	10.7	7,575	7,929	10.8	7,206	7,654	10.3	7,576	8,395	11.2
U.S. dependent areas	830	863	21.0	767	810	19.8	753	807	20.0	679	741	18.6	616	711	18.0
Total^f	44,692	45,802	14.6	42,800	44,320	14.0	41,950	43,972	13.8	40,628	43,307	13.5	40,493	44,784	13.9

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited. Rates are not calculated by transmission category because of the lack of denominator data.

^b Hispanics/Latinos can be of any race.

^c Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^d Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 2a. Stage 3 (AIDS), by year of diagnosis and selected characteristics, 2010–2014 and cumulative—United States

	2010			2011			2012			2013			2014			Cumulative ^b	
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Est. No. ^a
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		No.	Rate		
Age at diagnosis (yr)																	
<13 ^c	21	21	0.0	15	15	0.0	10	10	0.0	8	8	0.0	94	104	0.2	9,561	9,588
13–14	49	50	0.6	40	42	0.5	30	31	0.4	28	30	0.4	20	22	0.3	1,462	1,477
15–19	451	461	2.1	426	440	2.0	349	363	1.7	401	423	2.0	207	227	1.1	8,909	9,027
20–24	1,989	2,038	9.4	1,970	2,042	9.2	1,895	1,981	8.8	2,001	2,112	9.2	1,339	1,467	6.4	51,371	51,986
25–29	2,886	2,956	14.0	2,776	2,877	13.5	2,730	2,853	13.3	2,809	2,974	13.8	2,303	2,531	11.5	141,002	142,069
30–34	3,305	3,382	16.9	3,133	3,242	15.8	3,236	3,377	16.1	2,931	3,099	14.6	2,368	2,598	12.1	224,940	226,254
35–39	3,717	3,804	18.9	3,205	3,318	16.9	2,919	3,047	15.6	2,848	3,010	15.3	2,330	2,556	12.8	243,597	245,042
40–44	4,324	4,425	21.2	3,776	3,909	18.6	3,431	3,577	17.0	3,095	3,270	15.7	2,424	2,659	12.9	204,493	205,990
45–49	4,292	4,392	19.4	3,965	4,107	18.5	3,650	3,805	17.5	3,344	3,534	16.6	2,611	2,866	13.7	139,973	141,345
50–54	3,168	3,240	14.5	2,908	3,009	13.3	2,888	3,010	13.3	2,807	2,963	13.1	2,253	2,468	10.9	83,736	84,738
55–59	1,761	1,801	9.1	1,802	1,863	9.2	1,771	1,848	8.9	1,796	1,893	8.9	1,548	1,702	7.9	46,497	47,105
60–64	936	957	5.6	906	936	5.3	991	1,033	5.8	1,031	1,087	6.0	835	913	4.9	24,762	25,078
≥65	742	759	1.9	721	744	1.8	722	751	1.7	769	811	1.8	713	784	1.7	20,882	21,137
Race/ethnicity																	
American Indian/Alaska Native	116	118	5.2	103	105	4.6	94	96	4.2	91	94	4.0	90	95	4.0	3,498	3,523
Asian ^d	363	372	2.5	362	375	2.5	354	369	2.3	358	379	2.3	323	352	2.1	9,689	9,815
Black/African American	13,419	13,745	36.2	12,363	12,816	33.4	12,030	12,574	32.4	11,548	12,219	31.2	9,119	10,045	25.4	499,734	504,354
Hispanic/Latino ^e	5,617	5,745	11.3	5,237	5,415	10.4	5,015	5,222	9.8	4,868	5,132	9.5	3,924	4,279	7.7	217,650	219,578
Native Hawaiian/ Other Pacific Islander	39	40	8.0	31	32	6.3	26	27	5.2	27	28	5.2	18	19	3.5	842	850
White	6,953	7,102	3.6	6,483	6,693	3.4	6,111	6,354	3.2	6,005	6,328	3.2	4,850	5,303	2.7	436,952	439,455
Multiple races	1,134	1,165	20.6	1,064	1,108	19.0	992	1,044	17.4	971	1,034	16.7	721	803	12.6	32,820	33,260
Transmission category																	
Male adult or adolescent																	
Male-to-male sexual contact	11,695	14,448	—	11,238	13,863	—	10,826	13,664	—	10,781	13,785	—	8,180	11,277	—	529,245	586,385
Injection drug use	1,485	2,072	—	1,240	1,773	—	1,116	1,649	—	957	1,477	—	812	1,268	—	165,386	187,372
Male-to-male sexual contact and injection drug use	1,177	1,431	—	1,018	1,242	—	929	1,166	—	774	1,005	—	555	782	—	78,824	85,290
Heterosexual contact ^f	2,255	3,092	—	2,112	2,896	—	1,946	2,728	—	1,891	2,730	—	1,453	2,211	—	66,501	84,200
Other ^g	4,063	118	—	3,613	127	—	3,699	117	—	3,696	133	—	3,243	85	—	107,624	11,627
Subtotal	20,675	21,161	16.9	19,221	19,901	15.8	18,516	19,324	15.2	18,099	19,130	14.9	14,243	15,624	12.0	947,580	954,875

Table 2a. Stage 3 (AIDS), by year of diagnosis and selected characteristics, 2010–2014 and cumulative—United States (cont)

	2010			2011			2012			2013			2014			Cumulative ^b	
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Est. No. ^a
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		No.	Rate		
Female adult or adolescent																	
Injection drug use	981	1,469	—	864	1,302	—	781	1,205	—	671	1,089	—	523	913	—	75,745	90,413
Heterosexual contact ^f	3,603	5,502	—	3,316	5,195	—	3,113	5,039	—	2,899	4,839	—	2,203	4,150	—	114,383	149,980
Other ^g	2,361	133	—	2,227	131	—	2,202	109	—	2,191	147	—	1,982	106	—	53,916	5,979
Subtotal	6,945	7,105	5.4	6,407	6,628	5.0	6,096	6,353	4.8	5,761	6,075	4.5	4,708	5,168	3.8	244,044	246,372
Child (<13 yrs at diagnosis)																	
Perinatal	15	15	—	12	12	—	8	8	—	8	8	—	83	92	—	8,690	8,715
Other ^h	6	6	—	3	3	—	2	2	—	0	0	—	11	12	—	871	874
Subtotal	21	21	0.0	15	15	0.0	10	10	0.0	8	8	0.0	94	104	0.2	9,561	9,588
Region of residence																	
Northeast	5,631	5,782	10.4	5,153	5,356	9.6	4,765	4,992	8.9	4,360	4,636	8.3	3,409	3,793	6.8	351,863	354,392
Midwest	3,365	3,431	5.1	3,248	3,341	5.0	3,106	3,215	4.8	2,952	3,087	4.6	2,431	2,627	3.9	127,677	128,670
South	13,747	14,059	12.2	12,949	13,402	11.5	12,608	13,155	11.2	12,667	13,376	11.3	10,003	10,965	9.2	483,173	487,539
West	4,898	5,016	7.0	4,293	4,445	6.1	4,143	4,325	5.9	3,889	4,115	5.5	3,202	3,511	4.7	238,472	240,235
Totalⁱ	27,641	28,287	9.1	25,643	26,545	8.5	24,622	25,687	8.2	23,868	25,214	8.0	19,045	20,896	6.6	1,201,185	1,210,835

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^b From the beginning of the epidemic through 2014.

^c The criteria for stage 3 (AIDS) classification among pediatric cases were expanded in 2014 (see Technical Notes).

^d Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^e Hispanics/Latinos can be of any race.

^f Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^g Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^h Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

ⁱ Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 2b. Stage 3 (AIDS), by year of diagnosis and selected characteristics, 2010–2014 and cumulative—United States and 6 dependent areas

	2010			2011			2012			2013			2014			Cumulative ^b	
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Est. No. ^a
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		No.	Rate		
Age at diagnosis (yr)																	
<13 ^c	21	21	0.0	15	15	0.0	10	10	0.0	8	8	0.0	94	104	0.2	9,986	10,014
13–14	49	50	0.6	41	43	0.5	31	32	0.4	29	31	0.4	20	22	0.3	1,520	1,536
15–19	455	465	2.1	434	449	2.0	350	364	1.7	405	427	2.0	211	233	1.1	9,147	9,268
20–24	2,007	2,056	9.3	1,986	2,059	9.2	1,914	2,001	8.7	2,025	2,138	9.2	1,353	1,485	6.4	52,793	53,421
25–29	2,919	2,990	14.0	2,810	2,913	13.5	2,762	2,887	13.3	2,838	3,005	13.7	2,321	2,555	11.5	145,224	146,317
30–34	3,356	3,435	16.9	3,179	3,290	15.8	3,276	3,419	16.1	2,974	3,145	14.6	2,400	2,641	12.1	231,855	233,212
35–39	3,791	3,880	19.1	3,277	3,393	17.1	2,973	3,103	15.7	2,899	3,064	15.4	2,375	2,615	13.0	250,834	252,332
40–44	4,416	4,520	21.3	3,847	3,983	18.7	3,502	3,651	17.1	3,151	3,330	15.7	2,474	2,726	13.1	210,319	211,872
45–49	4,399	4,502	19.6	4,060	4,205	18.7	3,731	3,890	17.7	3,418	3,614	16.8	2,659	2,928	13.8	143,924	145,345
50–54	3,240	3,314	14.7	2,989	3,093	13.5	2,947	3,072	13.4	2,874	3,034	13.3	2,293	2,520	11.0	86,093	87,132
55–59	1,802	1,843	9.2	1,834	1,896	9.2	1,810	1,889	9.0	1,841	1,942	9.0	1,583	1,749	8.0	47,907	48,541
60–64	966	988	5.7	921	952	5.3	1,011	1,054	5.8	1,056	1,114	6.1	853	938	5.0	25,555	25,886
≥65	762	779	1.9	753	777	1.9	752	783	1.8	791	834	1.8	728	803	1.7	21,837	22,105
Race/ethnicity																	
American Indian/Alaska Native	116	118	—	103	105	—	94	96	—	91	94	—	90	95	—	3,499	3,524
Asian ^d	363	372	—	363	376	—	354	369	—	358	379	—	324	353	—	9,747	9,873
Black/African American	13,428	13,755	—	12,369	12,823	—	12,040	12,585	—	11,552	12,223	—	9,126	10,053	—	500,208	504,833
Hispanic/Latino ^e	6,146	6,288	—	5,731	5,928	—	5,448	5,676	—	5,304	5,599	—	4,233	4,689	—	252,778	255,036
Native Hawaiian/ Other Pacific Islander	41	42	—	32	33	—	27	28	—	27	28	—	18	19	—	858	867
White	6,955	7,104	—	6,483	6,693	—	6,114	6,357	—	6,006	6,330	—	4,851	5,305	—	437,071	439,575
Multiple races	1,134	1,165	—	1,065	1,109	—	992	1,044	—	971	1,034	—	722	804	—	32,833	33,274
Transmission category																	
Male adult or adolescent																	
Male-to-male sexual contact	11,811	14,570	—	11,355	13,988	—	10,946	13,792	—	10,910	13,926	—	8,266	11,394	—	535,337	592,621
Injection drug use	1,614	2,207	—	1,340	1,881	—	1,208	1,748	—	1,031	1,560	—	869	1,346	—	178,756	201,009
Male-to-male sexual contact and injection drug use	1,205	1,460	—	1,037	1,263	—	946	1,185	—	785	1,017	—	564	794	—	81,355	87,850
Heterosexual contact ^f	2,354	3,196	—	2,217	3,007	—	2,019	2,807	—	1,985	2,833	—	1,516	2,301	—	70,522	88,352
Other ^g	4,074	121	—	3,625	130	—	3,709	120	—	3,706	135	—	3,253	85	—	108,156	11,828
Subtotal	21,058	21,554	17.0	19,574	20,268	15.9	18,828	19,651	15.2	18,417	19,471	14.9	14,468	15,920	12.1	974,126	981,660

Table 2b. Stage 3 (AIDS), by year of diagnosis and selected characteristics, 2010–2014 and cumulative—United States and 6 dependent areas (cont)

	2010			2011			2012			2013			2014			Cumulative ^b	
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Est. No. ^a
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		No.	Rate		
Female adult or adolescent																	
Injection drug use	1,012	1,501	—	891	1,331	—	804	1,230	—	697	1,118	—	546	944	—	78,776	93,515
Heterosexual contact ^f	3,728	5,632	—	3,432	5,318	—	3,218	5,154	—	2,989	4,940	—	2,264	4,241	—	119,886	155,667
Other ^g	2,364	134	—	2,234	135	—	2,209	111	—	2,198	150	—	1,992	108	—	54,220	6,125
Subtotal	7,104	7,268	5.5	6,557	6,784	5.1	6,231	6,494	4.8	5,884	6,207	4.5	4,802	5,293	3.8	252,882	255,307
Child (<13 yrs at diagnosis)																	
Perinatal	15	15	—	12	12	—	8	8	—	8	8	—	83	92	—	9,089	9,115
Other ^h	6	6	—	3	3	—	2	2	—	0	0	—	11	12	—	897	900
Subtotal	21	21	0.0	15	15	0.0	10	10	0.0	8	8	0.0	94	104	0.2	9,986	10,014
Region of residence																	
Northeast	5,631	5,782	10.4	5,153	5,356	9.6	4,765	4,992	8.9	4,360	4,636	8.3	3,409	3,793	6.8	351,863	354,392
Midwest	3,365	3,431	5.1	3,248	3,341	5.0	3,106	3,215	4.8	2,952	3,087	4.6	2,431	2,627	3.9	127,677	128,670
South	13,747	14,059	12.2	12,949	13,402	11.5	12,608	13,155	11.2	12,667	13,376	11.3	10,003	10,965	9.2	483,173	487,539
West	4,898	5,016	7.0	4,293	4,445	6.1	4,143	4,325	5.9	3,889	4,115	5.5	3,202	3,511	4.7	238,472	240,235
U.S. dependent areas	542	556	13.5	503	523	12.8	447	469	11.6	441	473	11.9	319	422	10.7	35,809	36,146
Totalⁱ	28,183	28,844	9.2	26,146	27,067	8.6	25,069	26,156	8.2	24,309	25,687	8.0	19,364	21,318	6.6	1,236,994	1,246,981

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited. Rates are not calculated by transmission category because of the lack of denominator data.

^b From the beginning of the epidemic through 2014.

^c The criteria for stage 3 (AIDS) classification among pediatric cases were expanded in 2014 (see Technical Notes).

^d Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^e Hispanics/Latinos can be of any race.

^f Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^g Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^h Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

ⁱ Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 3a. Diagnoses of HIV infection, by race/ethnicity and selected characteristics, 2014—United States

	American Indian/ Alaska Native			Asian			Black/African American			Hispanic/Latino ^a			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^b			Estimated ^b			Estimated ^b			Estimated ^b			Estimated ^b			Estimated ^b			Estimated ^b					
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^c	Rate
Age at diagnosis (yr)																								
<13	3	3	0.7	10	11	0.4	98	108	1.5	16	18	0.1	1	1	1.0	26	28	0.1	5	5	0.2	159	174	0.3
13–14	0	0	0.0	3	3	0.9	17	18	1.5	5	5	0.3	0	0	0.0	8	8	0.2	0	0	0.0	33	35	0.4
15–19	8	9	4.7	20	22	2.2	1,044	1,148	38.2	330	364	7.8	2	2	5.9	203	221	1.9	57	62	9.4	1,664	1,828	8.7
20–24	37	39	19.4	141	157	12.7	3,896	4,296	124.8	1,574	1,740	36.4	8	9	18.6	1,295	1,416	11.2	193	211	36.7	7,144	7,868	34.3
25–29	46	49	27.7	171	190	13.5	3,280	3,646	122.6	1,711	1,893	42.6	6	7	13.1	1,738	1,904	15.2	162	180	41.5	7,114	7,870	35.8
30–34	32	34	21.3	163	182	12.6	2,104	2,345	85.1	1,527	1,687	38.0	8	9	17.9	1,510	1,654	13.4	105	116	31.0	5,449	6,026	28.0
35–39	12	13	8.8	137	152	11.1	1,568	1,744	68.9	1,194	1,326	32.1	10	11	26.2	1,198	1,314	11.5	93	103	33.4	4,212	4,662	23.4
40–44	23	25	16.9	116	129	9.4	1,397	1,556	60.0	967	1,070	27.7	7	8	20.8	1,198	1,309	10.6	91	100	36.2	3,799	4,196	20.4
45–49	20	21	14.2	78	85	7.1	1,338	1,490	57.3	813	893	26.3	5	6	16.1	1,315	1,440	10.8	78	86	35.6	3,647	4,021	19.3
50–54	19	20	12.5	52	58	5.2	1,114	1,242	45.6	515	569	19.4	2	2	6.3	1,171	1,289	8.4	55	61	25.5	2,928	3,242	14.4
55–59	6	7	4.6	25	28	2.8	908	1,014	40.6	297	330	14.2	1	1	3.6	682	753	4.9	30	33	15.8	1,949	2,166	10.1
60–64	1	1	0.9	11	12	1.4	443	501	25.2	158	173	10.0	2	2	10.4	335	368	2.7	10	11	7.0	960	1,069	5.8
≥65	1	1	0.5	14	16	0.8	385	431	10.7	120	133	3.7	1	1	2.5	288	321	0.9	10	11	3.5	819	914	2.0
Transmission category																								
Male adult or adolescent																								
Male-to-male sexual contact	120	142	—	586	798	—	7,777	11,201	—	5,737	7,552	—	33	45	—	6,791	9,008	—	522	672	—	21,566	29,418	—
Injection drug use	9	11	—	13	20	—	208	679	—	202	395	—	1	2	—	358	455	—	18	27	—	809	1,590	—
Male-to-male sexual contact and injection drug use	5	7	—	13	17	—	137	288	—	172	251	—	1	1	—	511	611	—	31	40	—	870	1,217	—
Heterosexual contact ^d	8	10	—	37	60	—	1,212	2,108	—	441	623	—	0	1	—	305	422	—	46	63	—	2,049	3,285	—
Other ^e	17	0	—	159	3	—	3,537	29	—	1,439	11	—	10	0	—	1,621	17	—	108	1	—	6,891	60	—
Subtotal	159	170	18.3	808	898	13.4	12,871	14,305	94.0	7,991	8,832	41.5	45	49	22.0	9,586	10,513	12.6	725	803	40.6	32,185	35,571	27.4

Table 3a. Diagnoses of HIV infection, by race/ethnicity and selected characteristics, 2014—United States (cont)

	American Indian/ Alaska Native			Asian			Black/African American			Hispanic/Latino ^a			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^b			Estimated ^b			Estimated ^b			Estimated ^b			Estimated ^b			Estimated ^b			Estimated ^b					
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^c	Rate
Female adult or adolescent																								
Injection drug use	9	13	—	1	7	—	134	451	—	87	183	—	0	1	—	247	360	—	17	31	—	495	1,045	—
Heterosexual contact ^d	21	36	—	59	129	—	1,988	4,654	—	564	1,159	—	3	7	—	570	1,115	—	77	142	—	3,282	7,242	—
Other ^e	16	0	—	63	1	—	2,501	23	—	569	9	—	4	0	—	538	8	—	65	1	—	3,756	41	—
Subtotal	46	49	5.1	123	137	1.8	4,623	5,128	30.0	1,220	1,350	6.5	7	8	3.5	1,355	1,483	1.7	159	174	8.2	7,533	8,328	6.1
Child (<13 yrs at diagnosis)																								
Perinatal	2	2	—	9	10	—	75	83	—	12	13	—	1	1	—	13	14	—	3	3	—	115	127	—
Other ^f	1	1	—	1	1	—	23	25	—	4	4	—	0	0	—	13	14	—	2	2	—	44	48	—
Subtotal	3	3	0.7	10	11	0.4	98	108	1.5	16	18	0.1	1	1	1.0	26	28	0.1	5	5	0.2	159	174	0.3
Region of residence																								
Northeast	10	11	8.5	182	205	5.8	3,055	3,385	53.6	1,952	2,190	28.3	4	5	21.9	1,720	1,922	5.1	214	236	26.7	7,137	7,953	14.2
Midwest	25	26	6.6	88	97	4.8	2,467	2,677	38.2	617	680	13.3	3	3	10.5	1,773	1,910	3.7	126	136	11.2	5,099	5,529	8.2
South	56	61	7.9	210	231	6.0	10,798	12,068	53.0	3,767	4,097	20.1	13	15	17.5	4,818	5,276	7.6	403	449	21.5	20,065	22,196	18.5
West	117	124	11.9	461	513	7.0	1,272	1,411	41.2	2,891	3,234	14.6	33	35	8.6	2,656	2,917	7.6	146	161	7.3	7,576	8,395	11.2
Total^g	208	222	9.5	941	1,046	6.2	17,592	19,540	49.4	9,227	10,201	18.4	53	58	10.6	10,967	12,025	6.1	889	982	15.4	39,877	44,073	13.8

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Hispanics/Latinos can be of any race.

^b Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^c Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 3b. Diagnoses of HIV infection, by race/ethnicity and selected characteristics, 2014—United States and 6 dependent areas

	American Indian/ Alaska Native		Asian		Black/ African American		Hispanic/Latino ^a		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total	
	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^{b,c}
Age at diagnosis (yr)																
<13	3	3	10	11	98	108	17	19	1	1	26	28	5	5	160	176
13–14	0	0	3	3	17	18	5	5	0	0	8	8	0	0	33	35
15–19	8	9	20	22	1,045	1,150	351	388	2	2	203	221	57	62	1,686	1,854
20–24	37	39	141	157	3,896	4,296	1,674	1,854	8	9	1,296	1,418	193	211	7,245	7,983
25–29	46	49	171	190	3,281	3,648	1,790	1,984	6	7	1,739	1,905	162	180	7,195	7,963
30–34	32	34	163	182	2,106	2,347	1,582	1,750	8	9	1,511	1,655	105	116	5,507	6,093
35–39	12	13	137	152	1,569	1,745	1,269	1,412	10	11	1,198	1,314	93	103	4,288	4,750
40–44	23	25	116	129	1,398	1,557	1,045	1,161	7	8	1,198	1,309	91	100	3,878	4,288
45–49	20	21	78	85	1,339	1,491	881	971	5	6	1,315	1,440	78	86	3,716	4,101
50–54	19	20	53	59	1,118	1,247	554	614	2	2	1,172	1,290	55	61	2,973	3,294
55–59	6	7	25	28	910	1,016	336	376	1	1	682	753	31	35	1,991	2,215
60–64	1	1	11	12	444	502	183	202	2	2	335	368	10	11	986	1,099
≥65	1	1	14	16	386	432	135	150	1	1	288	321	10	11	835	933
Transmission category																
Male adult or adolescent																
Male-to-male sexual contact	120	142	587	799	7,781	11,207	6,003	7,893	33	45	6,795	9,012	523	674	21,842	29,771
Injection drug use	9	11	13	20	208	682	252	474	1	2	358	455	18	27	859	1,672
Male-to-male sexual contact and injection drug use	5	7	13	17	137	289	175	257	1	1	511	611	31	40	873	1,222
Heterosexual contact ^d	8	10	37	60	1,215	2,113	533	744	0	1	305	422	46	63	2,144	3,412
Other ^e	17	0	159	3	3,542	29	1,502	11	10	0	1,621	17	108	1	6,959	61
Subtotal	159	170	809	900	12,883	14,319	8,465	9,379	45	49	9,590	10,518	726	804	32,677	36,138

Table 3b. Diagnoses of HIV infection, by race/ethnicity and selected characteristics, 2014—United States and 6 dependent areas (cont)

	American Indian/ Alaska Native		Asian		Black/ African American		Hispanic/Latino ^a		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total	
	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^{b,c}
Female adult or adolescent																
Injection drug use	9	13	1	7	134	451	99	198	0	1	247	360	17	31	507	1,060
Heterosexual contact ^d	21	36	59	129	1,989	4,657	657	1,282	3	7	570	1,115	77	142	3,376	7,369
Other ^e	16	0	63	1	2,503	23	584	9	4	0	538	8	65	1	3,773	42
Subtotal	46	49	123	137	4,626	5,131	1,340	1,490	7	8	1,355	1,483	159	174	7,656	8,471
Child (<13 yrs at diagnosis)																
Perinatal	2	2	9	10	75	83	13	15	1	1	13	14	3	3	116	128
Other ^f	1	1	1	1	23	25	4	4	0	0	13	14	2	2	44	48
Subtotal	3	3	10	11	98	108	17	19	1	1	26	28	5	5	160	176
Region of residence																
Northeast	10	11	182	205	3,055	3,385	1,952	2,190	4	5	1,720	1,922	214	236	7,137	7,953
Midwest	25	26	88	97	2,467	2,677	617	680	3	3	1,773	1,910	126	136	5,099	5,529
South	56	61	210	231	10,798	12,068	3,767	4,097	13	15	4,818	5,276	403	449	20,065	22,196
West	117	124	461	513	1,272	1,411	2,891	3,234	33	35	2,656	2,917	146	161	7,576	8,395
U.S. dependent areas	0	0	1	1	15	17	595	687	0	0	4	5	1	1	616	711
Total^g	208	222	942	1,047	17,607	19,557	9,822	10,887	53	58	10,971	12,029	890	983	40,493	44,784

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Hispanics/Latinos can be of any race.

^b Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting.

^c Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 4a. Stage 3 (AIDS), by race/ethnicity and selected characteristics, 2014—United States

	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total				
	Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c				
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^d
Age at diagnosis (yr)																										
<13	1	1	0.2	1	1	0.0	56	62	0.9	11	12	0.1	0	0	0.0	6	6	0.0	19	21	0.9	94	104	0.2		
13–14	0	0	0.0	1	1	0.3	13	15	1.2	4	4	0.2	0	0	0.0	0	0	0.0	2	2	0.7	20	22	0.3		
15–19	0	0	0.0	0	0	0.0	145	159	5.3	27	30	0.6	0	0	0.0	17	19	0.2	18	20	3.0	207	227	1.1		
20–24	4	4	2.1	20	21	1.7	798	878	25.5	274	299	6.3	0	0	0.0	184	199	1.6	59	65	11.3	1,339	1,467	6.4		
25–29	10	10	5.9	38	41	2.9	1,240	1,372	46.1	534	582	13.1	1	1	2.2	391	425	3.4	89	99	22.8	2,303	2,531	11.5		
30–34	14	15	9.1	46	50	3.5	1,130	1,245	45.2	600	653	14.7	5	5	11.3	502	550	4.5	71	79	21.2	2,368	2,598	12.1		
35–39	8	9	5.9	57	63	4.6	1,036	1,140	45.0	626	684	16.5	3	3	7.9	512	561	4.9	88	97	31.3	2,330	2,556	12.8		
40–44	19	20	13.7	58	64	4.6	1,073	1,182	45.6	535	584	15.1	2	2	5.7	650	709	5.8	87	98	35.4	2,424	2,659	12.9		
45–49	11	12	7.9	53	57	4.8	1,120	1,232	47.4	503	549	16.2	3	3	9.1	826	907	6.8	95	106	43.8	2,611	2,866	13.7		
50–54	14	15	9.0	21	23	2.1	1,004	1,103	40.5	357	389	13.2	1	1	3.2	774	844	5.5	82	93	38.5	2,253	2,468	10.9		
55–59	5	5	3.5	15	16	1.6	745	821	32.9	230	251	10.8	1	1	3.7	492	540	3.5	60	67	31.5	1,548	1,702	7.9		
60–64	3	3	2.6	4	4	0.5	413	454	22.8	128	139	8.0	2	2	9.3	258	281	2.0	27	30	18.3	835	913	4.9		
≥65	1	1	0.5	9	10	0.5	346	380	9.5	95	103	2.9	0	0	0.0	238	263	0.7	24	27	8.3	713	784	1.7		
Transmission category																										
Male adult or adolescent																										
Male-to-male sexual contact	38	46	—	163	234	—	3,036	4,343	—	1,973	2,665	—	11	13	—	2,656	3,564	—	303	412	—	8,180	11,277	—		
Injection drug use	10	11	—	6	10	—	317	587	—	200	313	—	0	0	—	246	300	—	33	47	—	812	1,268	—		
Male-to-male sexual contact and injection drug use	5	7	—	9	11	—	135	233	—	100	144	—	1	1	—	269	335	—	36	51	—	555	782	—		
Heterosexual contact ^e	4	5	—	20	31	—	899	1,451	—	279	388	—	1	1	—	203	268	—	47	67	—	1,453	2,211	—		
Other ^f	9	0	—	67	2	—	1,644	36	—	689	20	—	2	0	—	731	23	—	101	5	—	3,243	85	—		
Subtotal	66	69	7.5	265	289	4.3	6,031	6,649	43.7	3,241	3,530	16.6	15	16	7.2	4,105	4,489	5.4	520	582	29.4	14,243	15,624	12.0		

Table 4a. Stage 3 (AIDS), by race/ethnicity and selected characteristics, 2014—United States (cont)

	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c					
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^d	Rate
Female adult or adolescent																								
Injection drug use	7	9	—	2	4	—	220	462	—	87	144	—	0	0	—	178	247	—	29	46	—	523	913	—
Heterosexual contact ^e	10	15	—	32	58	—	1,442	2,809	—	313	568	—	1	3	—	315	550	—	90	146	—	2,203	4,150	—
Other ^f	6	0	—	23	1	—	1,370	62	—	272	25	—	2	0	—	246	10	—	63	8	—	1,982	106	—
Subtotal	23	24	2.5	57	62	0.8	3,032	3,334	19.5	672	737	3.5	3	3	1.4	739	808	0.9	182	200	9.4	4,708	5,168	3.8
Child (<13 yrs at diagnosis)																								
Perinatal	0	0	—	0	0	—	52	58	—	7	8	—	0	0	—	6	6	—	18	20	—	83	92	—
Other ^g	1	1	—	1	1	—	4	4	—	4	4	—	0	0	—	0	0	—	1	1	—	11	12	—
Subtotal	1	1	0.2	1	1	0.0	56	62	0.9	11	12	0.1	0	0	0.0	6	6	0.0	19	21	0.9	94	104	0.2
Region of residence																								
Northeast	2	2	1.6	54	60	1.7	1,569	1,739	27.5	908	1,011	13.1	0	0	0.0	716	801	2.1	160	181	20.4	3,409	3,793	6.8
Midwest	10	10	2.6	44	46	2.3	1,181	1,280	18.3	259	282	5.5	1	1	3.4	829	889	1.7	107	118	9.7	2,431	2,627	3.9
South	22	24	3.1	76	83	2.2	5,804	6,402	28.1	1,603	1,721	8.5	4	4	4.9	2,134	2,333	3.3	360	398	19.1	10,003	10,965	9.2
West	56	59	5.6	149	163	2.2	565	624	18.2	1,154	1,266	5.7	13	14	3.4	1,171	1,280	3.3	94	105	4.8	3,202	3,511	4.7
Total^h	90	95	4.0	323	352	2.1	9,119	10,045	25.4	3,924	4,279	7.7	18	19	3.5	4,850	5,303	2.7	721	803	12.6	19,045	20,896	6.6

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^g Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^h Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 4b. Stage 3 (AIDS), by race/ethnicity and selected characteristics, 2014—United States and 6 dependent areas

	American Indian/ Alaska Native		Asian ^a		Black/ African American		Hispanic/Latino ^b		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Age at diagnosis (yr)																
<13	1	1	1	1	56	62	11	12	0	0	6	6	19	21	94	104
13–14	0	0	1	1	13	15	4	4	0	0	0	0	2	2	20	22
15–19	0	0	0	0	145	159	31	36	0	0	17	19	18	20	211	233
20–24	4	4	20	21	798	878	288	317	0	0	184	199	59	65	1,353	1,485
25–29	10	10	38	41	1,240	1,372	552	606	1	1	391	425	89	99	2,321	2,555
30–34	14	15	46	50	1,130	1,245	632	696	5	5	502	550	71	79	2,400	2,641
35–39	8	9	57	63	1,036	1,140	671	743	3	3	512	561	88	97	2,375	2,615
40–44	19	20	58	64	1,074	1,183	584	650	2	2	650	709	87	98	2,474	2,726
45–49	11	12	53	57	1,121	1,234	550	609	3	3	826	907	95	106	2,659	2,928
50–54	14	15	22	24	1,008	1,108	391	434	1	1	775	845	82	93	2,293	2,520
55–59	5	5	15	16	745	821	264	297	1	1	492	540	61	68	1,583	1,749
60–64	3	3	4	4	414	456	145	163	2	2	258	281	27	30	853	938
≥65	1	1	9	10	346	380	110	122	0	0	238	263	24	27	728	803
Transmission category																
Male adult or adolescent																
Male-to-male sexual contact	38	46	164	235	3,037	4,344	2,055	2,776	11	13	2,657	3,565	304	414	8,266	11,394
Injection drug use	10	11	6	10	317	589	257	390	0	0	246	300	33	47	869	1,346
Male-to-male sexual contact and injection drug use	5	7	9	11	135	233	109	156	1	1	269	335	36	51	564	794
Heterosexual contact ^e	4	5	20	31	901	1,455	340	474	1	1	203	268	47	67	1,516	2,301
Other ^f	9	0	67	2	1,647	36	696	20	2	0	731	23	101	5	3,253	85
Subtotal	66	69	266	290	6,037	6,656	3,457	3,815	15	16	4,106	4,490	521	583	14,468	15,920

Table 4b. Stage 3 (AIDS), by race/ethnicity and selected characteristics, 2014—United States and 6 dependent areas (cont)

	American Indian/ Alaska Native		Asian ^a		Black/ African American		Hispanic/Latino ^b		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Female adult or adolescent																
Injection drug use	7	9	2	4	220	462	110	175	0	0	178	247	29	46	546	944
Heterosexual contact ^e	10	15	32	58	1,442	2,811	374	658	1	3	315	550	90	146	2,264	4,241
Other ^f	6	0	23	1	1,371	62	281	27	2	0	246	10	63	8	1,992	108
Subtotal	23	24	57	62	3,033	3,335	765	861	3	3	739	808	182	200	4,802	5,293
Child (<13 yrs at diagnosis)																
Perinatal	0	0	0	0	52	58	7	8	0	0	6	6	18	20	83	92
Other ^g	1	1	1	1	4	4	4	4	0	0	0	0	1	1	11	12
Subtotal	1	1	1	1	56	62	11	12	0	0	6	6	19	21	94	104
Region of residence																
Northeast	2	2	54	60	1,569	1,739	908	1,011	0	0	716	801	160	181	3,409	3,793
Midwest	10	10	44	46	1,181	1,280	259	282	1	1	829	889	107	118	2,431	2,627
South	22	24	76	83	5,804	6,402	1,603	1,721	4	4	2,134	2,333	360	398	10,003	10,965
West	56	59	149	163	565	624	1,154	1,266	13	14	1,171	1,280	94	105	3,202	3,511
U.S. dependent areas	0	0	1	1	7	9	309	409	0	0	1	1	1	2	319	422
Total^h	90	95	324	353	9,126	10,053	4,233	4,689	18	19	4,851	5,305	722	804	19,364	21,318

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^g Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^h Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 5a. Diagnoses of HIV infection, by transmission category, race/ethnicity, and selected characteristics, 2014—United States

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
American Indian/Alaska Native												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	0	0	0	0	0	0	0	0	0	0	0	0
15–19	5	5	0	0	0	0	0	0	0	0	5	5
20–24	32	36	1	1	1	1	0	0	2	0	36	38
25–29	33	36	2	2	0	0	0	0	1	0	36	38
30–34	18	22	0	0	0	1	3	3	3	0	24	26
35–39	6	6	0	0	0	0	1	1	0	0	7	7
40–44	9	13	3	4	0	1	1	2	5	0	18	19
45–49	9	12	0	0	2	2	2	2	3	0	16	17
50–54	7	8	3	3	2	2	0	0	1	0	13	14
55–59	1	2	0	0	0	0	1	1	1	0	3	4
60–64	0	0	0	0	0	0	0	0	0	0	0	0
≥65	0	1	0	0	0	0	0	0	1	0	1	1
Region of residence												
Northeast	6	6	0	0	0	0	2	2	0	0	8	8
Midwest	11	13	1	1	3	3	0	0	2	0	17	18
South	29	37	4	5	0	1	0	1	7	0	40	44
West	74	84	4	5	2	3	6	7	8	0	94	100
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	0	0	0	0	0
15–19	—	—	0	0	—	—	2	3	1	0	3	3
20–24	—	—	0	0	—	—	0	1	1	0	1	1
25–29	—	—	1	2	—	—	5	8	4	0	10	11
30–34	—	—	3	3	—	—	2	5	3	0	8	8
35–39	—	—	1	1	—	—	2	4	2	0	5	5
40–44	—	—	1	1	—	—	2	4	2	0	5	5
45–49	—	—	1	1	—	—	2	3	1	0	4	4
50–54	—	—	1	1	—	—	4	5	1	0	6	6
55–59	—	—	0	0	—	—	2	3	1	0	3	3
60–64	—	—	1	1	—	—	0	0	0	0	1	1
≥65	—	—	0	0	—	—	0	0	0	0	0	0
Region of residence												
Northeast	—	—	0	0	—	—	0	2	2	0	2	2
Midwest	—	—	1	2	—	—	3	7	4	0	8	9
South	—	—	3	5	—	—	4	12	9	0	16	17
West	—	—	5	6	—	—	14	16	1	0	20	21
Subtotal^e	120	142	18	24	5	7	29	46	33	1	205	219

Table 5a. Diagnoses of HIV infection, by transmission category, race/ethnicity, and selected characteristics, 2014—United States (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Asian												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	2	2	0	0	0	0	0	0	1	1	3	3
15–19	14	18	1	1	1	1	1	1	2	0	19	21
20–24	117	141	0	0	2	3	1	1	11	0	131	145
25–29	121	162	0	0	1	2	5	7	26	0	153	171
30–34	106	139	4	5	2	3	12	14	20	0	144	161
35–39	82	111	4	5	3	4	5	9	21	0	115	128
40–44	65	92	2	3	4	5	3	6	21	0	95	106
45–49	44	66	1	3	0	0	4	6	20	0	69	75
50–54	25	39	1	2	0	0	4	6	13	0	43	48
55–59	6	13	0	0	0	0	1	4	8	0	15	17
60–64	2	7	0	0	0	0	0	1	5	0	7	8
≥65	2	9	0	1	0	0	1	5	11	1	14	16
Region of residence												
Northeast	118	157	3	6	0	0	11	16	28	1	160	180
Midwest	42	63	1	2	4	5	7	10	19	0	73	80
South	104	159	7	9	2	3	6	13	49	1	168	185
West	322	420	2	3	7	9	13	21	63	1	407	453
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	0	0	0	0	0
15–19	—	—	0	0	—	—	1	1	0	0	1	1
20–24	—	—	0	0	—	—	5	11	5	0	10	11
25–29	—	—	0	0	—	—	9	19	9	0	18	19
30–34	—	—	0	1	—	—	10	20	9	0	19	22
35–39	—	—	0	1	—	—	13	23	9	0	22	24
40–44	—	—	0	1	—	—	9	22	12	0	21	24
45–49	—	—	0	1	—	—	5	9	4	0	9	10
50–54	—	—	0	1	—	—	4	9	5	0	9	10
55–59	—	—	1	2	—	—	1	9	8	1	10	11
60–64	—	—	0	0	—	—	2	4	2	0	4	4
≥65	—	—	0	0	—	—	0	0	0	0	0	0
Region of residence												
Northeast	—	—	0	1	—	—	13	23	9	0	22	25
Midwest	—	—	1	1	—	—	8	13	4	0	13	14
South	—	—	0	1	—	—	20	38	16	0	36	39
West	—	—	0	3	—	—	18	54	34	1	52	58
Subtotal^e	586	798	14	27	13	17	96	188	222	5	931	1,035

Table 5a. Diagnoses of HIV infection, by transmission category, race/ethnicity, and selected characteristics, 2014—United States (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Black/African American												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	2	5	0	0	0	0	1	1	6	3	9	10
15–19	658	856	1	8	7	17	21	35	153	6	840	923
20–24	2,677	3,460	10	52	26	60	108	175	581	7	3,402	3,754
25–29	1,955	2,628	16	63	24	53	140	213	527	2	2,662	2,959
30–34	909	1,367	12	57	24	44	149	227	423	1	1,517	1,696
35–39	462	791	11	59	11	26	139	220	358	1	981	1,097
40–44	334	622	17	67	11	23	159	264	355	1	876	978
45–49	327	585	29	89	9	19	138	257	349	1	852	951
50–54	207	404	30	84	10	20	151	264	295	2	693	773
55–59	139	276	39	97	8	15	98	205	247	1	531	594
60–64	60	119	24	52	5	8	54	121	123	2	266	301
≥65	47	91	19	48	2	4	54	126	120	2	242	271
Region of residence												
Northeast	1,158	1,575	49	229	31	49	275	461	582	2	2,095	2,316
Midwest	1,344	1,781	20	51	17	34	126	189	401	15	1,908	2,070
South	4,630	6,934	119	361	62	164	752	1,356	2,321	10	7,884	8,825
West	645	912	20	38	27	42	59	100	233	1	984	1,093
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	2	4	6	4	8	9
15–19	—	—	2	11	—	—	74	205	128	10	204	226
20–24	—	—	6	30	—	—	241	509	247	3	494	542
25–29	—	—	11	48	—	—	252	638	355	2	618	688
30–34	—	—	11	44	—	—	266	604	310	1	587	649
35–39	—	—	10	47	—	—	258	599	319	1	587	647
40–44	—	—	12	50	—	—	239	528	270	0	521	578
45–49	—	—	23	59	—	—	200	480	263	0	486	539
50–54	—	—	22	58	—	—	173	411	226	0	421	470
55–59	—	—	21	55	—	—	156	365	200	0	377	420
60–64	—	—	11	29	—	—	70	171	96	0	177	200
≥65	—	—	5	20	—	—	57	140	81	0	143	160
Region of residence												
Northeast	—	—	31	127	—	—	323	921	594	7	948	1,055
Midwest	—	—	15	47	—	—	233	533	289	2	537	582
South	—	—	73	239	—	—	1,343	2,938	1,446	9	2,862	3,185
West	—	—	15	38	—	—	89	262	172	4	276	304
Subtotal^e	7,777	11,201	342	1,130	137	288	3,200	6,761	6,038	52	17,494	19,433

Table 5a. Diagnoses of HIV infection, by transmission category, race/ethnicity, and selected characteristics, 2014—United States (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Hispanic/Latino^f												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	2	2	0	0	0	0	0	0	0	0	2	2
15–19	226	287	0	3	8	12	3	4	43	2	280	309
20–24	1,224	1,497	12	22	25	36	29	35	154	4	1,444	1,594
25–29	1,274	1,589	21	35	35	49	49	63	189	1	1,568	1,736
30–34	996	1,309	32	55	38	52	67	87	228	1	1,361	1,504
35–39	692	942	29	57	23	34	72	95	200	0	1,016	1,128
40–44	506	729	31	60	16	26	69	98	205	0	827	914
45–49	417	583	25	52	15	21	64	92	160	0	681	748
50–54	219	324	22	45	4	8	44	69	115	0	404	446
55–59	92	153	20	40	6	9	22	35	74	0	214	237
60–64	52	82	8	17	1	2	9	20	41	0	111	121
≥65	37	56	2	10	1	2	13	23	30	1	83	91
Region of residence												
Northeast	1,070	1,429	53	153	24	36	101	150	329	0	1,577	1,769
Midwest	411	536	5	12	7	12	26	36	92	2	541	598
South	2,311	3,010	67	125	59	93	240	330	601	7	3,278	3,564
West	1,945	2,578	77	105	82	111	74	107	417	2	2,595	2,902
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	1	3	2	3	3
15–19	—	—	2	7	—	—	17	45	31	4	50	55
20–24	—	—	12	19	—	—	59	126	59	0	130	145
25–29	—	—	7	17	—	—	66	139	70	2	143	157
30–34	—	—	12	23	—	—	86	159	68	0	166	183
35–39	—	—	13	26	—	—	87	172	78	0	178	198
40–44	—	—	14	25	—	—	60	131	66	0	140	156
45–49	—	—	6	15	—	—	70	130	56	0	132	145
50–54	—	—	10	21	—	—	51	102	50	0	111	123
55–59	—	—	6	16	—	—	34	77	43	0	83	93
60–64	—	—	3	7	—	—	19	45	25	0	47	52
≥65	—	—	2	7	—	—	15	34	20	0	37	41
Region of residence												
Northeast	—	—	33	80	—	—	133	339	209	2	375	421
Midwest	—	—	7	12	—	—	41	70	28	1	76	83
South	—	—	26	53	—	—	267	464	186	4	479	521
West	—	—	21	39	—	—	123	286	146	1	290	325
Subtotal^e	5,737	7,552	289	579	172	251	1,005	1,782	2,008	19	9,211	10,183

Table 5a. Diagnoses of HIV infection, by transmission category, race/ethnicity, and selected characteristics, 2014—United States (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Native Hawaiian/Other Pacific Islander												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	0	0	0	0	0	0	0	0	0	0	0	0
15–19	1	1	0	0	0	0	0	0	0	0	1	1
20–24	5	8	0	0	0	0	0	0	2	0	7	8
25–29	6	7	0	0	0	0	0	0	0	0	6	7
30–34	5	6	0	0	1	1	0	0	1	0	7	7
35–39	6	9	0	0	0	0	0	0	3	0	9	10
40–44	3	6	0	0	0	0	0	0	3	0	6	7
45–49	5	6	0	0	0	0	0	0	0	0	5	6
50–54	1	1	0	0	0	0	0	0	0	0	1	1
55–59	0	0	0	0	0	0	0	0	0	0	0	0
60–64	0	1	1	1	0	0	0	0	1	0	2	2
≥65	1	1	0	0	0	0	0	0	0	0	1	1
Region of residence												
Northeast	1	2	1	2	0	0	0	0	1	0	3	3
Midwest	3	3	0	0	0	0	0	0	0	0	3	3
South	5	12	0	0	0	0	0	1	6	0	11	13
West	24	28	0	0	1	1	0	0	3	0	28	30
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	0	0	0	0	0
15–19	—	—	0	0	—	—	0	1	1	0	1	1
20–24	—	—	0	0	—	—	1	1	0	0	1	1
25–29	—	—	0	0	—	—	0	0	0	0	0	0
30–34	—	—	0	0	—	—	0	1	1	0	1	1
35–39	—	—	0	0	—	—	0	1	1	0	1	1
40–44	—	—	0	0	—	—	1	1	0	0	1	1
45–49	—	—	0	0	—	—	0	0	0	0	0	0
50–54	—	—	0	0	—	—	1	1	0	0	1	1
55–59	—	—	0	0	—	—	0	1	1	0	1	1
60–64	—	—	0	0	—	—	0	0	0	0	0	0
≥65	—	—	0	0	—	—	0	0	0	0	0	0
Region of residence												
Northeast	—	—	0	0	—	—	0	1	1	0	1	1
Midwest	—	—	0	0	—	—	0	0	0	0	0	0
South	—	—	0	0	—	—	1	1	0	0	1	1
West	—	—	0	0	—	—	2	5	3	0	5	6
Subtotal^e	33	45	1	3	1	1	3	7	14	0	52	57

Table 5a. Diagnoses of HIV infection, by transmission category, race/ethnicity, and selected characteristics, 2014—United States (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
White												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	2	3	0	0	0	0	0	0	2	1	4	4
15–19	126	158	2	2	9	10	3	3	20	0	160	173
20–24	914	1,130	30	34	56	65	14	18	127	3	1,141	1,249
25–29	1,175	1,474	46	54	118	133	29	35	181	1	1,549	1,696
30–34	942	1,220	55	63	90	105	32	39	186	2	1,305	1,429
35–39	738	983	43	51	68	80	22	30	171	1	1,042	1,144
40–44	698	950	57	69	61	72	43	54	189	0	1,048	1,145
45–49	794	1,091	41	53	48	63	49	68	234	3	1,166	1,279
50–54	693	955	45	61	35	46	52	71	205	1	1,030	1,133
55–59	372	540	22	34	17	23	25	44	144	1	580	641
60–64	194	271	14	20	9	12	17	24	65	1	299	328
≥65	143	234	3	14	0	3	19	36	97	4	262	291
Region of residence												
Northeast	1,082	1,430	51	89	48	63	54	80	254	1	1,489	1,663
Midwest	1,142	1,473	49	59	83	98	43	56	255	8	1,572	1,694
South	2,885	3,920	141	172	192	234	138	199	781	7	4,137	4,532
West	1,682	2,184	117	135	188	217	70	86	331	1	2,388	2,623
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	2	3	2	1	4	4
15–19	—	—	10	13	—	—	16	34	17	0	43	47
20–24	—	—	24	36	—	—	63	130	67	1	154	167
25–29	—	—	44	59	—	—	75	148	70	1	189	208
30–34	—	—	43	60	—	—	89	163	73	2	205	225
35–39	—	—	36	46	—	—	68	123	52	0	156	169
40–44	—	—	28	41	—	—	66	122	56	0	150	163
45–49	—	—	29	43	—	—	55	118	65	0	149	162
50–54	—	—	18	29	—	—	65	125	58	1	141	156
55–59	—	—	10	19	—	—	48	93	44	0	102	112
60–64	—	—	3	7	—	—	15	32	18	0	36	40
≥65	—	—	2	6	—	—	8	24	16	0	26	30
Region of residence												
Northeast	—	—	40	71	—	—	87	184	102	2	229	256
Midwest	—	—	28	41	—	—	90	166	78	2	196	210
South	—	—	118	161	—	—	300	568	251	2	669	731
West	—	—	61	87	—	—	93	197	107	2	261	286
Subtotal^e	6,791	9,008	605	815	511	611	875	1,537	2,159	25	10,941	11,996

Table 5a. Diagnoses of HIV infection, by transmission category, race/ethnicity, and selected characteristics, 2014—United States (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Multiple races												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	0	0	0	0	0	0	0	0	0	0	0	0
15–19	42	50	0	0	1	1	1	1	4	0	48	53
20–24	149	170	1	1	9	10	3	3	7	0	169	185
25–29	116	140	5	5	6	7	6	7	10	0	143	159
30–34	60	78	2	3	4	6	6	8	13	0	85	94
35–39	43	65	3	4	3	4	9	11	17	0	75	84
40–44	41	55	2	3	4	5	3	4	11	0	61	67
45–49	33	46	1	3	3	4	11	14	13	0	61	67
50–54	23	41	1	3	0	1	3	4	17	0	44	49
55–59	13	20	2	3	1	1	3	5	7	0	26	29
60–64	0	3	1	2	0	0	0	1	4	0	5	6
≥65	2	5	0	1	0	0	1	3	5	0	8	9
Region of residence												
Northeast	121	154	4	9	4	5	17	23	28	0	174	192
Midwest	77	95	2	2	2	3	5	7	14	0	100	108
South	233	309	5	8	8	12	19	26	52	0	317	355
West	91	114	7	8	17	19	5	7	14	0	134	148
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	0	0	0	0	0
15–19	—	—	1	1	—	—	5	8	3	0	9	9
20–24	—	—	0	1	—	—	14	24	10	0	24	26
25–29	—	—	1	2	—	—	10	19	8	0	19	21
30–34	—	—	1	2	—	—	13	19	6	0	20	22
35–39	—	—	1	2	—	—	10	18	7	0	18	20
40–44	—	—	4	7	—	—	13	27	13	0	30	33
45–49	—	—	4	7	—	—	6	12	7	0	17	19
50–54	—	—	2	4	—	—	3	8	6	0	11	12
55–59	—	—	2	3	—	—	0	2	2	0	4	4
60–64	—	—	0	0	—	—	2	5	3	0	5	5
≥65	—	—	1	1	—	—	1	1	0	0	2	2
Region of residence												
Northeast	—	—	5	12	—	—	16	31	18	0	39	43
Midwest	—	—	0	1	—	—	15	25	9	0	24	26
South	—	—	9	14	—	—	42	77	33	0	84	91
West	—	—	3	4	—	—	4	9	5	0	12	13
Subtotal^e	522	672	35	58	31	40	123	205	173	1	884	976

Table 5a. Diagnoses of HIV infection, by transmission category, race/ethnicity, and selected characteristics, 2014—United States (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Total												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	8	12	0	0	0	0	1	1	9	5	18	19
15–19	1,072	1,374	4	15	26	43	29	45	222	9	1,353	1,485
20–24	5,118	6,442	54	112	119	174	155	233	884	14	6,330	6,974
25–29	4,680	6,034	90	159	184	244	229	325	934	4	6,117	6,766
30–34	3,036	4,141	105	183	159	211	269	378	874	4	4,443	4,917
35–39	2,029	2,905	90	177	108	148	248	366	770	2	3,245	3,598
40–44	1,656	2,467	112	206	96	133	278	428	789	2	2,931	3,236
45–49	1,629	2,389	97	200	77	109	268	440	779	4	2,850	3,142
50–54	1,175	1,772	102	198	51	77	254	415	646	3	2,228	2,464
55–59	623	1,004	83	173	32	48	150	294	481	2	1,369	1,521
60–64	308	483	48	92	15	22	80	166	239	3	690	766
≥65	232	396	24	74	3	8	88	193	264	8	611	680
Region of residence												
Northeast	3,556	4,753	161	488	107	153	460	733	1,222	5	5,506	6,132
Midwest	3,030	3,966	78	128	116	155	207	298	783	25	4,214	4,572
South	10,197	14,380	343	680	323	507	1,155	1,925	3,817	26	15,835	17,518
West	4,783	6,320	227	294	324	402	227	328	1,069	4	6,630	7,348
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	4	8	11	7	15	16
15–19	—	—	15	32	—	—	115	297	181	14	311	343
20–24	—	—	42	87	—	—	383	802	389	5	814	894
25–29	—	—	64	128	—	—	417	971	516	5	997	1,104
30–34	—	—	70	135	—	—	466	972	470	3	1,006	1,109
35–39	—	—	61	123	—	—	438	940	468	1	967	1,064
40–44	—	—	59	125	—	—	390	835	419	1	868	960
45–49	—	—	63	127	—	—	338	752	396	1	797	879
50–54	—	—	53	114	—	—	301	662	346	2	700	778
55–59	—	—	40	95	—	—	241	549	299	1	580	645
60–64	—	—	18	45	—	—	108	257	144	1	270	303
≥65	—	—	10	34	—	—	81	199	117	1	208	233
Region of residence												
Northeast	—	—	109	291	—	—	572	1,502	935	12	1,616	1,804
Midwest	—	—	52	104	—	—	390	814	412	6	854	924
South	—	—	229	473	—	—	1,977	4,098	1,941	16	4,147	4,587
West	—	—	105	177	—	—	343	828	468	8	916	1,013
Total^e	21,566	29,418	1,304	2,635	870	1,217	5,331	10,527	10,647	102	39,718	43,899

Abbreviation: IDU, injection drug use.

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

^f Hispanics/Latinos can be of any race.

Table 5b. Diagnoses of HIV infection, by transmission category, race/ethnicity, and selected characteristics, 2014—United States and 6 dependent areas

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
American Indian/Alaska Native												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	0	0	0	0	0	0	0	0	0	0	0	0
15–19	5	5	0	0	0	0	0	0	0	0	5	5
20–24	32	36	1	1	1	1	0	0	2	0	36	38
25–29	33	36	2	2	0	0	0	0	1	0	36	38
30–34	18	22	0	0	0	1	3	3	3	0	24	26
35–39	6	6	0	0	0	0	1	1	0	0	7	7
40–44	9	13	3	4	0	1	1	2	5	0	18	19
45–49	9	12	0	0	2	2	2	2	3	0	16	17
50–54	7	8	3	3	2	2	0	0	1	0	13	14
55–59	1	2	0	0	0	0	1	1	1	0	3	4
60–64	0	0	0	0	0	0	0	0	0	0	0	0
≥65	0	1	0	0	0	0	0	0	1	0	1	1
Region of residence												
Northeast	6	6	0	0	0	0	2	2	0	0	8	8
Midwest	11	13	1	1	3	3	0	0	2	0	17	18
South	29	37	4	5	0	1	0	1	7	0	40	44
West	74	84	4	5	2	3	6	7	8	0	94	100
U.S. dependent areas	0	0	0	0	0	0	0	0	0	0	0	0
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	0	0	0	0	0
15–19	—	—	0	0	—	—	2	3	1	0	3	3
20–24	—	—	0	0	—	—	0	1	1	0	1	1
25–29	—	—	1	2	—	—	5	8	4	0	10	11
30–34	—	—	3	3	—	—	2	5	3	0	8	8
35–39	—	—	1	1	—	—	2	4	2	0	5	5
40–44	—	—	1	1	—	—	2	4	2	0	5	5
45–49	—	—	1	1	—	—	2	3	1	0	4	4
50–54	—	—	1	1	—	—	4	5	1	0	6	6
55–59	—	—	0	0	—	—	2	3	1	0	3	3
60–64	—	—	1	1	—	—	0	0	0	0	1	1
≥65	—	—	0	0	—	—	0	0	0	0	0	0
Region of residence												
Northeast	—	—	0	0	—	—	0	2	2	0	2	2
Midwest	—	—	1	2	—	—	3	7	4	0	8	9
South	—	—	3	5	—	—	4	12	9	0	16	17
West	—	—	5	6	—	—	14	16	1	0	20	21
U.S. dependent areas	—	—	0	0	—	—	0	0	0	0	0	0
Subtotal^e	120	142	18	24	5	7	29	46	33	1	205	219

Table 5b. Diagnoses of HIV infection, by transmission category, race/ethnicity, and selected characteristics, 2014—United States and 6 dependent areas (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Asian												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	2	2	0	0	0	0	0	0	1	1	3	3
15–19	14	18	1	1	1	1	1	1	2	0	19	21
20–24	117	141	0	0	2	3	1	1	11	0	131	145
25–29	121	162	0	0	1	2	5	7	26	0	153	171
30–34	106	139	4	5	2	3	12	14	20	0	144	161
35–39	82	111	4	5	3	4	5	9	21	0	115	128
40–44	65	92	2	3	4	5	3	6	21	0	95	106
45–49	44	66	1	3	0	0	4	6	20	0	69	75
50–54	26	40	1	2	0	0	4	6	13	0	44	49
55–59	6	13	0	0	0	0	1	4	8	0	15	17
60–64	2	7	0	0	0	0	0	1	5	0	7	8
≥65	2	9	0	1	0	0	1	5	11	1	14	16
Region of residence												
Northeast	118	157	3	6	0	0	11	16	28	1	160	180
Midwest	42	63	1	2	4	5	7	10	19	0	73	80
South	104	159	7	9	2	3	6	13	49	1	168	185
West	322	420	2	3	7	9	13	21	63	1	407	453
U.S. dependent areas	1	1	0	0	0	0	0	0	0	0	1	1
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	0	0	0	0	0
15–19	—	—	0	0	—	—	1	1	0	0	1	1
20–24	—	—	0	0	—	—	5	11	5	0	10	11
25–29	—	—	0	0	—	—	9	19	9	0	18	19
30–34	—	—	0	1	—	—	10	20	9	0	19	22
35–39	—	—	0	1	—	—	13	23	9	0	22	24
40–44	—	—	0	1	—	—	9	22	12	0	21	24
45–49	—	—	0	1	—	—	5	9	4	0	9	10
50–54	—	—	0	1	—	—	4	9	5	0	9	10
55–59	—	—	1	2	—	—	1	9	8	1	10	11
60–64	—	—	0	0	—	—	2	4	2	0	4	4
≥65	—	—	0	0	—	—	0	0	0	0	0	0
Region of residence												
Northeast	—	—	0	1	—	—	13	23	9	0	22	25
Midwest	—	—	1	1	—	—	8	13	4	0	13	14
South	—	—	0	1	—	—	20	38	16	0	36	39
West	—	—	0	3	—	—	18	54	34	1	52	58
U.S. dependent areas	—	—	0	0	—	—	0	0	0	0	0	0
Subtotal^e	587	799	14	27	13	17	96	188	222	5	932	1,036

Table 5b. Diagnoses of HIV infection, by transmission category, race/ethnicity, and selected characteristics, 2014—United States and 6 dependent areas (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Black/African American												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	2	5	0	0	0	0	1	1	6	3	9	10
15–19	659	857	1	8	7	17	21	35	153	6	841	924
20–24	2,677	3,460	10	52	26	60	108	175	581	7	3,402	3,754
25–29	1,955	2,628	16	64	24	53	140	213	528	2	2,663	2,960
30–34	910	1,368	12	57	24	44	149	227	423	1	1,518	1,697
35–39	463	792	11	59	11	26	139	220	358	1	982	1,099
40–44	334	622	17	68	11	23	159	265	356	1	877	979
45–49	328	586	29	89	9	19	138	257	349	1	853	952
50–54	207	404	30	84	10	20	153	267	296	2	696	776
55–59	139	276	39	98	8	15	98	205	248	1	532	595
60–64	60	119	24	53	5	8	54	121	124	2	267	302
≥65	47	91	19	48	2	4	55	127	120	2	243	272
Region of residence												
Northeast	1,158	1,575	49	229	31	49	275	461	582	2	2,095	2,316
Midwest	1,344	1,781	20	51	17	34	126	189	401	15	1,908	2,070
South	4,630	6,934	119	361	62	164	752	1,356	2,321	10	7,884	8,825
West	645	912	20	38	27	42	59	100	233	1	984	1,093
U.S. dependent areas	4	5	0	3	0	0	3	5	5	0	12	14
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	2	4	6	4	8	9
15–19	—	—	2	11	—	—	74	205	128	10	204	226
20–24	—	—	6	30	—	—	241	509	247	3	494	542
25–29	—	—	11	48	—	—	252	638	355	2	618	688
30–34	—	—	11	44	—	—	266	605	311	1	588	650
35–39	—	—	10	47	—	—	258	599	319	1	587	647
40–44	—	—	12	50	—	—	239	528	270	0	521	578
45–49	—	—	23	59	—	—	200	480	263	0	486	539
50–54	—	—	22	58	—	—	174	413	226	0	422	471
55–59	—	—	21	55	—	—	156	366	201	0	378	422
60–64	—	—	11	29	—	—	70	171	96	0	177	200
≥65	—	—	5	20	—	—	57	140	81	0	143	160
Region of residence												
Northeast	—	—	31	127	—	—	323	921	594	7	948	1,055
Midwest	—	—	15	47	—	—	233	533	289	2	537	582
South	—	—	73	239	—	—	1,343	2,938	1,446	9	2,862	3,185
West	—	—	15	38	—	—	89	262	172	4	276	304
U.S. dependent areas	—	—	0	0	—	—	1	3	2	0	3	3
Subtotal^e	7,781	11,207	342	1,133	137	289	3,204	6,770	6,045	52	17,509	19,450

Table 5b. Diagnoses of HIV infection, by transmission category, race/ethnicity, and selected characteristics, 2014—United States and 6 dependent areas (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Hispanic/Latino^f												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	2	2	0	0	0	0	0	0	0	0	2	2
15–19	240	305	0	4	8	12	3	4	45	2	296	327
20–24	1,300	1,591	13	25	25	36	36	45	164	4	1,538	1,702
25–29	1,324	1,652	27	43	36	50	58	75	197	1	1,642	1,821
30–34	1,021	1,342	37	63	38	52	74	96	235	1	1,405	1,555
35–39	723	979	33	63	24	35	85	111	203	0	1,068	1,188
40–44	532	765	37	71	16	27	81	115	216	0	882	978
45–49	434	606	37	70	15	21	74	107	170	0	730	805
50–54	231	340	25	50	4	8	53	81	120	0	433	480
55–59	98	161	27	49	7	10	34	51	77	0	243	271
60–64	58	90	11	22	1	3	17	31	45	0	132	145
≥65	40	59	5	13	1	2	18	29	30	1	94	104
Region of residence												
Northeast	1,070	1,429	53	153	24	36	101	150	329	0	1,577	1,769
Midwest	411	536	5	12	7	12	26	36	92	2	541	598
South	2,311	3,010	67	125	59	93	240	330	601	7	3,278	3,564
West	1,945	2,578	77	105	82	111	74	107	417	2	2,595	2,902
U.S. dependent areas	266	341	50	79	3	6	92	121	63	0	474	546
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	1	3	2	3	3
15–19	—	—	2	7	—	—	19	50	34	4	55	61
20–24	—	—	13	21	—	—	63	131	60	1	136	152
25–29	—	—	7	17	—	—	70	144	71	2	148	163
30–34	—	—	13	24	—	—	95	171	69	0	177	195
35–39	—	—	17	31	—	—	104	194	80	0	201	224
40–44	—	—	17	29	—	—	78	154	68	0	163	183
45–49	—	—	7	16	—	—	84	150	60	0	151	167
50–54	—	—	11	22	—	—	59	112	51	0	121	134
55–59	—	—	7	17	—	—	43	87	43	0	93	104
60–64	—	—	3	7	—	—	23	49	25	0	51	57
≥65	—	—	2	7	—	—	19	39	20	0	41	46
Region of residence												
Northeast	—	—	33	80	—	—	133	339	209	2	375	421
Midwest	—	—	7	12	—	—	41	70	28	1	76	83
South	—	—	26	53	—	—	267	464	186	4	479	521
West	—	—	21	39	—	—	123	286	146	1	290	325
U.S. dependent areas	—	—	12	15	—	—	93	123	15	1	120	139
Subtotal^e	6,003	7,893	351	672	175	257	1,190	2,026	2,086	20	9,805	10,868

Table 5b. Diagnoses of HIV infection, by transmission category, race/ethnicity, and selected characteristics, 2014—United States and 6 dependent areas (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Native Hawaiian/Other Pacific Islander												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	0	0	0	0	0	0	0	0	0	0	0	0
15–19	1	1	0	0	0	0	0	0	0	0	1	1
20–24	5	8	0	0	0	0	0	0	2	0	7	8
25–29	6	7	0	0	0	0	0	0	0	0	6	7
30–34	5	6	0	0	1	1	0	0	1	0	7	7
35–39	6	9	0	0	0	0	0	0	3	0	9	10
40–44	3	6	0	0	0	0	0	0	3	0	6	7
45–49	5	6	0	0	0	0	0	0	0	0	5	6
50–54	1	1	0	0	0	0	0	0	0	0	1	1
55–59	0	0	0	0	0	0	0	0	0	0	0	0
60–64	0	1	1	1	0	0	0	0	1	0	2	2
≥65	1	1	0	0	0	0	0	0	0	0	1	1
Region of residence												
Northeast	1	2	1	2	0	0	0	0	1	0	3	3
Midwest	3	3	0	0	0	0	0	0	0	0	3	3
South	5	12	0	0	0	0	0	1	6	0	11	13
West	24	28	0	0	1	1	0	0	3	0	28	30
U.S. dependent areas	0	0	0	0	0	0	0	0	0	0	0	0
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	0	0	0	0	0
15–19	—	—	0	0	—	—	0	1	1	0	1	1
20–24	—	—	0	0	—	—	1	1	0	0	1	1
25–29	—	—	0	0	—	—	0	0	0	0	0	0
30–34	—	—	0	0	—	—	0	1	1	0	1	1
35–39	—	—	0	0	—	—	0	1	1	0	1	1
40–44	—	—	0	0	—	—	1	1	0	0	1	1
45–49	—	—	0	0	—	—	0	0	0	0	0	0
50–54	—	—	0	0	—	—	1	1	0	0	1	1
55–59	—	—	0	0	—	—	0	1	1	0	1	1
60–64	—	—	0	0	—	—	0	0	0	0	0	0
≥65	—	—	0	0	—	—	0	0	0	0	0	0
Region of residence												
Northeast	—	—	0	0	—	—	0	1	1	0	1	1
Midwest	—	—	0	0	—	—	0	0	0	0	0	0
South	—	—	0	0	—	—	1	1	0	0	1	1
West	—	—	0	0	—	—	2	5	3	0	5	6
U.S. dependent areas	—	—	0	0	—	—	0	0	0	0	0	0
Subtotal^e	33	45	1	3	1	1	3	7	14	0	52	57

Table 5b. Diagnoses of HIV infection, by transmission category, race/ethnicity, and selected characteristics, 2014—United States and 6 dependent areas (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
White												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	2	3	0	0	0	0	0	0	2	1	4	4
15–19	126	158	2	2	9	10	3	3	20	0	160	173
20–24	915	1,131	30	34	56	65	14	18	127	3	1,142	1,251
25–29	1,176	1,475	46	54	118	133	29	35	181	1	1,550	1,697
30–34	943	1,221	55	63	90	105	32	39	186	2	1,306	1,430
35–39	738	983	43	51	68	80	22	30	171	1	1,042	1,144
40–44	698	950	57	69	61	72	43	54	189	0	1,048	1,145
45–49	794	1,091	41	53	48	63	49	68	234	3	1,166	1,279
50–54	694	956	45	61	35	46	52	71	205	1	1,031	1,134
55–59	372	540	22	34	17	23	25	44	144	1	580	641
60–64	194	271	14	20	9	12	17	24	65	1	299	328
≥65	143	234	3	14	0	3	19	36	97	4	262	291
Region of residence												
Northeast	1,082	1,430	51	89	48	63	54	80	254	1	1,489	1,663
Midwest	1,142	1,473	49	59	83	98	43	56	255	8	1,572	1,694
South	2,885	3,920	141	172	192	234	138	199	781	7	4,137	4,532
West	1,682	2,184	117	135	188	217	70	86	331	1	2,388	2,623
U.S. dependent areas	4	5	0	0	0	0	0	0	0	0	4	5
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	2	3	2	1	4	4
15–19	—	—	10	13	—	—	16	34	17	0	43	47
20–24	—	—	24	36	—	—	63	130	67	1	154	167
25–29	—	—	44	59	—	—	75	148	70	1	189	208
30–34	—	—	43	60	—	—	89	163	73	2	205	225
35–39	—	—	36	46	—	—	68	123	52	0	156	169
40–44	—	—	28	41	—	—	66	122	56	0	150	163
45–49	—	—	29	43	—	—	55	118	65	0	149	162
50–54	—	—	18	29	—	—	65	125	58	1	141	156
55–59	—	—	10	19	—	—	48	93	44	0	102	112
60–64	—	—	3	7	—	—	15	32	18	0	36	40
≥65	—	—	2	6	—	—	8	24	16	0	26	30
Region of residence												
Northeast	—	—	40	71	—	—	87	184	102	2	229	256
Midwest	—	—	28	41	—	—	90	166	78	2	196	210
South	—	—	118	161	—	—	300	568	251	2	669	731
West	—	—	61	87	—	—	93	197	107	2	261	286
U.S. dependent areas	—	—	0	0	—	—	0	0	0	0	0	0
Subtotal^e	6,795	9,012	605	815	511	611	875	1,537	2,159	25	10,945	12,001

Table 5b. Diagnoses of HIV infection, by transmission category, race/ethnicity, and selected characteristics, 2014—United States and 6 dependent areas (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Multiple races												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	0	0	0	0	0	0	0	0	0	0	0	0
15–19	42	50	0	0	1	1	1	1	4	0	48	53
20–24	149	170	1	1	9	10	3	3	7	0	169	185
25–29	116	140	5	5	6	7	6	7	10	0	143	159
30–34	60	78	2	3	4	6	6	8	13	0	85	94
35–39	43	65	3	4	3	4	9	11	17	0	75	84
40–44	41	55	2	3	4	5	3	4	11	0	61	67
45–49	33	46	1	3	3	4	11	14	13	0	61	67
50–54	23	41	1	3	0	1	3	4	17	0	44	49
55–59	14	22	2	3	1	1	3	5	7	0	27	30
60–64	0	3	1	2	0	0	0	1	4	0	5	6
≥65	2	5	0	1	0	0	1	3	5	0	8	9
Region of residence												
Northeast	121	154	4	9	4	5	17	23	28	0	174	192
Midwest	77	95	2	2	2	3	5	7	14	0	100	108
South	233	309	5	8	8	12	19	26	52	0	317	355
West	91	114	7	8	17	19	5	7	14	0	134	148
U.S. dependent areas	1	1	0	0	0	0	0	0	0	0	1	1
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	0	0	0	0	0
15–19	—	—	1	1	—	—	5	8	3	0	9	9
20–24	—	—	0	1	—	—	14	24	10	0	24	26
25–29	—	—	1	2	—	—	10	19	8	0	19	21
30–34	—	—	1	2	—	—	13	19	6	0	20	22
35–39	—	—	1	2	—	—	10	18	7	0	18	20
40–44	—	—	4	7	—	—	13	27	13	0	30	33
45–49	—	—	4	7	—	—	6	12	7	0	17	19
50–54	—	—	2	4	—	—	3	8	6	0	11	12
55–59	—	—	2	3	—	—	0	2	2	0	4	4
60–64	—	—	0	0	—	—	2	5	3	0	5	5
≥65	—	—	1	1	—	—	1	1	0	0	2	2
Region of residence												
Northeast	—	—	5	12	—	—	16	31	18	0	39	43
Midwest	—	—	0	1	—	—	15	25	9	0	24	26
South	—	—	9	14	—	—	42	77	33	0	84	91
West	—	—	3	4	—	—	4	9	5	0	12	13
U.S. dependent areas	—	—	0	0	—	—	0	0	0	0	0	0
Subtotal^e	523	674	35	58	31	40	123	205	173	1	885	977

Table 5b. Diagnoses of HIV infection, by transmission category, race/ethnicity, and selected characteristics, 2014—United States and 6 dependent areas (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Total												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	8	12	0	0	0	0	1	1	9	5	18	19
15–19	1,087	1,393	4	16	26	43	29	45	224	9	1,370	1,505
20–24	5,195	6,537	55	115	119	174	162	242	894	14	6,425	7,083
25–29	4,731	6,099	96	168	185	246	238	337	943	4	6,193	6,853
30–34	3,063	4,176	110	190	159	211	276	388	881	4	4,489	4,970
35–39	2,061	2,944	94	182	109	150	261	382	773	2	3,298	3,659
40–44	1,682	2,503	118	218	96	133	290	445	801	2	2,987	3,301
45–49	1,647	2,413	109	218	77	109	278	455	789	4	2,900	3,200
50–54	1,189	1,790	105	204	51	77	265	430	652	3	2,262	2,504
55–59	630	1,013	90	183	33	49	162	309	485	2	1,400	1,558
60–64	314	490	51	98	15	22	88	178	244	3	712	792
≥65	235	400	27	78	3	8	94	200	264	8	623	694
Region of residence												
Northeast	3,556	4,753	161	488	107	153	460	733	1,222	5	5,506	6,132
Midwest	3,030	3,966	78	128	116	155	207	298	783	25	4,214	4,572
South	10,197	14,380	343	680	323	507	1,155	1,925	3,817	26	15,835	17,518
West	4,783	6,320	227	294	324	402	227	328	1,069	4	6,630	7,348
U.S. dependent areas	276	353	50	82	3	6	95	127	68	0	492	567
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	4	8	11	7	15	16
15–19	—	—	15	33	—	—	117	302	184	14	316	349
20–24	—	—	43	89	—	—	387	807	390	5	820	901
25–29	—	—	64	128	—	—	421	977	517	5	1,002	1,110
30–34	—	—	71	136	—	—	475	984	472	3	1,018	1,123
35–39	—	—	65	128	—	—	455	962	470	1	990	1,091
40–44	—	—	62	128	—	—	408	858	421	1	891	987
45–49	—	—	64	128	—	—	352	772	400	1	816	901
50–54	—	—	54	115	—	—	310	674	347	2	711	791
55–59	—	—	41	97	—	—	250	560	300	1	591	658
60–64	—	—	18	45	—	—	112	261	144	1	274	307
≥65	—	—	10	34	—	—	85	203	117	1	212	238
Region of residence												
Northeast	—	—	109	291	—	—	572	1,502	935	12	1,616	1,804
Midwest	—	—	52	104	—	—	390	814	412	6	854	924
South	—	—	229	473	—	—	1,977	4,098	1,941	16	4,147	4,587
West	—	—	105	177	—	—	343	828	468	8	916	1,013
U.S. dependent areas	—	—	12	15	—	—	94	127	17	1	123	143
Total^e	21,842	29,771	1,366	2,732	873	1,222	5,520	10,781	10,732	103	40,333	44,608

Abbreviation: IDU, injection drug use.

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

^f Hispanics/Latinos can be of any race.

Table 6a. Stage 3 (AIDS), by transmission category, race/ethnicity, and selected characteristics, 2014—United States

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
American Indian/Alaska Native												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	0	0	0	0	0	0	0	0	0	0	0	0
15–19	0	0	0	0	0	0	0	0	0	0	0	0
20–24	4	4	0	0	0	0	0	0	0	0	4	4
25–29	5	6	2	2	0	0	0	0	1	0	8	8
30–34	6	7	1	1	1	1	2	2	1	0	11	11
35–39	3	4	0	0	0	0	0	0	1	0	4	4
40–44	7	10	4	4	0	0	1	1	3	0	15	16
45–49	6	8	0	0	2	2	0	1	2	0	10	11
50–54	5	5	2	2	2	2	0	0	0	0	9	10
55–59	2	2	1	1	0	0	1	1	0	0	4	4
60–64	0	0	0	0	0	0	0	0	0	0	0	0
≥65	0	1	0	0	0	0	0	0	1	0	1	1
Region of residence												
Northeast	0	0	0	0	0	0	1	1	0	0	1	1
Midwest	3	4	1	1	2	2	0	0	1	0	7	7
South	8	11	4	5	1	1	0	1	3	0	16	17
West	27	32	5	5	2	3	3	3	5	0	42	44
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	0	0	0	0	0
15–19	—	—	0	0	—	—	0	0	0	0	0	0
20–24	—	—	0	0	—	—	0	0	0	0	0	0
25–29	—	—	0	0	—	—	1	2	1	0	2	2
30–34	—	—	1	1	—	—	0	2	2	0	3	3
35–39	—	—	2	3	—	—	0	2	2	0	4	4
40–44	—	—	2	3	—	—	1	2	1	0	4	4
45–49	—	—	0	0	—	—	1	1	0	0	1	1
50–54	—	—	1	1	—	—	4	4	0	0	5	5
55–59	—	—	0	0	—	—	1	1	0	0	1	1
60–64	—	—	1	1	—	—	2	2	0	0	3	3
≥65	—	—	0	0	—	—	0	0	0	0	0	0
Region of residence												
Northeast	—	—	0	0	—	—	1	1	0	0	1	1
Midwest	—	—	0	0	—	—	2	3	1	0	3	3
South	—	—	3	4	—	—	1	3	2	0	6	6
West	—	—	4	5	—	—	6	9	3	0	13	14
Subtotal^e	38	46	17	20	5	7	14	20	15	0	89	94

Table 6a. Stage 3 (AIDS), by transmission category, race/ethnicity, and selected characteristics, 2014—United States (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Asian^f												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	0	0	0	0	0	0	0	0	1	1	1	1
15–19	0	0	0	0	0	0	0	0	0	0	0	0
20–24	13	16	0	0	0	0	0	0	2	0	15	16
25–29	23	28	0	0	2	2	2	2	3	0	30	32
30–34	28	38	1	1	0	1	4	4	8	0	41	45
35–39	27	41	1	2	2	3	2	4	13	0	45	50
40–44	31	43	2	2	5	6	3	5	11	0	52	57
45–49	28	42	1	2	0	0	4	5	13	0	46	50
50–54	7	13	1	2	0	0	3	4	6	0	17	19
55–59	3	5	0	0	0	0	2	3	3	0	8	9
60–64	1	2	0	0	0	0	0	0	1	0	2	2
≥65	2	6	0	1	0	0	0	2	6	0	8	9
Region of residence												
Northeast	32	45	1	3	0	0	2	5	13	0	48	53
Midwest	18	26	0	1	2	2	7	10	10	0	37	39
South	30	53	3	4	0	0	4	7	22	0	59	65
West	83	109	2	3	7	8	7	10	22	1	121	132
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	0	0	0	0	0
15–19	—	—	0	0	—	—	0	0	0	0	0	0
20–24	—	—	0	0	—	—	3	5	2	0	5	5
25–29	—	—	0	0	—	—	4	9	4	0	8	9
30–34	—	—	0	0	—	—	4	5	1	0	5	5
35–39	—	—	0	0	—	—	8	13	4	0	12	13
40–44	—	—	1	1	—	—	2	6	3	0	6	7
45–49	—	—	1	1	—	—	5	7	1	0	7	8
50–54	—	—	0	0	—	—	1	4	3	0	4	4
55–59	—	—	0	0	—	—	2	7	5	0	7	8
60–64	—	—	0	0	—	—	2	2	0	0	2	2
≥65	—	—	0	0	—	—	1	1	0	0	1	1
Region of residence												
Northeast	—	—	0	0	—	—	3	6	3	0	6	7
Midwest	—	—	0	0	—	—	7	7	0	0	7	7
South	—	—	0	0	—	—	10	17	6	0	16	17
West	—	—	2	3	—	—	12	27	14	1	28	31
Subtotal^e	163	234	8	14	9	11	52	89	90	2	322	351

Table 6a. Stage 3 (AIDS), by transmission category, race/ethnicity, and selected characteristics, 2014—United States (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Black/African American												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	0	1	0	0	0	0	1	1	4	3	5	5
15–19	82	103	0	1	2	3	3	5	23	9	110	120
20–24	511	643	1	8	7	13	29	45	105	9	653	718
25–29	734	941	3	19	16	28	56	83	161	3	970	1,075
30–34	456	650	11	30	11	24	93	132	188	2	759	839
35–39	306	458	26	50	7	19	109	159	176	2	624	688
40–44	276	442	24	55	13	25	124	188	208	1	645	710
45–49	255	415	47	91	21	35	147	231	230	1	700	772
50–54	187	309	60	101	23	34	144	230	204	2	618	677
55–59	130	216	64	104	22	30	92	170	164	1	472	521
60–64	62	100	49	74	8	13	46	96	92	0	257	283
≥65	37	64	32	54	5	8	55	110	89	3	218	239
Region of residence												
Northeast	425	593	74	168	19	30	171	280	287	10	976	1,081
Midwest	524	683	24	44	20	30	78	121	170	6	816	885
South	1,826	2,716	196	343	71	139	622	1,003	1,104	17	3,819	4,219
West	261	351	23	31	25	33	28	47	83	2	420	464
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	1	8	8	8	9
15–19	—	—	1	2	—	—	8	20	26	17	35	39
20–24	—	—	2	10	—	—	57	124	86	26	145	160
25–29	—	—	4	22	—	—	135	269	131	7	270	298
30–34	—	—	5	25	—	—	206	379	160	2	371	406
35–39	—	—	10	40	—	—	206	411	196	0	412	451
40–44	—	—	34	74	—	—	204	397	190	0	428	472
45–49	—	—	46	84	—	—	197	375	177	1	420	460
50–54	—	—	55	90	—	—	177	335	154	0	386	426
55–59	—	—	42	67	—	—	115	233	116	0	273	301
60–64	—	—	12	26	—	—	79	145	65	0	156	171
≥65	—	—	9	21	—	—	58	120	61	1	128	141
Region of residence												
Northeast	—	—	57	125	—	—	244	512	287	14	588	652
Midwest	—	—	35	64	—	—	141	313	179	8	355	385
South	—	—	116	249	—	—	1,000	1,855	831	37	1,947	2,141
West	—	—	12	24	—	—	57	129	73	4	142	156
Subtotal^e	3,036	4,343	537	1,050	135	233	2,341	4,260	3,014	98	9,063	9,982

Table 6a. Stage 3 (AIDS), by transmission category, race/ethnicity, and selected characteristics, 2014—United States (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Hispanic/Latino^g												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	0	0	0	0	0	0	0	0	1	1	1	1
15–19	12	13	0	0	0	0	0	0	4	4	16	17
20–24	193	237	2	4	5	8	8	9	37	8	245	267
25–29	365	446	10	15	13	18	30	37	57	1	475	518
30–34	352	464	19	29	20	27	33	41	93	2	517	563
35–39	314	440	32	48	14	21	40	56	119	0	519	565
40–44	242	346	27	41	13	21	44	59	103	1	429	468
45–49	227	319	26	44	13	18	47	64	96	0	409	446
50–54	139	194	42	58	8	11	31	43	63	1	283	308
55–59	71	110	26	40	6	8	21	32	50	0	174	190
60–64	32	58	10	20	5	8	9	21	42	0	98	106
≥65	26	38	6	13	3	4	16	26	24	0	75	81
Region of residence												
Northeast	321	466	77	138	16	24	75	109	184	12	673	749
Midwest	139	194	8	13	7	11	14	21	52	2	220	240
South	799	1,075	61	93	35	52	143	192	285	6	1,323	1,418
West	714	929	54	70	42	57	47	67	168	0	1,025	1,123
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	1	3	2	3	3
15–19	—	—	0	1	—	—	3	6	8	5	11	12
20–24	—	—	2	3	—	—	10	17	17	11	29	32
25–29	—	—	8	13	—	—	21	47	30	5	59	64
30–34	—	—	10	15	—	—	48	75	25	0	83	90
35–39	—	—	10	18	—	—	56	101	41	0	107	119
40–44	—	—	14	21	—	—	60	94	32	1	106	116
45–49	—	—	12	23	—	—	39	80	43	0	94	103
50–54	—	—	20	27	—	—	29	54	25	0	74	81
55–59	—	—	8	13	—	—	27	49	21	0	56	61
60–64	—	—	3	6	—	—	14	27	13	0	30	33
≥65	—	—	0	3	—	—	6	18	14	0	20	22
Region of residence												
Northeast	—	—	40	69	—	—	86	184	109	10	235	262
Midwest	—	—	4	7	—	—	23	34	12	1	39	42
South	—	—	27	45	—	—	152	244	94	7	273	295
West	—	—	16	24	—	—	52	107	57	7	125	138
Subtotal^e	1,973	2,665	287	457	100	144	592	956	961	45	3,913	4,267

Table 6a. Stage 3 (AIDS), by transmission category, race/ethnicity, and selected characteristics, 2014—United States (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Native Hawaiian/Other Pacific Islander												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	0	0	0	0	0	0	0	0	0	0	0	0
15–19	0	0	0	0	0	0	0	0	0	0	0	0
20–24	0	0	0	0	0	0	0	0	0	0	0	0
25–29	1	1	0	0	0	0	0	0	0	0	1	1
30–34	3	3	0	0	1	1	0	0	0	0	4	4
35–39	2	3	0	0	0	0	0	0	1	0	3	3
40–44	1	1	0	0	0	0	1	1	0	0	2	2
45–49	3	3	0	0	0	0	0	0	0	0	3	3
50–54	1	1	0	0	0	0	0	0	0	0	1	1
55–59	0	0	0	0	0	0	0	0	0	0	0	0
60–64	0	1	0	0	0	0	0	0	1	0	1	1
≥65	0	0	0	0	0	0	0	0	0	0	0	0
Region of residence												
Northeast	0	0	0	0	0	0	0	0	0	0	0	0
Midwest	1	1	0	0	0	0	0	0	0	0	1	1
South	1	2	0	0	0	0	1	1	1	0	3	3
West	9	10	0	0	1	1	0	0	1	0	11	12
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	0	0	0	0	0
15–19	—	—	0	0	—	—	0	0	0	0	0	0
20–24	—	—	0	0	—	—	0	0	0	0	0	0
25–29	—	—	0	0	—	—	0	0	0	0	0	0
30–34	—	—	0	0	—	—	0	1	1	0	1	1
35–39	—	—	0	0	—	—	0	0	0	0	0	0
40–44	—	—	0	0	—	—	0	0	0	0	0	0
45–49	—	—	0	0	—	—	0	0	0	0	0	0
50–54	—	—	0	0	—	—	0	0	0	0	0	0
55–59	—	—	0	0	—	—	0	1	1	0	1	1
60–64	—	—	0	0	—	—	1	1	0	0	1	1
≥65	—	—	0	0	—	—	0	0	0	0	0	0
Region of residence												
Northeast	—	—	0	0	—	—	0	0	0	0	0	0
Midwest	—	—	0	0	—	—	0	0	0	0	0	0
South	—	—	0	0	—	—	1	1	0	0	1	1
West	—	—	0	0	—	—	0	2	2	0	2	2
Subtotal^e	11	13	0	0	1	1	2	4	4	0	18	19

Table 6a. Stage 3 (AIDS), by transmission category, race/ethnicity, and selected characteristics, 2014—United States (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
White												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	0	0	0	0	0	0	0	0	0	0	0	0
15–19	7	10	1	1	1	1	0	0	5	3	14	15
20–24	108	128	7	8	15	17	4	4	16	4	150	162
25–29	232	292	11	13	26	31	13	15	44	3	326	355
30–34	304	392	19	22	29	36	14	17	60	1	426	466
35–39	271	363	30	34	29	36	18	22	69	1	417	456
40–44	332	457	40	46	44	53	26	31	99	2	541	590
45–49	474	631	47	56	46	59	36	47	119	1	722	793
50–54	407	563	39	49	47	59	44	55	131	3	668	729
55–59	254	353	29	38	20	25	19	31	85	0	407	447
60–64	147	194	12	15	9	11	18	25	41	2	227	247
≥65	120	181	11	19	3	5	11	21	62	2	207	228
Region of residence												
Northeast	374	509	34	52	28	36	47	62	111	5	594	664
Midwest	479	639	29	35	40	51	28	35	139	6	715	766
South	1,138	1,541	100	118	101	129	87	120	328	11	1,754	1,919
West	665	875	83	94	100	119	41	51	153	1	1,042	1,140
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	0	0	0	0	0
15–19	—	—	0	0	—	—	1	2	2	1	3	3
20–24	—	—	9	12	—	—	11	21	14	5	34	38
25–29	—	—	19	23	—	—	33	47	13	0	65	70
30–34	—	—	9	15	—	—	42	69	25	0	76	84
35–39	—	—	25	33	—	—	47	72	23	0	95	104
40–44	—	—	29	40	—	—	41	79	39	0	109	119
45–49	—	—	29	39	—	—	36	74	39	0	104	113
50–54	—	—	30	40	—	—	42	75	34	0	106	115
55–59	—	—	17	27	—	—	36	65	32	1	85	93
60–64	—	—	10	14	—	—	9	19	12	0	31	34
≥65	—	—	1	4	—	—	17	29	13	2	31	35
Region of residence												
Northeast	—	—	29	48	—	—	43	85	50	4	122	137
Midwest	—	—	17	25	—	—	55	94	39	1	111	120
South	—	—	87	116	—	—	174	290	116	5	377	411
West	—	—	45	59	—	—	43	81	41	1	129	140
Subtotal^e	2,656	3,564	424	547	269	335	518	818	977	33	4,844	5,297

Table 6a. Stage 3 (AIDS), by transmission category, race/ethnicity, and selected characteristics, 2014—United States (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Multiple races												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	0	0	0	0	0	0	0	0	2	2	2	2
15–19	8	9	0	0	0	0	1	1	1	1	10	11
20–24	43	49	1	1	2	2	0	0	2	0	48	53
25–29	59	70	1	1	5	6	2	2	5	0	72	80
30–34	30	43	2	2	6	8	5	6	12	1	55	61
35–39	31	47	5	6	4	6	5	6	13	0	58	65
40–44	37	50	1	3	2	3	9	11	10	0	59	67
45–49	38	51	4	5	5	7	11	15	13	0	71	80
50–54	31	48	5	8	5	7	4	7	17	0	62	69
55–59	18	28	10	13	5	7	9	12	11	0	53	59
60–64	3	7	3	4	2	3	0	2	6	0	14	16
≥65	5	9	1	3	0	1	1	4	9	0	16	18
Region of residence												
Northeast	52	74	13	21	5	8	12	20	28	1	110	124
Midwest	54	70	2	3	6	8	9	11	13	1	84	93
South	151	206	10	14	9	16	26	35	49	2	245	274
West	46	62	8	9	16	19	0	1	11	0	81	91
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	0	0	0	0	0
15–19	—	—	0	0	—	—	4	4	4	4	8	9
20–24	—	—	0	0	—	—	8	10	3	1	11	12
25–29	—	—	3	5	—	—	7	13	7	1	17	19
30–34	—	—	3	4	—	—	9	13	4	1	16	18
35–39	—	—	0	3	—	—	15	29	15	0	30	32
40–44	—	—	3	5	—	—	17	26	8	0	28	31
45–49	—	—	7	11	—	—	8	16	9	0	24	26
50–54	—	—	5	8	—	—	8	15	7	0	20	23
55–59	—	—	3	4	—	—	2	4	2	0	7	8
60–64	—	—	3	3	—	—	8	10	2	0	13	14
≥65	—	—	2	3	—	—	4	6	2	0	8	9
Region of residence												
Northeast	—	—	14	20	—	—	19	31	16	5	49	55
Midwest	—	—	2	4	—	—	12	19	8	1	22	24
South	—	—	9	17	—	—	53	87	36	2	98	106
West	—	—	4	6	—	—	6	9	3	0	13	15
Subtotal^e	303	412	62	93	36	51	137	213	164	13	702	782

Table 6a. Stage 3 (AIDS), by transmission category, race/ethnicity, and selected characteristics, 2014—United States (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Total												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	0	1	0	0	0	0	1	1	8	8	9	10
15–19	109	134	1	2	3	4	4	6	33	18	150	164
20–24	872	1,078	11	21	29	40	41	59	162	22	1,115	1,221
25–29	1,419	1,785	27	50	62	87	103	140	271	8	1,882	2,070
30–34	1,179	1,598	53	86	68	98	151	202	362	5	1,813	1,990
35–39	954	1,356	94	141	56	86	174	246	392	3	1,670	1,832
40–44	926	1,349	98	151	77	109	208	296	434	4	1,743	1,910
45–49	1,031	1,470	125	198	87	121	245	364	473	1	1,961	2,155
50–54	777	1,133	149	220	85	114	226	340	421	6	1,658	1,813
55–59	478	713	130	195	53	70	144	249	313	1	1,118	1,229
60–64	245	361	74	113	24	35	73	144	183	2	599	655
≥65	190	299	50	91	11	18	83	163	191	6	525	576
Region of residence												
Northeast	1,204	1,687	199	382	68	99	308	477	623	28	2,402	2,672
Midwest	1,218	1,617	64	96	77	104	136	197	385	16	1,880	2,031
South	3,953	5,605	374	577	217	338	883	1,359	1,792	36	7,219	7,915
West	1,805	2,369	175	212	193	241	126	179	443	5	2,742	3,006
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	2	11	10	11	12
15–19	—	—	1	4	—	—	16	32	40	27	57	63
20–24	—	—	13	26	—	—	89	178	122	43	224	246
25–29	—	—	34	63	—	—	201	386	186	13	421	461
30–34	—	—	28	61	—	—	309	543	218	4	555	608
35–39	—	—	47	96	—	—	332	627	281	0	660	723
40–44	—	—	83	144	—	—	325	604	273	2	681	749
45–49	—	—	95	159	—	—	286	551	269	2	650	712
50–54	—	—	111	167	—	—	261	487	223	0	595	655
55–59	—	—	70	111	—	—	183	360	177	2	430	473
60–64	—	—	29	51	—	—	115	206	92	1	236	258
≥65	—	—	12	32	—	—	86	174	90	2	188	208
Region of residence												
Northeast	—	—	140	262	—	—	396	819	465	32	1,001	1,114
Midwest	—	—	58	100	—	—	240	470	239	11	537	581
South	—	—	242	430	—	—	1,391	2,497	1,085	50	2,718	2,977
West	—	—	83	121	—	—	176	363	193	12	452	496
Total^e	8,180	11,277	1,335	2,181	555	782	3,656	6,361	5,225	191	18,951	20,792

Abbreviation: IDU, injection drug use.

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

^f Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^g Hispanics/Latinos can be of any race.

Table 6b. Stage 3 (AIDS), by transmission category, race/ethnicity, and selected characteristics, 2014—United States and 6 dependent areas

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
American Indian/Alaska Native												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	0	0	0	0	0	0	0	0	0	0	0	0
15–19	0	0	0	0	0	0	0	0	0	0	0	0
20–24	4	4	0	0	0	0	0	0	0	0	4	4
25–29	5	6	2	2	0	0	0	0	1	0	8	8
30–34	6	7	1	1	1	1	2	2	1	0	11	11
35–39	3	4	0	0	0	0	0	0	1	0	4	4
40–44	7	10	4	4	0	0	1	1	3	0	15	16
45–49	6	8	0	0	2	2	0	1	2	0	10	11
50–54	5	5	2	2	2	2	0	0	0	0	9	10
55–59	2	2	1	1	0	0	1	1	0	0	4	4
60–64	0	0	0	0	0	0	0	0	0	0	0	0
≥65	0	1	0	0	0	0	0	0	1	0	1	1
Region of residence												
Northeast	0	0	0	0	0	0	1	1	0	0	1	1
Midwest	3	4	1	1	2	2	0	0	1	0	7	7
South	8	11	4	5	1	1	0	1	3	0	16	17
West	27	32	5	5	2	3	3	3	5	0	42	44
U.S. dependent areas	0	0	0	0	0	0	0	0	0	0	0	0
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	0	0	0	0	0
15–19	—	—	0	0	—	—	0	0	0	0	0	0
20–24	—	—	0	0	—	—	0	0	0	0	0	0
25–29	—	—	0	0	—	—	1	2	1	0	2	2
30–34	—	—	1	1	—	—	0	2	2	0	3	3
35–39	—	—	2	3	—	—	0	2	2	0	4	4
40–44	—	—	2	3	—	—	1	2	1	0	4	4
45–49	—	—	0	0	—	—	1	1	0	0	1	1
50–54	—	—	1	1	—	—	4	4	0	0	5	5
55–59	—	—	0	0	—	—	1	1	0	0	1	1
60–64	—	—	1	1	—	—	2	2	0	0	3	3
≥65	—	—	0	0	—	—	0	0	0	0	0	0
Region of residence												
Northeast	0	0	0	0	0	0	1	1	0	0	1	1
Midwest	0	0	0	0	0	0	2	3	1	0	3	3
South	0	0	3	4	0	0	1	3	2	0	6	6
West	0	0	4	5	0	0	6	9	3	0	13	14
U.S. dependent areas	0	0	0	0	0	0	0	0	0	0	0	0
Subtotal^e	38	46	17	20	5	7	14	20	15	0	89	94

Table 6b. Stage 3 (AIDS), by transmission category, race/ethnicity, and selected characteristics, 2014—United States and 6 dependent areas (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Asian^f												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	0	0	0	0	0	0	0	0	1	1	1	1
15–19	0	0	0	0	0	0	0	0	0	0	0	0
20–24	13	16	0	0	0	0	0	0	2	0	15	16
25–29	23	28	0	0	2	2	2	2	3	0	30	32
30–34	28	38	1	1	0	1	4	4	8	0	41	45
35–39	27	41	1	2	2	3	2	4	13	0	45	50
40–44	31	43	2	2	5	6	3	5	11	0	52	57
45–49	28	42	1	2	0	0	4	5	13	0	46	50
50–54	8	14	1	2	0	0	3	4	6	0	18	20
55–59	3	5	0	0	0	0	2	3	3	0	8	9
60–64	1	2	0	0	0	0	0	0	1	0	2	2
≥65	2	6	0	1	0	0	0	2	6	0	8	9
Region of residence												
Northeast	32	45	1	3	0	0	2	5	13	0	48	53
Midwest	18	26	0	1	2	2	7	10	10	0	37	39
South	30	53	3	4	0	0	4	7	22	0	59	65
West	83	109	2	3	7	8	7	10	22	1	121	132
U.S. dependent areas	1	1	0	0	0	0	0	0	0	0	1	1
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	0	0	0	0	0
15–19	—	—	0	0	—	—	0	0	0	0	0	0
20–24	—	—	0	0	—	—	3	5	2	0	5	5
25–29	—	—	0	0	—	—	4	9	4	0	8	9
30–34	—	—	0	0	—	—	4	5	1	0	5	5
35–39	—	—	0	0	—	—	8	13	4	0	12	13
40–44	—	—	1	1	—	—	2	6	3	0	6	7
45–49	—	—	1	1	—	—	5	7	1	0	7	8
50–54	—	—	0	0	—	—	1	4	3	0	4	4
55–59	—	—	0	0	—	—	2	7	5	0	7	8
60–64	—	—	0	0	—	—	2	2	0	0	2	2
≥65	—	—	0	0	—	—	1	1	0	0	1	1
Region of residence												
Northeast	—	—	0	0	—	—	3	6	3	0	6	7
Midwest	—	—	0	0	—	—	7	7	0	0	7	7
South	—	—	0	0	—	—	10	17	6	0	16	17
West	—	—	2	3	—	—	12	27	14	1	28	31
U.S. dependent areas	—	—	0	0	—	—	0	0	0	0	0	0
Subtotal^e	164	235	8	14	9	11	52	89	90	2	323	352

Table 6b. Stage 3 (AIDS), by transmission category, race/ethnicity, and selected characteristics, 2014—United States and 6 dependent areas (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Black/African American												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	0	1	0	0	0	0	1	1	4	3	5	5
15–19	82	103	0	1	2	3	3	5	23	9	110	120
20–24	511	643	1	8	7	13	29	45	105	9	653	718
25–29	734	941	3	19	16	28	56	83	161	3	970	1,075
30–34	456	650	11	30	11	24	93	132	188	2	759	839
35–39	306	458	26	50	7	19	109	159	176	2	624	688
40–44	276	442	24	55	13	25	124	188	209	1	646	712
45–49	256	416	47	91	21	35	147	231	230	1	701	773
50–54	187	309	60	102	23	34	146	233	205	2	621	681
55–59	130	216	64	104	22	30	92	170	164	1	472	521
60–64	62	101	49	74	8	13	46	96	93	0	258	284
≥65	37	64	32	54	5	8	55	110	89	3	218	239
Region of residence												
Northeast	425	593	74	168	19	30	171	280	287	10	976	1,081
Midwest	524	683	24	44	20	30	78	121	170	6	816	885
South	1,826	2,716	196	343	71	139	622	1,003	1,104	17	3,819	4,219
West	261	351	23	31	25	33	28	47	83	2	420	464
U.S. dependent areas	1	2	0	1	0	0	2	4	3	0	6	7
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	1	8	8	8	9
15–19	—	—	1	2	—	—	8	20	26	17	35	39
20–24	—	—	2	10	—	—	57	124	86	26	145	160
25–29	—	—	4	22	—	—	135	269	131	7	270	298
30–34	—	—	5	25	—	—	206	379	160	2	371	406
35–39	—	—	10	40	—	—	206	411	196	0	412	451
40–44	—	—	34	74	—	—	204	397	190	0	428	472
45–49	—	—	46	84	—	—	197	375	177	1	420	460
50–54	—	—	55	90	—	—	177	336	155	0	387	427
55–59	—	—	42	67	—	—	115	233	116	0	273	301
60–64	—	—	12	26	—	—	79	145	65	0	156	171
≥65	—	—	9	21	—	—	58	120	61	1	128	141
Region of residence												
Northeast	—	—	57	125	—	—	244	512	287	14	588	652
Midwest	—	—	35	64	—	—	141	313	179	8	355	385
South	—	—	116	249	—	—	1,000	1,855	831	37	1,947	2,141
West	—	—	12	24	—	—	57	129	73	4	142	156
U.S. dependent areas	—	—	0	0	—	—	0	1	1	0	1	1
Subtotal^e	3,037	4,344	537	1,051	135	233	2,343	4,265	3,018	98	9,070	9,991

Table 6b. Stage 3 (AIDS), by transmission category, race/ethnicity, and selected characteristics, 2014—United States and 6 dependent areas (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Hispanic/Latino^g												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	0	0	0	0	0	0	0	0	1	1	1	1
15–19	13	15	0	0	0	0	0	0	4	4	17	19
20–24	205	252	2	4	5	8	8	9	37	8	257	282
25–29	376	462	11	16	13	18	33	41	57	1	490	538
30–34	363	479	21	33	20	27	36	45	94	2	534	586
35–39	322	451	45	65	16	23	51	71	120	0	554	611
40–44	252	359	36	54	15	24	52	70	104	1	459	508
45–49	239	335	39	60	16	22	56	77	97	0	447	494
50–54	145	202	51	70	10	14	34	48	64	1	304	335
55–59	75	115	34	51	6	8	30	44	50	0	195	219
60–64	36	63	11	22	5	8	17	34	44	0	113	128
≥65	29	42	7	14	3	4	23	35	24	0	86	95
Region of residence												
Northeast	321	466	77	138	16	24	75	109	184	12	673	749
Midwest	139	194	8	13	7	11	14	21	52	2	220	240
South	799	1,075	61	93	35	52	143	192	285	6	1,323	1,418
West	714	929	54	70	42	57	47	67	168	0	1,025	1,123
U.S. dependent areas	82	111	57	76	9	12	61	86	7	0	216	285
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	1	3	2	3	3
15–19	—	—	0	1	—	—	4	9	10	6	14	17
20–24	—	—	2	3	—	—	11	19	18	12	31	35
25–29	—	—	9	14	—	—	23	49	30	5	62	68
30–34	—	—	15	21	—	—	57	88	26	0	98	110
35–39	—	—	12	21	—	—	64	112	41	0	117	133
40–44	—	—	18	27	—	—	72	114	35	1	125	142
45–49	—	—	15	28	—	—	44	88	44	0	103	116
50–54	—	—	24	33	—	—	37	66	26	0	87	99
55–59	—	—	12	18	—	—	36	60	21	0	69	78
60–64	—	—	3	6	—	—	16	29	13	0	32	35
≥65	—	—	0	3	—	—	10	23	14	0	24	26
Region of residence												
Northeast	—	—	40	69	—	—	86	184	109	10	235	262
Midwest	—	—	4	7	—	—	23	34	12	1	39	42
South	—	—	27	45	—	—	152	244	94	7	273	295
West	—	—	16	24	—	—	52	107	57	7	125	138
U.S. dependent areas	—	—	23	31	—	—	61	90	9	3	93	124
Subtotal^e	2,055	2,776	367	565	109	156	714	1,132	977	47	4,222	4,676

Table 6b. Stage 3 (AIDS), by transmission category, race/ethnicity, and selected characteristics, 2014—United States and 6 dependent areas (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Native Hawaiian/Other Pacific Islander												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	0	0	0	0	0	0	0	0	0	0	0	0
15–19	0	0	0	0	0	0	0	0	0	0	0	0
20–24	0	0	0	0	0	0	0	0	0	0	0	0
25–29	1	1	0	0	0	0	0	0	0	0	1	1
30–34	3	3	0	0	1	1	0	0	0	0	4	4
35–39	2	3	0	0	0	0	0	0	1	0	3	3
40–44	1	1	0	0	0	0	1	1	0	0	2	2
45–49	3	3	0	0	0	0	0	0	0	0	3	3
50–54	1	1	0	0	0	0	0	0	0	0	1	1
55–59	0	0	0	0	0	0	0	0	0	0	0	0
60–64	0	1	0	0	0	0	0	0	1	0	1	1
≥65	0	0	0	0	0	0	0	0	0	0	0	0
Region of residence												
Northeast	0	0	0	0	0	0	0	0	0	0	0	0
Midwest	1	1	0	0	0	0	0	0	0	0	1	1
South	1	2	0	0	0	0	1	1	1	0	3	3
West	9	10	0	0	1	1	0	0	1	0	11	12
U.S. dependent areas	0	0	0	0	0	0	0	0	0	0	0	0
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	0	0	0	0	0
15–19	—	—	0	0	—	—	0	0	0	0	0	0
20–24	—	—	0	0	—	—	0	0	0	0	0	0
25–29	—	—	0	0	—	—	0	0	0	0	0	0
30–34	—	—	0	0	—	—	0	1	1	0	1	1
35–39	—	—	0	0	—	—	0	0	0	0	0	0
40–44	—	—	0	0	—	—	0	0	0	0	0	0
45–49	—	—	0	0	—	—	0	0	0	0	0	0
50–54	—	—	0	0	—	—	0	0	0	0	0	0
55–59	—	—	0	0	—	—	0	1	1	0	1	1
60–64	—	—	0	0	—	—	1	1	0	0	1	1
≥65	—	—	0	0	—	—	0	0	0	0	0	0
Region of residence												
Northeast	—	—	0	0	—	—	0	0	0	0	0	0
Midwest	—	—	0	0	—	—	0	0	0	0	0	0
South	—	—	0	0	—	—	1	1	0	0	1	1
West	—	—	0	0	—	—	0	2	2	0	2	2
U.S. dependent areas	—	—	0	0	—	—	0	0	0	0	0	0
Subtotal^e	11	13	0	0	1	1	2	4	4	0	18	19

Table 6b. Stage 3 (AIDS), by transmission category, race/ethnicity, and selected characteristics, 2014—United States and 6 dependent areas (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
White												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	0	0	0	0	0	0	0	0	0	0	0	0
15–19	7	10	1	1	1	1	0	0	5	3	14	15
20–24	108	128	7	8	15	17	4	4	16	4	150	162
25–29	232	292	11	13	26	31	13	15	44	3	326	355
30–34	304	392	19	22	29	36	14	17	60	1	426	466
35–39	271	363	30	34	29	36	18	22	69	1	417	456
40–44	332	457	40	46	44	53	26	31	99	2	541	590
45–49	474	631	47	56	46	59	36	47	119	1	722	793
50–54	408	564	39	49	47	59	44	55	131	3	669	731
55–59	254	353	29	38	20	25	19	31	85	0	407	447
60–64	147	194	12	15	9	11	18	25	41	2	227	247
≥65	120	181	11	19	3	5	11	21	62	2	207	228
Region of residence												
Northeast	374	509	34	52	28	36	47	62	111	5	594	664
Midwest	479	639	29	35	40	51	28	35	139	6	715	766
South	1,138	1,541	100	118	101	129	87	120	328	11	1,754	1,919
West	665	875	83	94	100	119	41	51	153	1	1,042	1,140
U.S. dependent areas	1	1	0	0	0	0	0	0	0	0	1	1
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	0	0	0	0	0
15–19	—	—	0	0	—	—	1	2	2	1	3	3
20–24	—	—	9	12	—	—	11	21	14	5	34	38
25–29	—	—	19	23	—	—	33	47	13	0	65	70
30–34	—	—	9	15	—	—	42	69	25	0	76	84
35–39	—	—	25	33	—	—	47	72	23	0	95	104
40–44	—	—	29	40	—	—	41	79	39	0	109	119
45–49	—	—	29	39	—	—	36	74	39	0	104	113
50–54	—	—	30	40	—	—	42	75	34	0	106	115
55–59	—	—	17	27	—	—	36	65	32	1	85	93
60–64	—	—	10	14	—	—	9	19	12	0	31	34
≥65	—	—	1	4	—	—	17	29	13	2	31	35
Region of residence												
Northeast	—	—	29	48	—	—	43	85	50	4	122	137
Midwest	—	—	17	25	—	—	55	94	39	1	111	120
South	—	—	87	116	—	—	174	290	116	5	377	411
West	—	—	45	59	—	—	43	81	41	1	129	140
U.S. dependent areas	—	—	0	0	—	—	0	0	0	0	0	0
Subtotal^e	2,657	3,565	424	547	269	335	518	818	977	33	4,845	5,298

Table 6b. Stage 3 (AIDS), by transmission category, race/ethnicity, and selected characteristics, 2014—United States and 6 dependent areas (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Multiple races												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	0	0	0	0	0	0	0	0	2	2	2	2
15–19	8	9	0	0	0	0	1	1	1	1	10	11
20–24	43	49	1	1	2	2	0	0	2	0	48	53
25–29	59	70	1	1	5	6	2	2	5	0	72	80
30–34	30	43	2	2	6	8	5	6	12	1	55	61
35–39	31	47	5	6	4	6	5	6	13	0	58	65
40–44	37	50	1	3	2	3	9	11	10	0	59	67
45–49	38	51	4	5	5	7	11	15	13	0	71	80
50–54	31	48	5	8	5	7	4	7	17	0	62	69
55–59	19	29	10	13	5	7	9	12	11	0	54	61
60–64	3	7	3	4	2	3	0	2	6	0	14	16
≥65	5	9	1	3	0	1	1	4	9	0	16	18
Region of residence												
Northeast	52	74	13	21	5	8	12	20	28	1	110	124
Midwest	54	70	2	3	6	8	9	11	13	1	84	93
South	151	206	10	14	9	16	26	35	49	2	245	274
West	46	62	8	9	16	19	0	1	11	0	81	91
U.S. dependent areas	1	2	0	0	0	0	0	0	0	0	1	2
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	0	0	0	0	0
15–19	—	—	0	0	—	—	4	4	4	4	8	9
20–24	—	—	0	0	—	—	8	10	3	1	11	12
25–29	—	—	3	5	—	—	7	13	7	1	17	19
30–34	—	—	3	4	—	—	9	13	4	1	16	18
35–39	—	—	0	3	—	—	15	29	15	0	30	32
40–44	—	—	3	5	—	—	17	26	8	0	28	31
45–49	—	—	7	11	—	—	8	16	9	0	24	26
50–54	—	—	5	8	—	—	8	15	7	0	20	23
55–59	—	—	3	4	—	—	2	4	2	0	7	8
60–64	—	—	3	3	—	—	8	10	2	0	13	14
≥65	—	—	2	3	—	—	4	6	2	0	8	9
Region of residence												
Northeast	—	—	14	20	—	—	19	31	16	5	49	55
Midwest	—	—	2	4	—	—	12	19	8	1	22	24
South	—	—	9	17	—	—	53	87	36	2	98	106
West	—	—	4	6	—	—	6	9	3	0	13	15
U.S. dependent areas	—	—	0	0	—	—	0	0	0	0	0	0
Subtotal^e	304	414	62	93	36	51	137	213	164	13	703	783

Table 6b. Stage 3 (AIDS), by transmission category, race/ethnicity, and selected characteristics, 2014—United States and 6 dependent areas (cont)

	Male-to-male sexual contact		IDU		Male-to-male sexual contact and IDU		Heterosexual contact ^a		Other ^b		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Total												
Male adult or adolescent												
Age at diagnosis (yr)												
13–14	0	1	0	0	0	0	1	1	8	8	9	10
15–19	110	136	1	2	3	4	4	6	33	18	151	166
20–24	884	1,093	11	21	29	40	41	59	162	22	1,127	1,236
25–29	1,430	1,801	28	51	62	87	106	144	271	8	1,897	2,090
30–34	1,190	1,613	55	90	68	98	154	207	363	5	1,830	2,013
35–39	962	1,367	107	158	58	88	185	262	393	3	1,705	1,878
40–44	936	1,363	107	164	79	112	216	308	436	4	1,774	1,951
45–49	1,044	1,487	138	214	90	125	254	376	474	1	2,000	2,204
50–54	785	1,143	158	232	87	117	231	347	423	6	1,684	1,846
55–59	483	720	138	206	53	70	153	262	313	1	1,140	1,259
60–64	249	368	75	116	24	35	81	157	186	2	615	678
≥65	193	303	51	92	11	18	90	172	191	6	536	590
Region of residence												
Northeast	1,204	1,687	199	382	68	99	308	477	623	28	2,402	2,672
Midwest	1,218	1,617	64	96	77	104	136	197	385	16	1,880	2,031
South	3,953	5,605	374	577	217	338	883	1,359	1,792	36	7,219	7,915
West	1,805	2,369	175	212	193	241	126	179	443	5	2,742	3,006
U.S. dependent areas	86	117	57	78	9	12	63	90	10	0	225	297
Female adult or adolescent												
Age at diagnosis (yr)												
13–14	—	—	0	0	—	—	0	2	11	10	11	12
15–19	—	—	1	4	—	—	17	35	42	29	60	68
20–24	—	—	13	26	—	—	90	179	123	44	226	249
25–29	—	—	35	64	—	—	203	388	186	13	424	465
30–34	—	—	33	67	—	—	318	557	219	4	570	628
35–39	—	—	49	99	—	—	340	638	281	0	670	737
40–44	—	—	87	149	—	—	337	624	276	2	700	775
45–49	—	—	98	163	—	—	291	559	270	2	659	724
50–54	—	—	115	173	—	—	269	500	225	0	609	673
55–59	—	—	74	116	—	—	192	371	177	2	443	489
60–64	—	—	29	51	—	—	117	209	92	1	238	260
≥65	—	—	12	32	—	—	90	179	90	2	192	213
Region of residence												
Northeast	—	—	140	262	—	—	396	819	465	32	1,001	1,114
Midwest	—	—	58	100	—	—	240	470	239	11	537	581
South	—	—	242	430	—	—	1,391	2,497	1,085	50	2,718	2,977
West	—	—	83	121	—	—	176	363	193	12	452	496
U.S. dependent areas	—	—	23	31	—	—	61	91	10	3	94	125
Total^e	8,266	11,394	1,415	2,290	564	794	3,780	6,542	5,245	193	19,270	21,214

Abbreviation: IDU, injection drug use.

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

^f Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^g Hispanics/Latinos can be of any race.

Table 7a. Diagnoses of HIV infection among children aged <13 years, by race/ethnicity, 2010–2014—United States

Race/ethnicity	2010			2011			2012			2013			2014		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		No.	Rate
American Indian/Alaska Native	0	0	0.0	1	1	0.2	0	0	0.0	0	0	0.0	3	3	0.7
Asian	3	3	0.1	6	6	0.3	13	14	0.6	15	16	0.6	10	11	0.4
Black/African American	147	150	2.1	117	121	1.7	168	175	2.4	120	127	1.8	98	108	1.5
Hispanic/Latino ^b	37	38	0.3	22	23	0.2	15	16	0.1	10	11	0.1	16	18	0.1
Native Hawaiian/Other Pacific Islander	0	0	0.0	0	0	0.0	1	1	1.0	0	0	0.0	1	1	1.0
White	31	32	0.1	34	35	0.1	32	33	0.1	23	24	0.1	26	28	0.1
Multiple races	14	14	0.7	12	13	0.6	11	12	0.5	12	13	0.6	5	5	0.2
Total^c	232	238	0.4	192	198	0.4	240	250	0.5	180	191	0.4	159	174	0.3

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^b Hispanics/Latinos can be of any race.

^c Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 7b. Diagnoses of HIV infection among children aged <13 years, by race/ethnicity, 2010–2014—United States and 6 dependent areas

Race/ethnicity	2010		2011		2012		2013		2014	
	No.	Est. No. ^a	No.	Est. No. ^a	No.	Est. No. ^a	No.	Est. No. ^a	No.	Est. No. ^a
		Est. No. ^a		Est. No. ^a		Est. No. ^a		Est. No. ^a		
American Indian/Alaska Native	0	0	1	1	0	0	0	0	3	3
Asian	3	3	6	6	13	14	15	16	10	11
Black/African American	147	150	117	121	168	175	120	127	98	108
Hispanic/Latino ^b	41	42	24	25	15	16	10	11	17	19
Native Hawaiian/Other Pacific Islander	0	0	0	0	1	1	0	0	1	1
White	31	32	34	35	32	33	23	24	26	28
Multiple races	14	14	12	13	11	12	12	13	5	5
Total^c	236	242	194	200	240	250	180	191	160	176

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting.

^b Hispanics/Latinos can be of any race.

^c Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 8a. Stage 3 (AIDS) among children aged <13 years, by race/ethnicity, 2010–2014 and cumulative—United States

Race/ethnicity	2010			2011			2012			2013			2014			Cumulative ^b	
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Est. No. ^a
American Indian/Alaska Native	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0	1	1	0.2	27	27
Asian ^c	1	1	0.0	0	0	0.0	0	0	0.0	0	0	0.0	1	1	0.0	46	46
Black/African American	13	13	0.2	11	11	0.2	9	9	0.1	4	4	0.1	56	62	0.9	5,674	5,691
Hispanic/Latino ^d	4	4	0.0	2	2	0.0	0	0	0.0	2	2	0.0	11	12	0.1	1,988	1,993
Native Hawaiian/Other Pacific Islander	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0	6	6
White	3	3	0.0	2	2	0.0	1	1	0.0	0	0	0.0	6	6	0.0	1,554	1,556
Multiple races	0	0	0.0	0	0	0.0	0	0	0.0	2	2	0.1	19	21	0.9	266	269
Total^e	21	21	0.0	15	15	0.0	10	10	0.0	8	8	0.0	94	104	0.2	9,561	9,588

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

The criteria for stage 3 (AIDS) classification among pediatric cases were expanded in 2014 (see Technical Notes).

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^b From the beginning of the epidemic through 2014.

^c Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^d Hispanics/Latinos can be of any race.

^e Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 8b. Stage 3 (AIDS) among children aged <13 years, by race/ethnicity, 2010–2014 and cumulative—United States and 6 dependent areas

Race/ethnicity	2010		2011		2012		2013		2014		Cumulative ^a	
	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b
American Indian/Alaska Native	0	0	0	0	0	0	0	0	1	1	27	27
Asian ^c	1	1	0	0	0	0	0	0	1	1	47	47
Black/African American	13	13	11	11	9	9	4	4	56	62	5,686	5,703
Hispanic/Latino ^d	4	4	2	2	0	0	2	2	11	12	2,400	2,406
Native Hawaiian/Other Pacific Islander	0	0	0	0	0	0	0	0	0	0	6	6
White	3	3	2	2	1	1	0	0	6	6	1,554	1,556
Multiple races	0	0	0	0	0	0	2	2	19	21	266	269
Total^e	21	21	15	15	10	10	8	8	94	104	9,986	10,014

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

The criteria for stage 3 (AIDS) classification among pediatric cases were expanded in 2014 (see Technical Notes).

^a From the beginning of the epidemic through 2014.

^b Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting.

^c Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^d Hispanics/Latinos can be of any race.

^e Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 9. Stage 3 (AIDS) among children aged <13 years, by year of diagnosis, 1992–2014—United States and 6 dependent areas

Year of diagnosis	United States		United States and dependent areas	
	No.	Est. No. ^a	No.	Est. No. ^a
1992	908	909	957	958
1993	894	895	932	933
1994	819	820	845	846
1995	675	676	697	698
1996	528	529	541	542
1997	341	342	353	354
1998	250	251	257	258
1999	202	203	206	207
2000	137	137	140	140
2001	132	133	134	135
2002	109	109	116	116
2003	76	76	78	78
2004	59	59	60	60
2005	55	55	55	55
2006	40	41	43	44
2007	35	35	36	36
2008	40	40	41	41
2009	17	17	17	17
2010	21	21	21	21
2011	15	15	15	15
2012	10	10	10	10
2013	8	8	8	8
2014	94	104	94	104

Note. The criteria for stage 3 (AIDS) classification among pediatric cases were expanded in 2014 (see Technical Notes).

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting.

Table 10. Diagnoses of HIV infection among adult and adolescent Hispanics/Latinos, by transmission category and place of birth, 2014—United States and 6 dependent areas

Transmission category	Central America			Cuba			Mexico			Puerto Rico			South America			United States			Total		
	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No. ^b	Est. No. ^{a,b}	%
Male adult or adolescent																					
Male-to-male sexual contact	289	436	78.9	328	369	88.2	896	1,169	85.2	418	531	65.7	395	465	91.4	2,554	3,176	86.6	6,003	7,893	84.2
Injection drug use	6	21	3.9	2	5	1.3	32	53	3.9	85	123	15.2	2	6	1.1	89	152	4.2	252	474	5.1
Male-to-male sexual contact and injection drug use	5	11	1.9	3	4	0.9	13	25	1.8	8	12	1.5	10	12	2.4	111	144	3.9	175	257	2.7
Heterosexual contact ^c	54	82	14.9	33	40	9.6	75	125	9.1	104	141	17.4	18	26	5.1	153	190	5.2	533	744	7.9
Other ^d	145	2	0.4	28	0	0.0	226	0	0.0	95	1	0.2	39	0	0.0	404	5	0.1	1,502	11	0.1
Subtotal	499	553	100	394	418	100	1,242	1,373	100	710	808	100	464	509	100	3,311	3,667	100	8,465	9,379	100
Female adult or adolescent																					
Injection drug use	0	5	3.6	4	5	13.2	4	10	6.1	17	26	11.3	0	2	4.3	59	103	20.3	99	198	13.3
Heterosexual contact ^c	68	140	96.2	26	32	86.8	74	146	93.6	128	200	88.0	20	36	92.7	219	403	79.0	657	1,282	86.1
Other ^d	65	0	0.2	5	0	0.0	63	0	0.2	54	2	0.7	16	1	3.0	183	4	0.8	584	9	0.6
Subtotal	133	146	100	35	37	100	141	156	100	199	228	100	36	39	100	461	510	100	1,340	1,490	100
Total^e	632	698	100	429	455	100	1,383	1,528	100	909	1,035	100	500	548	100	3,772	4,177	100	9,805	10,868	100

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting.

^b Entries include persons whose place of birth is not among those listed and persons whose place of birth is unknown.

^c Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^d Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^e Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 11. Stage 3 (AIDS) among adult and adolescent Hispanics/Latinos, by transmission category and place of birth, 2014—United States and 6 dependent areas

Transmission category	Central America			Cuba			Mexico			Puerto Rico			South America			United States			Total		
	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No. ^b	Est. No. ^{a,b}	%
Male adult or adolescent																					
Male-to-male sexual contact	123	197	71.0	95	108	77.7	381	525	77.7	125	164	40.2	95	112	82.8	869	1,090	75.5	2,055	2,776	72.8
Injection drug use	5	15	5.4	2	4	3.0	24	41	6.0	96	123	30.3	3	4	3.1	101	144	9.9	257	390	10.2
Male-to-male sexual contact and injection drug use	2	5	1.7	3	4	2.7	9	18	2.6	17	22	5.3	4	5	3.6	65	85	5.9	109	156	4.1
Heterosexual contact ^c	42	60	21.5	18	23	16.5	56	93	13.7	68	96	23.6	11	14	10.5	91	111	7.7	340	474	12.4
Other ^d	80	1	0.5	12	0	0.0	153	0	0.0	22	2	0.6	10	0	0.0	201	15	1.0	696	20	0.5
Subtotal	252	277	100	130	139	100	623	676	100	328	407	100	123	135	100	1,327	1,444	100	3,457	3,815	100
Female adult or adolescent																					
Injection drug use	1	6	7.1	1	1	13.3	3	7	7.2	30	41	22.7	3	4	19.3	63	91	28.2	110	175	20.3
Heterosexual contact ^c	39	74	92.8	5	8	86.7	47	85	91.4	85	134	74.6	10	15	70.0	137	215	66.8	374	658	76.5
Other ^d	33	0	0.1	3	0	0.0	36	1	1.4	29	5	2.7	6	2	10.7	93	16	5.0	281	27	3.2
Subtotal	73	80	100	9	10	100	86	93	100	144	180	100	19	21	100	293	322	100	765	861	100
Total^e	325	357	100	139	149	100	709	769	100	472	587	100	142	156	100	1,620	1,766	100	4,222	4,676	100

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting.

^b Entries include persons whose place of birth is not among those listed and persons whose place of birth is unknown.

^c Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^d Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^e Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 12a. Deaths of persons with diagnosed HIV infection, by year of death and selected characteristics, 2010–2013—United States

	2010			2011			2012			2013		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate
Age at death (yr)												
<13	7	8	0.0	0	0	0.0	4	5	0.0	1	1	0.0
13–14	3	3	0.0	1	1	0.0	2	2	0.0	1	1	0.0
15–19	39	41	0.2	25	27	0.1	23	25	0.1	11	14	0.1
20–24	195	205	0.9	198	211	1.0	176	193	0.9	153	182	0.8
25–29	420	443	2.1	413	443	2.1	417	457	2.1	391	468	2.2
30–34	757	795	4.0	688	735	3.6	693	756	3.6	572	667	3.1
35–39	1,192	1,248	6.2	1,051	1,117	5.7	943	1,029	5.3	763	882	4.5
40–44	2,113	2,213	10.6	1,826	1,943	9.2	1,610	1,751	8.3	1,298	1,509	7.2
45–49	3,211	3,361	14.8	2,921	3,111	14.0	2,609	2,833	13.0	2,104	2,422	11.4
50–54	3,181	3,327	14.9	3,144	3,345	14.8	3,105	3,370	14.9	2,708	3,127	13.8
55–59	2,563	2,682	13.6	2,651	2,818	13.9	2,640	2,862	13.8	2,386	2,757	13.0
60–64	1,689	1,767	10.4	1,727	1,837	10.3	1,868	2,032	11.4	1,715	1,985	10.9
≥65	1,690	1,773	4.4	1,825	1,947	4.7	1,973	2,144	5.0	1,950	2,265	5.1
Race/ethnicity												
American Indian/Alaska Native	75	79	3.5	58	61	2.7	54	59	2.6	58	66	2.8
Asian ^b	73	76	0.5	86	91	0.6	72	78	0.5	56	64	0.4
Black/African American	8,257	8,642	22.7	7,928	8,434	22.0	7,682	8,352	21.5	6,492	7,581	19.4
Hispanic/Latino ^c	2,703	2,812	5.5	2,673	2,825	5.4	2,418	2,594	4.9	2,348	2,664	4.9
Native Hawaiian/Other Pacific Islander	7	7	1.4	10	10	2.0	7	7	1.4	4	5	0.9
White	5,135	5,405	2.7	4,972	5,322	2.7	5,039	5,512	2.8	4,327	5,028	2.5
Multiple races	810	846	15.0	743	789	13.5	791	857	14.2	768	874	14.1
Transmission category												
Male adult or adolescent												
Male-to-male sexual contact	5,410	6,881	—	5,287	6,865	—	5,203	6,868	—	4,827	6,660	—
Injection drug use	2,540	3,064	—	2,312	2,871	—	2,192	2,758	—	1,841	2,479	—
Male-to-male sexual contact and injection drug use	1,203	1,376	—	1,170	1,368	—	1,148	1,370	—	1,037	1,296	—
Heterosexual contact ^d	1,284	1,789	—	1,261	1,787	—	1,205	1,740	—	1,078	1,618	—
Perinatal	29	30	—	36	38	—	41	44	—	36	42	—
Other ^e	2,162	89	—	2,148	93	—	2,043	87	—	1,678	75	—
Subtotal	12,628	13,230	10.6	12,214	13,023	10.3	11,832	12,867	10.1	10,497	12,170	9.4

Table 12a. Deaths of persons with diagnosed HIV infection, by year of death and selected characteristics, 2010–2013—United States (cont)

	2010			2011			2012			2013		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate
Female adult or adolescent												
Injection drug use	1,329	1,680	—	1,327	1,669	—	1,229	1,612	—	1,088	1,504	—
Heterosexual contact ^d	1,970	2,887	—	1,846	2,758	—	1,879	2,889	—	1,593	2,544	—
Perinatal	31	32	—	45	48	—	44	47	—	27	31	—
Other ^e	1,095	30	—	1,038	35	—	1,075	40	—	847	31	—
Subtotal	4,425	4,629	3.5	4,256	4,510	3.4	4,227	4,588	3.4	3,555	4,110	3.0
Child (<13 yrs at death)												
Perinatal	6	7	—	0	0	—	3	3	—	1	1	—
Other ^e	1	1	—	0	0	—	1	1	—	0	0	—
Subtotal	7	8	0.0	0	0	0.0	4	5	0.0	1	1	0.0
Region of residence												
Northeast	4,596	4,786	8.6	4,367	4,610	8.3	4,019	4,313	7.7	3,798	4,302	7.7
Midwest	1,928	2,034	3.0	1,860	1,999	3.0	1,844	2,024	3.0	1,581	1,888	2.8
South	8,035	8,426	7.3	7,753	8,272	7.1	7,710	8,437	7.2	6,461	7,620	6.4
West	2,501	2,621	3.6	2,490	2,653	3.6	2,490	2,685	3.6	2,213	2,472	3.3
Total^f	17,060	17,866	5.8	16,470	17,534	5.6	16,063	17,459	5.6	14,053	16,281	5.1

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Deaths of persons with diagnosed HIV infection may be due to any cause.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 12b. Deaths of persons with diagnosed HIV infection, by year of death and selected characteristics, 2010–2013—United States and 6 dependent areas

	2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Age at death (yr)												
<13	7	8	0.0	0	0	0.0	4	5	0.0	1	1	0.0
13–14	3	3	0.0	1	1	0.0	2	2	0.0	1	1	0.0
15–19	41	43	0.2	26	28	0.1	25	27	0.1	11	14	0.1
20–24	199	209	1.0	203	217	1.0	178	195	0.9	157	187	0.8
25–29	433	457	2.1	427	458	2.1	428	470	2.2	400	480	2.2
30–34	799	840	4.1	711	760	3.7	722	789	3.7	594	694	3.2
35–39	1,249	1,308	6.4	1,097	1,167	5.9	988	1,079	5.5	789	914	4.6
40–44	2,181	2,285	10.8	1,900	2,022	9.5	1,659	1,804	8.5	1,334	1,553	7.3
45–49	3,325	3,481	15.2	3,024	3,221	14.3	2,705	2,940	13.4	2,188	2,522	11.7
50–54	3,268	3,419	15.1	3,251	3,460	15.1	3,179	3,451	15.1	2,788	3,221	14.1
55–59	2,627	2,750	13.7	2,721	2,894	14.1	2,706	2,936	14.0	2,445	2,827	13.2
60–64	1,737	1,818	10.6	1,775	1,889	10.5	1,910	2,079	11.5	1,761	2,038	11.1
≥65	1,749	1,835	4.5	1,886	2,012	4.8	2,037	2,215	5.1	2,008	2,333	5.1
Race/ethnicity												
American Indian/Alaska Native	75	79	—	58	61	—	54	59	—	58	66	—
Asian ^b	73	76	—	88	93	—	73	79	—	56	64	—
Black/African American	8,263	8,648	—	7,935	8,442	—	7,691	8,362	—	6,497	7,587	—
Hispanic/Latino ^c	3,251	3,391	—	3,213	3,409	—	2,883	3,110	—	2,763	3,155	—
Native Hawaiian/Other Pacific Islander	7	7	—	11	11	—	7	7	—	6	7	—
White	5,138	5,408	—	4,973	5,324	—	5,041	5,514	—	4,327	5,028	—
Multiple races	811	847	—	744	790	—	794	860	—	770	876	—
Transmission category												
Male adult or adolescent												
Male-to-male sexual contact	5,477	6,956	—	5,370	6,958	—	5,260	6,936	—	4,895	6,741	—
Injection drug use	2,742	3,286	—	2,509	3,094	—	2,356	2,948	—	1,960	2,626	—
Male-to-male sexual contact and injection drug use	1,243	1,420	—	1,200	1,401	—	1,182	1,408	—	1,060	1,323	—
Heterosexual contact ^d	1,366	1,881	—	1,336	1,873	—	1,286	1,834	—	1,153	1,711	—
Perinatal	29	30	—	38	40	—	43	46	—	38	44	—
Other ^e	2,182	90	—	2,165	93	—	2,059	88	—	1,687	75	—
Subtotal	13,039	13,664	10.8	12,618	13,459	10.5	12,186	13,260	10.3	10,793	12,519	9.6

Table 12b. Deaths of persons with diagnosed HIV infection, by year of death and selected characteristics, 2010–2013—United States and 6 dependent areas (cont)

	2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Female adult or adolescent												
Injection drug use	1,376	1,730	—	1,376	1,723	—	1,265	1,653	—	1,135	1,560	—
Heterosexual contact ^d	2,066	2,990	—	1,936	2,861	—	1,960	2,984	—	1,666	2,638	—
Perinatal	33	34	—	48	51	—	46	49	—	29	33	—
Other ^e	1,097	30	—	1,044	36	—	1,082	41	—	853	31	—
Subtotal	4,572	4,784	3.6	4,404	4,671	3.5	4,353	4,728	3.5	3,683	4,262	3.1
Child (<13 yrs at death)												
Perinatal	6	7	—	0	0	—	3	3	—	1	1	—
Other ^e	1	1	—	0	0	—	1	1	—	0	0	—
Subtotal	7	8	0.0	0	0	0.0	4	5	0.0	1	1	0.0
Region of residence												
Northeast	4,596	4,786	8.6	4,367	4,610	8.3	4,019	4,313	7.7	3,798	4,302	7.7
Midwest	1,928	2,034	3.0	1,860	1,999	3.0	1,844	2,024	3.0	1,581	1,888	2.8
South	8,035	8,426	7.3	7,753	8,272	7.1	7,710	8,437	7.2	6,461	7,620	6.4
West	2,501	2,621	3.6	2,490	2,653	3.6	2,490	2,685	3.6	2,213	2,472	3.3
U.S. dependent areas	558	589	14.3	552	597	14.6	480	533	13.2	424	502	12.6
Total^f	17,618	18,456	5.9	17,022	18,130	5.7	16,543	17,992	5.7	14,477	16,783	5.2

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Deaths of persons with diagnosed HIV infection may be due to any cause.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 13a. Deaths of persons with diagnosed HIV infection ever classified as stage 3 (AIDS), by year of death and selected characteristics, 2010–2013 and cumulative—United States

	2010			2011			2012			2013			Cumulative ^b		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Est. No. ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Age at death (yr)															
<13	0	0	0.0	0	0	0.0	1	1	0.0	1	1	0.0	4,988	4,998	
13–14	2	2	0.0	1	1	0.0	1	1	0.0	0	0	0.0	294	296	
15–19	27	28	0.1	20	21	0.1	16	17	0.1	8	10	0.0	1,308	1,318	
20–24	147	152	0.7	149	155	0.7	133	142	0.6	119	134	0.6	9,897	9,967	
25–29	313	324	1.5	310	325	1.5	319	341	1.6	290	327	1.5	47,146	47,389	
30–34	614	637	3.2	553	581	2.8	551	588	2.8	476	534	2.5	102,204	102,708	
35–39	990	1,025	5.1	888	931	4.7	774	828	4.2	645	719	3.7	128,451	129,198	
40–44	1,786	1,852	8.9	1,524	1,598	7.6	1,338	1,428	6.8	1,095	1,227	5.9	122,261	123,245	
45–49	2,688	2,784	12.3	2,465	2,590	11.7	2,221	2,373	10.9	1,790	1,986	9.3	95,240	96,371	
50–54	2,611	2,697	12.1	2,614	2,739	12.1	2,573	2,739	12.1	2,291	2,551	11.3	65,453	66,507	
55–59	2,056	2,122	10.7	2,163	2,261	11.2	2,180	2,316	11.1	1,987	2,204	10.4	41,007	41,777	
60–64	1,371	1,416	8.3	1,395	1,457	8.2	1,498	1,594	8.9	1,413	1,566	8.6	24,182	24,673	
≥65	1,316	1,360	3.4	1,401	1,462	3.5	1,527	1,616	3.7	1,544	1,703	3.8	24,597	25,091	
Race/ethnicity															
American Indian/Alaska Native	58	60	2.6	48	50	2.2	39	41	1.8	48	51	2.2	1,899	1,919	
Asian ^c	58	60	0.4	70	73	0.5	63	67	0.4	52	58	0.4	3,431	3,462	
Black/African American	6,734	6,959	18.3	6,435	6,729	17.5	6,248	6,653	17.2	5,338	5,987	15.3	274,025	277,059	
Hispanic/Latino ^d	2,309	2,377	4.7	2,268	2,362	4.5	2,079	2,191	4.1	2,029	2,220	4.1	102,923	103,819	
Native Hawaiian/Other Pacific Islander	7	7	1.4	9	9	1.8	6	6	1.2	3	3	0.6	359	361	
White	4,063	4,220	2.1	3,999	4,211	2.1	4,024	4,309	2.2	3,513	3,897	2.0	271,923	274,206	
Multiple races	692	715	12.7	654	687	11.8	673	717	11.9	676	746	12.0	12,468	12,710	
Transmission category															
Male adult or adolescent															
Male-to-male sexual contact	4,481	5,512	—	4,451	5,508	—	4,350	5,480	—	4,038	5,237	—	292,934	317,564	
Injection drug use	2,096	2,463	—	1,911	2,296	—	1,847	2,242	—	1,574	2,001	—	119,888	132,455	
Male-to-male sexual contact and injection drug use	1,050	1,182	—	1,007	1,151	—	990	1,154	—	894	1,072	—	48,032	51,413	
Heterosexual contact ^e	1,079	1,451	—	1,037	1,408	—	1,000	1,388	—	897	1,264	—	28,571	36,648	
Perinatal	26	27	—	33	35	—	38	40	—	33	37	—	407	419	
Other ^f	1,616	77	—	1,544	78	—	1,507	73	—	1,254	55	—	52,356	8,751	
Subtotal	10,348	10,712	8.6	9,983	10,475	8.3	9,732	10,377	8.1	8,690	9,667	7.5	542,188	547,251	

Table 13a. Deaths of persons with diagnosed HIV infection ever classified as stage 3 (AIDS), by year of death and selected characteristics, 2010–2013 and cumulative—United States (cont)

	2010			2011			2012			2013			Cumulative ^b	
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Est. No. ^a
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		
Female adult or adolescent														
Injection drug use	1,110	1,367	—	1,126	1,373	—	1,029	1,302	—	938	1,233	—	50,505	57,596
Heterosexual contact ^e	1,636	2,266	—	1,563	2,199	—	1,534	2,224	—	1,352	2,005	—	47,301	59,511
Perinatal	29	30	—	43	45	—	44	47	—	27	31	—	511	527
Other ^f	798	24	—	768	28	—	792	32	—	651	26	—	21,535	3,656
Subtotal	3,573	3,687	2.8	3,500	3,647	2.8	3,399	3,605	2.7	2,968	3,295	2.4	119,852	121,289
Child (<13 yrs at death)														
Perinatal	0	0	—	0	0	—	1	1	—	1	1	—	4,525	4,534
Other ^f	0	0	—	0	0	—	0	0	—	0	0	—	463	464
Subtotal	0	0	0.0	0	0	0.0	1	1	0.0	1	1	0.0	4,988	4,998
Region of residence														
Northeast	3,744	3,847	6.9	3,574	3,708	6.7	3,314	3,477	6.2	3,152	3,403	6.1	210,310	211,694
Midwest	1,536	1,601	2.4	1,468	1,553	2.3	1,458	1,567	2.3	1,268	1,456	2.2	67,858	68,733
South	6,556	6,791	5.9	6,344	6,659	5.7	6,266	6,726	5.7	5,355	6,093	5.1	257,293	260,531
West	2,085	2,160	3.0	2,097	2,202	3.0	2,094	2,213	3.0	1,884	2,010	2.7	131,567	132,579
Total^g	13,921	14,399	4.7	13,483	14,122	4.5	13,132	13,984	4.5	11,659	12,963	4.1	667,028	673,538

Note. Deaths of persons with diagnosed HIV infection may be due to any cause.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^b From the beginning of the epidemic through 2013.

^c Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^d Hispanics/Latinos can be of any race.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 13b. Deaths of persons with diagnosed HIV infection ever classified as stage 3 (AIDS), by year of death and selected characteristics, 2010–2013 and cumulative—United States and 6 dependent areas

	2010			2011			2012			2013			Cumulative ^b		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Est. No. ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Age at death (yr)															
<13	0	0	0.0	0	0	0.0	1	1	0.0	1	1	0.0	5,237	5,248	
13–14	2	2	0.0	1	1	0.0	1	1	0.0	0	0	0.0	303	305	
15–19	29	30	0.1	21	22	0.1	18	19	0.1	8	10	0.0	1,374	1,385	
20–24	148	153	0.7	151	158	0.7	135	144	0.6	122	138	0.6	10,393	10,465	
25–29	321	333	1.6	322	338	1.6	326	349	1.6	293	331	1.5	49,263	49,516	
30–34	639	663	3.3	565	593	2.9	569	607	2.9	488	547	2.5	106,183	106,707	
35–39	1,029	1,065	5.2	919	964	4.8	800	856	4.3	662	738	3.7	133,373	134,152	
40–44	1,843	1,911	9.0	1,578	1,655	7.8	1,376	1,468	6.9	1,119	1,254	5.9	126,609	127,634	
45–49	2,791	2,891	12.6	2,558	2,689	12.0	2,292	2,450	11.1	1,850	2,052	9.5	98,449	99,626	
50–54	2,680	2,769	12.2	2,696	2,827	12.4	2,631	2,802	12.3	2,350	2,616	11.4	67,524	68,615	
55–59	2,103	2,171	10.8	2,218	2,320	11.3	2,226	2,366	11.2	2,033	2,254	10.5	42,182	42,976	
60–64	1,411	1,457	8.5	1,430	1,495	8.3	1,528	1,626	9.0	1,452	1,609	8.8	24,965	25,473	
≥65	1,359	1,405	3.4	1,448	1,512	3.6	1,571	1,664	3.8	1,586	1,750	3.9	25,603	26,117	
Race/ethnicity															
American Indian/Alaska Native	58	60	—	48	50	—	39	41	—	48	51	—	1,899	1,919	
Asian ^c	58	60	—	72	75	—	64	68	—	52	58	—	3,466	3,497	
Black/African American	6,738	6,964	—	6,441	6,736	—	6,255	6,660	—	5,341	5,990	—	274,298	277,336	
Hispanic/Latino ^d	2,739	2,825	—	2,682	2,802	—	2,409	2,548	—	2,328	2,550	—	126,953	128,095	
Native Hawaiian/Other Pacific Islander	7	7	—	10	10	—	6	6	—	5	5	—	366	368	
White	4,063	4,220	—	3,999	4,211	—	4,025	4,310	—	3,513	3,897	—	272,001	274,285	
Multiple races	692	715	—	655	688	—	676	720	—	677	747	—	12,475	12,718	
Transmission category															
Male adult or adolescent															
Male-to-male sexual contact	4,535	5,569	—	4,519	5,582	—	4,390	5,525	—	4,095	5,300	—	296,703	321,408	
Injection drug use	2,258	2,634	—	2,059	2,456	—	1,965	2,373	—	1,655	2,090	—	130,221	143,010	
Male-to-male sexual contact and injection drug use	1,083	1,216	—	1,035	1,181	—	1,018	1,184	—	914	1,094	—	49,888	53,298	
Heterosexual contact ^e	1,140	1,516	—	1,095	1,472	—	1,052	1,446	—	955	1,328	—	31,001	39,151	
Perinatal	26	27	—	35	37	—	40	42	—	34	38	—	435	448	
Other ^f	1,622	78	—	1,552	78	—	1,515	74	—	1,254	55	—	52,698	8,877	
Subtotal	10,664	11,040	8.7	10,295	10,806	8.5	9,980	10,645	8.2	8,907	9,906	7.6	560,946	566,193	

Table 13b. Deaths of persons with diagnosed HIV infection ever classified as stage 3 (AIDS), by year of death and selected characteristics, 2010–2013 and cumulative—United States and 6 dependent areas (cont)

	2010			2011			2012			2013			Cumulative ^b	
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Est. No. ^a
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		
Female adult or adolescent														
Injection drug use	1,147	1,406	—	1,166	1,417	—	1,059	1,335	—	970	1,268	—	52,630	59,774
Heterosexual contact ^e	1,715	2,348	—	1,629	2,273	—	1,593	2,290	—	1,403	2,065	—	50,386	62,688
Perinatal	31	32	—	45	48	—	46	49	—	29	33	—	545	562
Other ^f	798	24	—	772	29	—	795	33	—	654	26	—	21,714	3,754
Subtotal	3,691	3,810	2.9	3,612	3,766	2.8	3,493	3,707	2.7	3,056	3,392	2.5	125,275	126,778
Child (<13 yrs at death)														
Perinatal	0	0	—	0	0	—	1	1	—	1	1	—	4,757	4,767
Other ^f	0	0	—	0	0	—	0	0	—	0	0	—	480	481
Subtotal	0	0	0.0	0	0	0.0	1	1	0.0	1	1	0.0	5,237	5,248
Region of residence														
Northeast	3,744	3,847	6.9	3,574	3,708	6.7	3,314	3,477	6.2	3,152	3,403	6.1	210,310	211,694
Midwest	1,536	1,601	2.4	1,468	1,553	2.3	1,458	1,567	2.3	1,268	1,456	2.2	67,858	68,733
South	6,556	6,791	5.9	6,344	6,659	5.7	6,266	6,726	5.7	5,355	6,093	5.1	257,293	260,531
West	2,085	2,160	3.0	2,097	2,202	3.0	2,094	2,213	3.0	1,884	2,010	2.7	131,567	132,579
U.S. dependent areas	434	452	11.0	424	451	11.0	342	369	9.2	305	336	8.4	24,430	24,681
Total^g	14,355	14,851	4.7	13,907	14,573	4.6	13,474	14,353	4.5	11,964	13,299	4.1	691,458	698,219

Note. Deaths of persons with diagnosed HIV infection may be due to any cause.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited. Rates are not calculated by transmission category because of the lack of denominator data.

^b From the beginning of the epidemic through 2013.

^c Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^d Hispanics/Latinos can be of any race.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 14a. Deaths of persons with diagnosed HIV infection, by race/ethnicity and selected characteristics, 2013—United States

	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c					
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^d	Rate
Age at death (yr)																								
<13	0	0	0.0	0	0	0.0	1	1	0.0	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0	1	1	0.0
13–14	0	0	0.0	0	0	0.0	1	1	0.1	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0	1	1	0.0
15–19	0	0	0.0	0	0	0.0	7	8	0.3	4	5	0.1	0	0	0.0	0	0	0.0	0	0	0.0	11	14	0.1
20–24	0	0	0.0	0	0	0.0	107	127	3.7	24	29	0.6	0	0	0.0	13	16	0.1	9	10	1.9	153	182	0.8
25–29	3	4	2.1	4	5	0.3	227	271	9.5	69	81	1.8	0	0	0.0	62	78	0.6	26	29	7.1	391	468	2.2
30–34	5	6	3.8	4	5	0.3	297	349	12.8	104	118	2.7	1	1	2.3	127	150	1.2	34	39	10.8	572	667	3.1
35–39	5	5	3.7	5	6	0.4	395	458	18.5	143	164	4.0	0	0	0.0	170	199	1.8	45	50	16.9	763	882	4.5
40–44	6	6	4.3	4	5	0.3	570	664	25.4	250	286	7.6	0	0	0.0	389	458	3.6	79	90	33.0	1,298	1,509	7.2
45–49	11	13	8.1	8	9	0.7	911	1,053	39.7	368	415	12.5	1	1	3.2	682	792	5.8	123	139	58.0	2,104	2,422	11.4
50–54	13	16	9.5	8	9	0.9	1,149	1,339	49.2	469	533	18.9	2	3	7.6	900	1,038	6.7	167	190	79.9	2,708	3,127	13.8
55–59	8	9	6.0	8	10	1.0	1,134	1,321	54.3	342	385	17.4	0	0	0.0	780	904	5.9	114	130	63.6	2,386	2,757	13.0
60–64	2	2	1.9	6	7	0.9	805	949	49.4	283	319	19.5	0	0	0.0	532	610	4.5	87	98	63.4	1,715	1,985	10.9
≥65	5	5	2.4	9	10	0.6	888	1,039	27.1	292	330	9.8	0	0	0.0	672	783	2.2	84	97	31.8	1,950	2,265	5.1
Transmission category																								
Male adult or adolescent																								
Male-to-male sexual contact	23	31	—	24	39	—	1,436	2,168	—	748	1,019	—	1	1	—	2,342	3,059	—	253	342	—	4,827	6,660	—
Injection drug use	6	7	—	1	4	—	901	1,278	—	451	590	—	0	0	—	382	473	—	100	127	—	1,841	2,479	—
Male-to-male sexual contact and injection drug use	2	4	—	1	1	—	332	445	—	193	234	—	1	1	—	430	514	—	78	98	—	1,037	1,296	—
Heterosexual contact ^e	4	6	—	4	7	—	714	1,108	—	172	236	—	0	0	—	139	199	—	45	62	—	1,078	1,618	—
Perinatal	0	0	—	0	0	—	27	31	—	6	7	—	0	0	—	1	1	—	2	2	—	36	42	—
Other ^f	7	0	—	16	0	—	907	23	—	275	8	—	0	0	—	394	39	—	79	4	—	1,678	75	—
Subtotal	42	48	5.2	46	52	0.8	4317	5,053	33.6	1,845	2,094	10.1	2	3	1.2	3,688	4,285	5.1	557	635	33.3	10,497	12,170	9.4

Table 14a. Deaths of persons with diagnosed HIV infection, by race/ethnicity and selected characteristics, 2013—United States (cont)

	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c					
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^d	Rate
Female adult or adolescent																								
Injection drug use	3	5	—	0	0	—	570	818	—	188	245	—	0	0	—	263	342	—	64	94	—	1,088	1,504	—
Heterosexual contact ^e	6	12	—	7	11	—	1,029	1,676	—	206	316	—	1	2	—	258	387	—	86	140	—	1,593	2,544	—
Perinatal	0	0	—	0	0	—	18	21	—	4	5	—	0	0	—	2	2	—	3	3	—	27	31	—
Other ^f	7	0	—	3	0	—	557	12	—	105	4	—	1	0	—	116	12	—	58	2	—	847	31	—
Subtotal	16	18	1.8	10	12	0.2	2,174	2,527	15.0	503	570	2.8	2	2	1.0	639	743	0.9	211	239	11.6	3,555	4,110	3.0
Child (<13 yrs at death)																								
Perinatal	0	0	—	0	0	—	1	1	—	0	0	—	0	0	—	0	0	—	0	0	—	1	1	—
Other ^f	0	0	—	0	0	—	0	0	—	0	0	—	0	0	—	0	0	—	0	0	—	0	0	—
Subtotal	0	0	0.0	0	0	0.0	1	1	0.0	0	0	0.0	0	0.0	0.0	0	0	0.0	0	0	0.0	1	1	0.0
Region of residence																								
Northeast	1	1	1.1	11	12	0.4	1,671	1,891	30.1	1,032	1,159	15.3	0	0	0.0	806	929	2.5	277	309	35.8	3,798	4,302	7.7
Midwest	6	7	1.7	6	8	0.4	728	883	12.7	118	148	2.9	0	0	0.0	648	755	1.5	75	88	7.4	1,581	1,888	2.8
South	20	23	3.0	16	19	0.5	3,693	4,360	19.4	687	797	4.0	1	1	1.4	1,757	2,087	3.0	287	332	16.5	6,461	7,620	6.4
West	31	34	3.3	23	25	0.3	400	447	13.2	511	560	2.6	3	4	0.9	1,116	1,257	3.3	129	146	6.8	2,213	2,472	3.3
Total^g	58	66	2.8	56	64	0.4	6,492	7,581	19.4	2,348	2,664	4.9	4	5	0.9	4,327	5,028	2.5	768	874	14.1	14,053	16,281	5.1

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Deaths of persons with a diagnosis of HIV infection may be due to any cause.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 14b. Deaths of persons with diagnosed HIV infection, by race/ethnicity and selected characteristics, 2013—United States and 6 dependent areas

	American Indian/ Alaska Native		Asian ^a		Black/African American		Hispanic/Latino ^b		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Age at death (yr)																
<13	0	0	0	0	1	1	0	0	0	0	0	0	0	0	1	1
13–14	0	0	0	0	1	1	0	0	0	0	0	0	0	0	1	1
15–19	0	0	0	0	7	8	4	5	0	0	0	0	0	0	11	14
20–24	0	0	0	0	107	127	28	33	0	0	13	16	9	10	157	187
25–29	3	4	4	5	227	271	78	93	0	0	62	78	26	29	400	480
30–34	5	6	4	5	297	349	126	144	1	1	127	150	34	39	594	694
35–39	5	5	5	6	395	458	168	194	1	1	170	199	45	50	789	914
40–44	6	6	4	5	571	665	285	328	0	0	389	458	79	90	1,334	1,553
45–49	11	13	8	9	911	1,053	451	514	2	2	682	792	123	139	2,188	2,522
50–54	13	16	8	9	1,151	1,342	546	622	2	3	900	1,038	168	191	2,788	3,221
55–59	8	9	8	10	1,134	1,321	400	453	0	0	780	904	115	131	2,445	2,827
60–64	2	2	6	7	805	949	329	372	0	0	532	610	87	98	1,761	2,038
≥65	5	5	9	10	890	1,042	348	396	0	0	672	783	84	97	2,008	2,333
Transmission category																
Male adult or adolescent																
Male-to-male sexual contact	23	31	24	39	1,437	2,169	813	1,097	2	2	2,342	3,059	254	344	4,895	6,741
Injection drug use	6	7	1	4	901	1,278	570	736	0	0	382	473	100	127	1,960	2,626
Male-to-male sexual contact and injection drug use	2	4	1	1	332	445	216	261	1	1	430	514	78	98	1,060	1,323
Heterosexual contact ^e	4	6	4	7	714	1,109	245	325	1	1	139	199	46	63	1,153	1,711
Perinatal	0	0	0	0	27	31	8	9	0	0	1	1	2	2	38	44
Other ^f	7	0	16	0	908	23	283	8	0	0	394	39	79	4	1,687	75
Subtotal	42	48	46	52	4,319	5,055	2,135	2,437	4	5	3,688	4,285	559	638	10,793	12,519

Table 14b. Deaths of persons with diagnosed HIV infection, by race/ethnicity and selected characteristics, 2013—United States and 6 dependent areas (cont)

	American Indian/ Alaska Native		Asian ^a		Black/African American		Hispanic/Latino ^b		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Female adult or adolescent																
Injection drug use	3	5	0	0	570	819	235	300	0	0	263	342	64	94	1,135	1,560
Heterosexual contact ^e	6	12	7	11	1,029	1,679	279	407	1	2	258	387	86	140	1,666	2,638
Perinatal	0	0	0	0	18	21	6	7	0	0	2	2	3	3	29	33
Other ^f	7	0	3	0	560	12	108	4	1	0	116	12	58	2	853	31
Subtotal	16	18	10	12	2,177	2,531	628	718	2	2	639	743	211	239	3,683	4,262
Child (<13 yrs at end of year)																
Perinatal	0	0	0	0	1	1	0	0	0	0	0	0	0	0	1	1
Other ^f	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Subtotal	0	0	0	0	1	1	0	0	0	0	0	0	0	0	1	1
Region of residence																
Northeast	1	1	11	12	1,671	1,891	1,032	1,159	0	0	806	929	277	309	3,798	4,302
Midwest	6	7	6	8	728	883	118	148	0	0	648	755	75	88	1,581	1,888
South	20	23	16	19	3,693	4,360	687	797	1	1	1,757	2,087	287	332	6,461	7,620
West	31	34	23	25	400	447	511	560	3	4	1,116	1,257	129	146	2,213	2,472
U.S. dependent areas	0	0	0	0	5	6	415	491	2	2	0	0	2	2	424	502
Total^g	58	66	56	64	6,497	7,587	2,763	3,155	6	7	4,327	5,028	770	876	14,477	16,783

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Deaths of persons with a diagnosis of HIV infection may be due to any cause.

Reported numbers less than 12, as well as estimated numbers (and accompanying trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 15a. Deaths of persons with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and selected characteristics, 2013—United States

	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c					
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^d	Rate
Age at death (yr)																								
<13	0	0	0.0	0	0	0.0	1	1	0.0	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0	1	1	0.0
13–14	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0
15–19	0	0	0.0	0	0	0.0	4	4	0.1	4	5	0.1	0	0	0.0	0	0	0.0	0	0	0.0	8	10	0.0
20–24	0	0	0.0	0	0	0.0	85	97	2.8	19	21	0.4	0	0	0.0	8	9	0.1	7	8	1.4	119	134	0.6
25–29	2	2	1.3	3	3	0.2	180	206	7.2	52	57	1.3	0	0	0.0	30	33	0.3	23	26	6.2	290	327	1.5
30–34	4	4	2.8	4	5	0.3	254	289	10.6	94	103	2.4	1	1	2.3	91	100	0.8	28	31	8.6	476	534	2.5
35–39	5	5	3.7	4	5	0.4	343	387	15.6	121	132	3.3	0	0	0.0	128	141	1.2	44	49	16.5	645	719	3.7
40–44	6	6	4.3	4	5	0.3	497	562	21.5	215	238	6.3	0	0	0.0	308	346	2.7	65	71	25.9	1,095	1,227	5.9
45–49	8	8	5.5	9	10	0.8	767	855	32.2	324	353	10.7	1	1	3.2	570	636	4.6	111	123	51.1	1,790	1,986	9.3
50–54	9	10	5.9	8	9	0.9	962	1,081	39.7	412	455	16.1	1	1	3.2	750	830	5.3	149	165	69.5	2,291	2,551	11.3
55–59	7	7	5.1	7	8	0.8	921	1,026	42.2	296	322	14.5	0	0	0.0	655	730	4.8	101	111	54.5	1,987	2,204	10.4
60–64	2	2	1.9	4	4	0.5	637	717	37.4	245	266	16.2	0	0	0.0	447	491	3.6	78	86	55.2	1,413	1,566	8.6
≥65	5	5	2.4	9	10	0.6	687	761	19.8	247	269	8.0	0	0	0.0	526	581	1.6	70	78	25.3	1,544	1,703	3.8
Transmission category																								
Male adult or adolescent																								
Male-to-male sexual contact	19	23	—	23	36	—	1,198	1,679	—	671	869	—	1	1	—	1,904	2,342	—	222	287	—	4,038	5,237	—
Injection drug use	6	7	—	1	4	—	768	1,023	—	386	478	—	0	0	—	323	381	—	90	109	—	1,574	2,001	—
Male-to-male sexual contact and injection drug use	1	2	—	1	1	—	286	366	—	169	197	—	0	0	—	366	419	—	71	86	—	894	1,072	—
Heterosexual contact ^e	3	4	—	4	7	—	593	863	—	146	187	—	0	0	—	109	148	—	42	55	—	897	1,264	—
Perinatal	0	0	—	0	0	—	23	26	—	7	8	—	0	0	—	1	1	—	2	2	—	33	37	—
Other ^f	5	0	—	15	0	—	665	14	—	215	7	—	0	0	—	292	33	—	62	1	—	1,254	55	—
Subtotal	34	36	4.0	44	49	0.8	3,533	3,971	26.4	1,594	1,746	8.4	1	1	0.5	2,995	3,323	4.0	489	541	28.4	8,690	9,667	7.5

Table 15a. Deaths of persons with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and selected characteristics, 2013—United States (cont)

	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c					
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^d	Rate
Female adult or adolescent																								
Injection drug use	3	5	—	0	0	—	484	661	—	171	214	—	0	0	—	222	270	—	58	82	—	938	1,233	—
Heterosexual contact ^e	5	10	—	6	9	—	876	1,325	—	172	252	—	1	2	—	214	290	—	78	118	—	1,352	2,005	—
Perinatal	0	0	—	0	0	—	18	21	—	4	5	—	0	0	—	2	2	—	3	3	—	27	31	—
Other ^f	6	0	—	2	0	—	426	9	—	88	4	—	1	0	—	80	11	—	48	2	—	651	26	—
Subtotal	14	15	1.6	8	9	0.1	1,804	2,015	11.9	435	474	2.3	2	2	1.0	518	574	0.7	187	205	10.0	2,968	3,295	2.4
Child (<13 yrs at death)																								
Perinatal	0	0	—	0	0	—	1	1	—	0	0	—	0	0	—	0	0	—	0	0	—	1	1	—
Other ^f	0	0	—	0	0	—	0	0	—	0	0	—	0	0	—	0	0	—	0	0	—	0	0	—
Subtotal	0	0	0.0	0	0	0.0	1	1	0.0	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0	1	1	0.0
Region of residence																								
Northeast	0	0	0.0	10	11	0.3	1,373	1,478	23.5	878	945	12.5	0	0	0.0	650	712	1.9	241	258	29.9	3,152	3,403	6.1
Midwest	3	3	0.8	5	6	0.3	579	677	9.7	99	122	2.4	0	0	0.0	520	576	1.1	62	71	6.0	1,268	1,456	2.2
South	18	20	2.6	14	17	0.5	3,047	3,472	15.4	591	665	3.4	1	1	1.4	1,427	1,628	2.3	257	291	14.5	5,355	6,093	5.1
West	27	28	2.7	23	24	0.3	339	360	10.6	461	489	2.2	2	2	0.5	916	980	2.6	116	126	5.9	1,884	2,010	2.7
Total^g	48	51	2.2	52	58	0.4	5,338	5,987	15.3	2,029	2,220	4.1	3	3	0.6	3,513	3,897	2.0	676	746	12.0	11,659	12,963	4.1

Note. Deaths of persons with a diagnosis of HIV infection may be due to any cause.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 15b. Deaths of persons with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and selected characteristics, 2013—United States and 6 dependent areas

	American Indian/ Alaska Native		Asian ^a		Black/ African American		Hispanic/Latino ^b		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Age at death (yr)																
<13	0	0	0	0	1	1	0	0	0	0	0	0	0	0	1	1
13–14	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
15–19	0	0	0	0	4	4	4	5	0	0	0	0	0	0	8	10
20–24	0	0	0	0	85	97	22	24	0	0	8	9	7	8	122	138
25–29	2	2	3	3	180	206	55	60	0	0	30	33	23	26	293	331
30–34	4	4	4	5	254	289	106	116	1	1	91	100	28	31	488	547
35–39	5	5	4	5	343	387	137	150	1	1	128	141	44	49	662	738
40–44	6	6	4	5	498	563	238	263	0	0	308	346	65	71	1,119	1,254
45–49	8	8	9	10	767	855	383	419	2	2	570	636	111	123	1,850	2,052
50–54	9	10	8	9	963	1,082	470	518	1	1	750	830	149	165	2,350	2,616
55–59	7	7	7	8	921	1,026	341	371	0	0	655	730	102	112	2,033	2,254
60–64	2	2	4	4	637	717	284	308	0	0	447	491	78	86	1,452	1,609
≥65	5	5	9	10	688	762	288	314	0	0	526	581	70	78	1,586	1,750
Transmission category																
Male adult or adolescent																
Male-to-male sexual contact	19	23	23	36	1,199	1,680	725	928	2	2	1,904	2,342	223	288	4,095	5,300
Injection drug use	6	7	1	4	768	1,023	467	567	0	0	323	381	90	109	1,655	2,090
Male-to-male sexual contact and injection drug use	1	2	1	1	286	366	189	219	0	0	366	419	71	86	914	1,094
Heterosexual contact ^e	3	4	4	7	593	863	203	250	1	1	109	148	42	55	955	1,328
Perinatal	0	0	0	0	23	26	8	9	0	0	1	1	2	2	34	38
Other ^f	5	0	15	0	665	14	215	7	0	0	292	33	62	1	1,254	55
Subtotal	34	36	44	49	3,534	3,972	1,807	1,981	3	3	2,995	3,323	490	542	8,907	9,906

Table 15b. Deaths of persons with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and selected characteristics, 2013—United States and 6 dependent areas (cont)

	American Indian/ Alaska Native		Asian ^a		Black/ African American		Hispanic/Latino ^b		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Female adult or adolescent																
Injection drug use	3	5	0	0	484	661	203	249	0	0	222	270	58	82	970	1,268
Heterosexual contact ^e	5	10	6	9	876	1,327	223	309	1	2	214	290	78	118	1,403	2,065
Perinatal	0	0	0	0	18	21	6	7	0	0	2	2	3	3	29	33
Other ^f	6	0	2	0	428	9	89	4	1	0	80	11	48	2	654	26
Subtotal	14	15	8	9	1,806	2,018	521	569	2	2	518	574	187	205	3,056	3,392
Child (<13 yrs at end of year)																
Perinatal	0	0	0	0	1	1	0	0	0	0	0	0	0	0	1	1
Other ^f	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Subtotal	0	0	0	0	1	1	0	0	0	0	0	0	0	0	1	1
Region of residence																
Northeast	0	0	10	11	1,373	1,478	878	945	0	0	650	712	241	258	3,152	3,403
Midwest	3	3	5	6	579	677	99	122	0	0	520	576	62	71	1,268	1,456
South	18	20	14	17	3,047	3,472	591	665	1	1	1,427	1,628	257	291	5,355	6,093
West	27	28	23	24	339	360	461	489	2	2	916	980	116	126	1,884	2,010
U.S. dependent areas	0	0	0	0	3	4	299	330	2	2	0	0	1	1	305	336
Total^g	48	51	52	58	5,341	5,990	2,328	2,550	5	5	3,513	3,897	677	747	11,964	13,299

Note. Deaths of persons with a diagnosis of HIV infection may be due to any cause.

Reported numbers less than 12, as well as estimated numbers (and accompanying trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 16a. Survival for more than 12, 24, and 36 months after a diagnosis of HIV infection during 2005–2010, by selected characteristics—United States

	No. of persons	Proportion survived (in months)		
		>12	>24	>36
Age at diagnosis (yr)				
<13	1,526	0.98	0.98	0.98
13–14	264	0.99	0.99	0.99
15–19	11,725	0.99	0.99	0.99
20–24	36,203	0.99	0.98	0.98
25–29	38,495	0.98	0.97	0.97
30–34	35,958	0.97	0.96	0.95
35–39	39,229	0.96	0.95	0.94
40–44	40,942	0.94	0.93	0.92
45–49	33,585	0.92	0.90	0.89
50–54	22,257	0.90	0.87	0.85
55–59	12,800	0.87	0.83	0.81
60–64	6,134	0.83	0.79	0.76
≥65	5,041	0.74	0.69	0.64
Race/ethnicity				
American Indian/Alaska Native	982	0.92	0.90	0.88
Asian	3,949	0.96	0.95	0.94
Black/African American	131,958	0.94	0.93	0.91
Hispanic/Latino ^a	57,303	0.95	0.94	0.93
Native Hawaiian/Other Pacific Islander	356	0.97	0.96	0.95
White	79,444	0.95	0.93	0.92
Multiple races	10,167	0.95	0.93	0.92
Transmission category				
Male adult or adolescent				
Male-to-male sexual contact	129,329	0.97	0.96	0.95
Injection drug use	12,499	0.90	0.87	0.85
Male-to-male sexual contact and injection drug use	9,442	0.96	0.95	0.94
Heterosexual contact ^b	20,436	0.94	0.92	0.90
Other ^c	42,600	0.89	0.87	0.85
Subtotal	214,306	0.95	0.93	0.92
Female adult or adolescent				
Injection drug use	7,424	0.94	0.91	0.88
Heterosexual contact ^b	34,765	0.96	0.95	0.94
Other ^c	26,138	0.93	0.91	0.89
Subtotal	68,327	0.95	0.93	0.92
Child (<13 yrs at diagnosis)				
Perinatal	1,249	0.98	0.98	0.98
Other ^d	277	0.99	0.99	0.99
Subtotal	1,526	0.98	0.98	0.98
Region of residence				
Northeast	57,962	0.95	0.93	0.92
Midwest	34,829	0.96	0.94	0.93
South	141,089	0.94	0.93	0.91
West	50,279	0.95	0.94	0.93
Year of diagnosis				
2005	48,353	0.94	0.92	0.90
2006	48,591	0.94	0.92	0.91
2007	49,735	0.95	0.93	0.92
2008	48,218	0.95	0.94	0.92
2009	45,651	0.95	0.94	0.93
2010	43,611	0.96	0.95	0.94
Total	284,159	0.95	0.93	0.92

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data exclude persons whose month of diagnosis or month of death is unknown.

See Technical Notes for method for calculating proportion of persons surviving.

^a Hispanics/Latinos can be of any race.

^b Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^c Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^d Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

Table 16b. Survival for more than 12, 24, and 36 months after a diagnosis of HIV infection during 2005–2010, by selected characteristics—United States and 6 dependent areas

	No. of persons	Proportion survived (in months)		
		>12	>24	>36
Age at diagnosis (yr)				
<13	1,551	0.98	0.98	0.98
13–14	273	0.99	0.99	0.99
15–19	11,871	0.99	0.99	0.99
20–24	36,695	0.99	0.98	0.98
25–29	39,220	0.98	0.97	0.97
30–34	36,753	0.97	0.96	0.95
35–39	40,081	0.96	0.95	0.93
40–44	41,843	0.94	0.93	0.91
45–49	34,354	0.92	0.90	0.88
50–54	22,811	0.89	0.86	0.84
55–59	13,135	0.86	0.83	0.80
60–64	6,333	0.83	0.79	0.75
≥65	5,269	0.74	0.69	0.64
Race/ethnicity				
American Indian/Alaska Native	982	0.92	0.90	0.88
Asian	3,951	0.96	0.95	0.94
Black/African American	132,080	0.94	0.93	0.91
Hispanic/Latino ^a	63,178	0.94	0.93	0.92
Native Hawaiian/Other Pacific Islander	369	0.96	0.95	0.95
White	79,456	0.95	0.93	0.92
Multiple races	10,173	0.95	0.93	0.92
Transmission category				
Male adult or adolescent				
Male-to-male sexual contact	130,766	0.97	0.96	0.95
Injection drug use	13,808	0.89	0.86	0.84
Male-to-male sexual contact and injection drug use	9,653	0.96	0.95	0.93
Heterosexual contact ^b	21,439	0.94	0.92	0.90
Other ^c	42,894	0.89	0.86	0.85
Subtotal	218,560	0.94	0.93	0.92
Female adult or adolescent				
Injection drug use	7,716	0.93	0.91	0.88
Heterosexual contact ^b	36,073	0.96	0.95	0.94
Other ^c	26,289	0.93	0.91	0.89
Subtotal	70,078	0.95	0.93	0.91
Child (<13 yrs at diagnosis)				
Perinatal	1,269	0.98	0.98	0.98
Other ^d	282	0.99	0.99	0.99
Subtotal	1,551	0.98	0.98	0.98
Region of residence				
Northeast	57,962	0.95	0.93	0.92
Midwest	34,829	0.96	0.94	0.93
South	141,089	0.94	0.93	0.91
West	50,279	0.95	0.94	0.93
U.S. dependent areas	6,030	0.87	0.85	0.83
Year of diagnosis				
2005	49,610	0.94	0.92	0.90
2006	49,684	0.94	0.92	0.91
2007	50,730	0.94	0.93	0.92
2008	49,194	0.95	0.93	0.92
2009	46,532	0.95	0.94	0.93
2010	44,439	0.96	0.94	0.93
Total	290,189	0.95	0.93	0.92

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data exclude persons whose month of diagnosis or month of death is unknown.

See Technical Notes for method for calculating proportion of persons surviving.

^a Hispanics/Latinos can be of any race.

^b Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^c Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^d Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

Table 17a. Survival for more than 12, 24, and 36 months after a stage 3 (AIDS) classification during 2005–2010, by selected characteristics—United States

	No. of persons	Proportion survived (in months)		
		>12	>24	>36
Age at diagnosis (yr)				
<13	205	0.93	0.92	0.92
13–14	374	0.99	0.98	0.97
15–19	2,542	0.98	0.97	0.95
20–24	10,600	0.96	0.94	0.93
25–29	18,343	0.95	0.93	0.91
30–34	23,035	0.94	0.92	0.90
35–39	30,113	0.92	0.90	0.88
40–44	34,622	0.91	0.88	0.86
45–49	29,901	0.88	0.85	0.82
50–54	20,033	0.85	0.81	0.78
55–59	11,245	0.81	0.77	0.73
60–64	5,495	0.77	0.71	0.67
≥65	4,468	0.68	0.62	0.57
Race/ethnicity				
American Indian/Alaska Native	670	0.86	0.83	0.79
Asian ^a	2,210	0.92	0.91	0.90
Black/African American	89,689	0.89	0.85	0.83
Hispanic/Latino ^b	39,261	0.91	0.89	0.87
Native Hawaiian/Other Pacific Islander	210	0.94	0.92	0.91
White	50,523	0.90	0.87	0.85
Multiple races	8,413	0.91	0.88	0.86
Transmission category				
Male adult or adolescent				
Male-to-male sexual contact	76,177	0.93	0.90	0.89
Injection drug use	13,350	0.85	0.81	0.77
Male-to-male sexual contact and injection drug use	9,539	0.92	0.88	0.86
Heterosexual contact ^c	16,004	0.89	0.86	0.83
Other ^d	25,425	0.82	0.79	0.77
Subtotal	140,495	0.90	0.87	0.85
Female adult or adolescent				
Injection drug use	8,610	0.88	0.83	0.78
Heterosexual contact ^c	26,479	0.92	0.89	0.86
Other ^d	15,187	0.86	0.83	0.81
Subtotal	50,276	0.89	0.86	0.83
Child (<13 yrs at diagnosis)				
Perinatal	186	0.93	0.92	0.92
Other ^e	19	0.89	0.89	0.89
Subtotal	205	0.93	0.92	0.92
Region of residence				
Northeast	44,309	0.91	0.88	0.86
Midwest	22,324	0.90	0.88	0.85
South	91,061	0.88	0.85	0.82
West	33,282	0.91	0.89	0.87
Year of diagnosis				
2005	35,280	0.89	0.85	0.83
2006	33,495	0.89	0.85	0.83
2007	32,427	0.89	0.86	0.84
2008	31,698	0.90	0.87	0.85
2009	30,479	0.90	0.88	0.86
2010	27,597	0.91	0.88	0.86
Total	190,976	0.90	0.87	0.84

Note. Data exclude persons whose month of diagnosis or month of death is unknown.

See Technical Notes for method for calculating proportion of persons surviving.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^d Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

Table 17b. Survival for more than 12, 24, and 36 months after a stage 3 (AIDS) classification during 2005–2010, by selected characteristics—United States and 6 dependent areas

	No. of persons	Proportion survived (in months)		
		>12	>24	>36
Age at diagnosis (yr)				
<13	210	0.92	0.91	0.91
13–14	381	0.99	0.98	0.97
15–19	2,591	0.98	0.97	0.95
20–24	10,731	0.96	0.94	0.93
25–29	18,662	0.94	0.93	0.91
30–34	23,541	0.93	0.91	0.90
35–39	30,757	0.92	0.89	0.88
40–44	35,442	0.90	0.88	0.85
45–49	30,630	0.88	0.84	0.82
50–54	20,563	0.84	0.80	0.77
55–59	11,540	0.81	0.76	0.73
60–64	5,665	0.76	0.71	0.67
≥65	4,649	0.68	0.61	0.57
Race/ethnicity				
American Indian/Alaska Native	670	0.86	0.83	0.79
Asian ^a	2,211	0.92	0.91	0.90
Black/African American	89,771	0.89	0.85	0.83
Hispanic/Latino ^b	43,545	0.89	0.87	0.85
Native Hawaiian/Other Pacific Islander	220	0.94	0.91	0.90
White	50,531	0.90	0.87	0.85
Multiple races	8,414	0.91	0.88	0.86
Transmission category				
Male adult or adolescent				
Male-to-male sexual contact	77,003	0.92	0.90	0.89
Injection drug use	14,606	0.84	0.79	0.75
Male-to-male sexual contact and injection drug use	9,755	0.91	0.88	0.86
Heterosexual contact ^c	16,676	0.89	0.85	0.83
Other ^d	25,518	0.82	0.79	0.77
Subtotal	143,558	0.89	0.86	0.84
Female adult or adolescent				
Injection drug use	8,925	0.88	0.82	0.78
Heterosexual contact ^c	27,433	0.91	0.88	0.86
Other ^d	15,236	0.86	0.83	0.81
Subtotal	51,594	0.89	0.86	0.83
Child (<13 yrs at diagnosis)				
Perinatal	191	0.92	0.92	0.91
Other ^e	19	0.89	0.89	0.89
Subtotal	210	0.92	0.91	0.91
Region of residence				
Northeast	44,309	0.91	0.88	0.86
Midwest	22,324	0.90	0.88	0.85
South	91,061	0.88	0.85	0.82
West	33,282	0.91	0.89	0.87
U.S. dependent areas	4,386	0.76	0.72	0.69
Year of diagnosis				
2005	36,216	0.88	0.85	0.82
2006	34,300	0.88	0.85	0.83
2007	33,163	0.89	0.86	0.84
2008	32,417	0.89	0.87	0.84
2009	31,127	0.90	0.87	0.85
2010	28,139	0.90	0.88	0.86
Total	195,362	0.89	0.86	0.84

Note. Data exclude persons whose month of diagnosis or month of death is unknown.

See Technical Notes for method for calculating proportion of persons surviving.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^d Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

Table 18a. Persons living with diagnosed HIV infection, by year and selected characteristics, 2010–2013—United States

	2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Age at end of year (yr)												
<13	2,944	2,971	5.6	2,671	2,701	5.1	2,526	2,563	4.9	2,370	2,415	4.6
13–14	1,171	1,177	14.2	1,036	1,042	12.6	872	878	10.6	751	759	9.1
15–19	7,223	7,293	33.2	6,751	6,842	31.6	6,231	6,349	29.7	5,637	5,778	27.3
20–24	28,998	29,400	135.5	30,612	31,167	140.6	31,852	32,622	144.3	31,945	32,980	144.3
25–29	49,217	49,774	235.4	51,438	52,172	245.0	53,785	54,780	256.0	57,276	58,652	271.5
30–34	67,908	68,482	341.2	69,832	70,558	343.7	71,538	72,473	346.3	72,783	73,984	347.4
35–39	89,171	89,707	446.7	85,525	86,152	439.2	84,038	84,798	434.4	84,253	85,193	433.6
40–44	136,072	136,593	653.4	129,908	130,501	619.8	123,946	124,641	591.9	117,410	118,185	565.7
45–49	168,318	168,608	744.8	168,721	169,061	762.2	165,109	165,505	762.1	159,884	160,327	754.3
50–54	135,093	135,105	604.4	144,048	144,034	637.9	153,033	152,984	676.9	162,207	162,034	717.2
55–59	86,897	86,743	438.2	96,115	95,882	473.0	105,582	105,251	506.3	114,839	114,306	538.9
60–64	46,254	46,106	271.4	52,913	52,679	295.6	59,605	59,243	332.3	66,899	66,334	365.8
≥65	34,243	34,017	84.0	39,502	39,159	94.7	46,089	45,568	105.6	53,848	52,994	118.5
Race/ethnicity												
American Indian/Alaska Native	2,514	2,514	110.8	2,615	2,615	114.3	2,748	2,750	119.0	2,868	2,870	123.2
Asian ^b	9,183	9,255	62.6	9,871	9,965	65.3	10,608	10,735	68.2	11,361	11,530	70.9
Black/African American	362,984	364,116	957.9	374,401	375,745	978.7	385,351	386,974	997.9	396,852	398,645	1,018.1
Hispanic/Latino ^c	169,564	170,378	335.7	175,810	176,782	340.3	182,346	183,560	345.8	188,827	190,282	350.8
Native Hawaiian/Other Pacific Islander	712	717	143.5	758	765	149.8	809	818	156.9	859	869	162.8
White	277,765	278,081	140.9	284,169	284,497	144.1	290,389	290,736	147.1	296,976	297,289	150.3
Multiple races	30,787	30,915	547.3	31,448	31,582	541.8	31,955	32,084	533.6	32,359	32,456	523.5
Transmission category												
Male adult or adolescent												
Male-to-male sexual contact	377,287	431,706	—	393,828	451,841	—	410,383	472,561	—	427,150	493,543	—
Injection drug use	67,465	82,781	—	66,233	81,729	—	64,957	80,613	—	63,990	79,709	—
Male-to-male sexual contact and injection drug use	45,693	50,933	—	45,623	50,958	—	45,511	50,929	—	45,398	50,849	—
Heterosexual contact ^d	54,122	67,751	—	55,600	69,847	—	56,895	71,724	—	58,215	73,652	—
Perinatal	3,922	3,929	—	4,100	4,106	—	4,232	4,237	—	4,335	4,336	—
Other ^e	89,985	3,138	—	94,090	3,104	—	98,539	3,086	—	103,032	3,075	—
Subtotal	638,474	640,239	512.0	659,474	661,585	524.0	680,517	683,152	535.7	702,120	705,165	547.4

Table 18a. Persons living with diagnosed HIV infection, by year and selected characteristics, 2010–2013—United States (cont)

	2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Female adult or adolescent												
Injection drug use	39,701	53,226	—	39,046	52,841	—	38,434	52,408	—	37,905	51,976	—
Heterosexual contact ^d	110,583	153,479	—	113,044	158,553	—	115,070	163,103	—	117,259	167,772	—
Perinatal	4,275	4,282	—	4,473	4,479	—	4,626	4,631	—	4,792	4,795	—
Other ^e	57,532	1,778	—	60,364	1,791	—	63,033	1,800	—	65,656	1,817	—
Subtotal	212,091	212,765	162.0	216,927	217,665	164.2	221,163	221,941	166.0	225,612	226,361	167.7
Child (<13 yrs at end of year)												
Perinatal	2,602	2,624	—	2,323	2,347	—	2,144	2,173	—	1,965	1,999	—
Other ^e	342	347	—	348	354	—	382	391	—	405	416	—
Subtotal	2,944	2,971	5.6	2,671	2,701	5.1	2,526	2,563	4.9	2,370	2,415	4.6
Region of residence												
Northeast	224,220	224,961	406.2	227,653	228,438	410.6	231,280	232,164	415.8	234,718	235,612	420.5
Midwest	100,550	100,658	150.3	104,114	104,239	155.2	107,777	107,933	160.3	111,572	111,699	165.3
South	365,470	366,476	319.0	379,033	380,283	327.6	391,792	393,326	335.2	405,462	407,214	343.6
West	163,269	163,879	227.2	168,272	168,990	232.0	173,357	174,234	236.7	178,350	179,415	241.2
Total^f	853,509	855,974	276.7	879,072	881,951	282.9	904,206	907,657	289.0	930,102	933,941	295.1

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 18b. Persons living with diagnosed HIV infection, by year and selected characteristics, 2010–2013—United States and 6 dependent areas

	2010			2011			2012			2013		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate
Age at end of year (yr)												
<13	2,991	3,018	5.6	2,713	2,744	5.1	2,554	2,592	4.8	2,388	2,434	4.6
13–14	1,189	1,195	14.2	1,051	1,058	12.6	894	901	10.7	776	784	9.2
15–19	7,389	7,460	33.5	6,894	6,987	31.8	6,350	6,471	29.8	5,730	5,874	27.3
20–24	29,431	29,840	135.7	31,049	31,614	140.7	32,323	33,108	144.6	32,432	33,486	144.7
25–29	50,030	50,600	236.3	52,244	52,994	245.9	54,577	55,593	256.7	58,045	59,444	272.0
30–34	69,381	69,973	344.0	71,265	72,013	346.3	72,914	73,873	348.6	74,121	75,352	349.6
35–39	91,150	91,703	450.7	87,420	88,065	443.0	85,854	86,632	438.1	86,030	86,990	437.1
40–44	139,017	139,554	659.0	132,658	133,269	625.0	126,529	127,245	596.8	119,796	120,590	570.1
45–49	171,952	172,254	751.7	172,326	172,676	769.0	168,637	169,041	768.8	163,294	163,741	760.9
50–54	138,036	138,051	610.3	147,132	147,120	644.0	156,326	156,278	683.6	165,666	165,490	724.1
55–59	88,779	88,626	442.2	98,209	97,973	477.6	107,842	107,504	511.0	117,357	116,813	544.4
60–64	47,355	47,208	274.1	54,132	53,896	298.5	60,954	60,587	335.5	68,332	67,757	369.0
≥65	35,295	35,068	85.4	40,680	40,333	96.1	47,426	46,896	107.1	55,364	54,492	120.1
Race/ethnicity												
American Indian/Alaska Native	2,515	2,515	—	2,616	2,616	—	2,749	2,751	—	2,869	2,871	—
Asian ^b	9,237	9,309	—	9,923	10,017	—	10,659	10,786	—	11,412	11,581	—
Black/African American	363,335	364,470	—	374,761	376,108	—	385,706	387,332	—	397,215	399,011	—
Hispanic/Latino ^c	187,545	188,446	—	194,000	195,057	—	200,817	202,118	—	207,546	209,073	—
Native Hawaiian/Other Pacific Islander	728	733	—	774	781	—	826	835	—	874	885	—
White	277,831	278,147	—	284,235	284,563	—	290,454	290,801	—	297,044	297,357	—
Multiple races	30,804	30,932	—	31,464	31,598	—	31,969	32,097	—	32,371	32,467	—
Transmission category												
Male adult or adolescent												
Male-to-male sexual contact	380,876	435,436	—	397,591	455,769	—	414,377	476,742	—	431,350	497,957	—
Injection drug use	72,357	87,865	—	71,026	86,711	—	69,669	85,505	—	68,656	84,547	—
Male-to-male sexual contact and injection drug use	46,681	51,940	—	46,597	51,951	—	46,469	51,907	—	46,348	51,816	—
Heterosexual contact ^d	56,555	70,291	—	58,094	72,456	—	59,443	74,391	—	60,804	76,360	—
Perinatal	4,070	4,077	—	4,249	4,255	—	4,383	4,388	—	4,491	4,492	—
Other ^e	90,435	3,188	—	94,572	3,154	—	99,038	3,139	—	103,555	3,128	—
Subtotal	650,974	652,797	515.4	672,129	674,297	527.3	693,379	696,072	539.0	715,204	718,300	550.8

Table 18b. Persons living with diagnosed HIV infection, by year and selected characteristics, 2010–2013—United States and 6 dependent areas (cont)

	2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Female adult or adolescent												
Injection drug use	41,120	54,701	—	40,448	54,298	—	39,821	53,849	—	39,271	53,393	—
Heterosexual contact ^d	114,685	157,812	—	117,221	162,969	—	119,322	167,611	—	121,548	172,336	—
Perinatal	4,402	4,409	—	4,599	4,605	—	4,756	4,760	—	4,922	4,925	—
Other ^e	57,823	1,814	—	60,663	1,828	—	63,348	1,837	—	65,998	1,857	—
Subtotal	218,030	218,736	164.3	222,931	223,700	166.5	227,247	228,057	168.3	231,739	232,511	170.1
Child (<13 yrs at end of year)												
Perinatal	2,642	2,664	—	2,360	2,385	—	2,171	2,200	—	1,983	2,018	—
Other ^e	349	354	—	353	359	—	383	392	—	405	416	—
Subtotal	2,991	3,018	5.6	2,713	2,744	5.1	2,554	2,592	4.8	2,388	2,434	4.6
Region of residence												
Northeast	224,220	224,961	406.2	227,653	228,438	410.6	231,280	232,164	415.8	234,718	235,612	420.5
Midwest	100,550	100,658	150.3	104,114	104,239	155.2	107,777	107,933	160.3	111,572	111,699	165.3
South	365,470	366,476	319.0	379,033	380,283	327.6	391,792	393,326	335.2	405,462	407,214	343.6
West	163,269	163,879	227.2	168,272	168,990	232.0	173,357	174,234	236.7	178,350	179,415	241.2
U.S. dependent areas	18,486	18,577	451.2	18,701	18,790	460.5	18,974	19,064	472.5	19,229	19,304	484.1
Total^f	871,995	874,551	279.0	897,773	900,741	285.2	923,180	926,721	291.3	949,331	953,245	297.4

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 19a. Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by year and selected characteristics, 2010–2013—United States

	2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Age at end of year (yr)												
<13	459	462	0.9	374	377	0.7	314	317	0.6	269	272	0.5
13–14	324	326	3.9	254	256	3.1	193	194	2.4	152	154	1.8
15–19	2,516	2,534	11.5	2,261	2,284	10.5	1,953	1,980	9.3	1,696	1,729	8.2
20–24	7,475	7,568	34.9	7,974	8,112	36.6	8,208	8,387	37.1	8,218	8,443	37.0
25–29	15,752	15,932	75.3	16,100	16,347	76.8	16,439	16,768	78.3	17,296	17,733	82.1
30–34	27,638	27,871	138.9	27,925	28,237	137.6	28,186	28,592	136.6	28,158	28,664	134.6
35–39	44,636	44,892	223.6	41,978	42,298	215.6	40,408	40,806	209.1	39,781	40,278	205.0
40–44	77,472	77,729	371.8	72,872	73,193	347.6	68,566	68,961	327.5	63,972	64,418	308.3
45–49	104,424	104,535	461.8	103,793	103,967	468.7	100,646	100,862	464.4	96,359	96,627	454.6
50–54	87,144	87,102	389.7	92,674	92,637	410.3	97,936	97,898	433.2	103,289	103,194	456.8
55–59	57,449	57,341	289.7	63,445	63,299	312.3	69,556	69,363	333.7	75,413	75,120	354.2
60–64	30,877	30,781	181.2	35,150	35,019	196.5	39,536	39,338	220.7	44,434	44,136	243.4
≥65	23,087	22,950	56.7	26,613	26,420	63.9	30,962	30,684	71.1	36,075	35,633	79.7
Race/ethnicity												
American Indian/Alaska Native	1,356	1,355	59.7	1,411	1,410	61.6	1,466	1,466	63.5	1,509	1,508	64.7
Asian ^b	5,046	5,077	34.4	5,338	5,378	35.3	5,629	5,681	36.1	5,935	6,001	36.9
Black/African American	198,670	199,009	523.5	204,598	205,096	534.2	210,380	211,018	544.2	216,590	217,250	554.8
Hispanic/Latino ^c	102,059	102,485	201.9	105,028	105,537	203.1	107,964	108,568	204.5	110,803	111,480	205.5
Native Hawaiian/Other Pacific Islander	399	402	80.5	421	425	83.2	441	446	85.5	465	470	88.1
White	153,116	152,986	77.5	155,600	155,468	78.7	157,687	157,514	79.7	160,179	159,945	80.9
Multiple races	18,607	18,710	331.3	19,017	19,132	328.2	19,336	19,459	323.6	19,631	19,747	318.5
Transmission category												
Male adult or adolescent												
Male-to-male sexual contact	208,125	232,456	—	214,912	240,811	—	221,388	248,996	—	228,131	257,544	—
Injection drug use	46,705	55,288	—	46,034	54,766	—	45,303	54,172	—	44,686	53,649	—
Male-to-male sexual contact and injection drug use	30,407	33,058	—	30,418	33,150	—	30,357	33,162	—	30,237	33,095	—
Heterosexual contact ^d	33,462	41,047	—	34,537	42,536	—	35,483	43,875	—	36,477	45,341	—
Perinatal	2,198	2,205	—	2,281	2,289	—	2,327	2,336	—	2,386	2,395	—
Other ^e	44,842	2,132	—	46,840	2,106	—	48,977	2,092	—	51,354	2,100	—
Subtotal	365,739	366,187	292.8	375,022	375,658	297.5	383,835	384,634	301.6	393,271	394,124	306.0

Table 19a. Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by year and selected characteristics, 2010–2013—United States (cont)

	2010			2011			2012			2013		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate
Female adult or adolescent												
Injection drug use	25,494	32,217	—	25,232	32,145	—	24,984	32,048	—	24,717	31,905	—
Heterosexual contact ^d	60,000	77,675	—	61,753	80,670	—	63,332	83,485	—	64,879	86,319	—
Perinatal	2,349	2,357	—	2,455	2,464	—	2,528	2,537	—	2,638	2,651	—
Other ^e	25,212	1,128	—	26,577	1,133	—	27,910	1,129	—	29,338	1,130	—
Subtotal	113,055	113,376	86.3	116,017	116,412	87.8	118,754	119,200	89.1	121,572	122,005	90.4
Child (<13 yrs at end of year)												
Perinatal	442	445	—	355	358	—	297	300	—	254	257	—
Other ^e	17	17	—	19	19	—	17	17	—	15	15	—
Subtotal	459	462	0.9	374	377	0.7	314	317	0.6	269	272	0.5
Region of residence												
Northeast	133,906	134,509	242.9	135,485	136,157	244.7	136,936	137,671	246.6	138,144	138,904	247.9
Midwest	52,276	52,242	78.0	54,056	54,031	80.5	55,704	55,679	82.7	57,388	57,311	84.8
South	195,618	195,588	170.3	202,223	202,331	174.3	208,565	208,760	177.9	215,877	216,042	182.3
West	97,453	97,685	135.4	99,649	99,928	137.2	101,698	102,040	138.6	103,703	104,144	140.0
Total^f	479,253	480,024	155.2	491,413	492,447	158.0	502,903	504,150	160.5	515,112	516,401	163.2

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 19b. Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by year and selected characteristics, 2010–2013—United States and 6 dependent areas

	2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Age at end of year (yr)												
<13	467	470	0.9	381	384	0.7	320	323	0.6	272	275	0.5
13–14	331	333	4.0	258	260	3.1	197	199	2.4	158	161	1.9
15–19	2,574	2,592	11.6	2,317	2,341	10.7	1,998	2,025	9.3	1,733	1,767	8.2
20–24	7,625	7,719	35.1	8,109	8,248	36.7	8,335	8,516	37.2	8,347	8,574	37.1
25–29	15,979	16,161	75.5	16,333	16,582	76.9	16,660	16,992	78.5	17,521	17,961	82.2
30–34	28,120	28,358	139.4	28,378	28,695	138.0	28,611	29,023	137.0	28,563	29,077	134.9
35–39	45,568	45,830	225.2	42,830	43,157	217.1	41,210	41,616	210.4	40,523	41,029	206.2
40–44	79,133	79,393	374.9	74,371	74,696	350.3	69,909	70,310	329.8	65,206	65,659	310.4
45–49	106,791	106,901	466.5	106,134	106,305	473.4	102,897	103,109	468.9	98,466	98,732	458.8
50–54	89,148	89,101	393.9	94,759	94,716	414.6	100,146	100,100	437.8	105,631	105,528	461.8
55–59	58,772	58,659	292.7	64,908	64,753	315.6	71,147	70,944	337.2	77,161	76,856	358.2
60–64	31,670	31,571	183.3	36,020	35,883	198.7	40,489	40,283	223.1	45,444	45,135	245.8
≥65	23,815	23,673	57.7	27,434	27,234	64.9	31,908	31,620	72.2	37,147	36,691	80.9
Race/ethnicity												
American Indian/Alaska Native	1,357	1,356	—	1,412	1,411	—	1,467	1,467	—	1,510	1,509	—
Asian ^b	5,070	5,101	—	5,361	5,401	—	5,651	5,702	—	5,957	6,023	—
Black/African American	198,860	199,200	—	204,788	205,287	—	210,573	211,211	—	216,784	217,444	—
Hispanic/Latino ^c	112,528	112,949	—	115,577	116,075	—	118,616	119,203	—	121,592	122,252	—
Native Hawaiian/Other Pacific Islander	409	412	—	431	435	—	452	457	—	474	479	—
White	153,153	153,023	—	155,637	155,505	—	157,726	157,553	—	160,219	159,986	—
Multiple races	18,616	18,720	—	19,026	19,141	—	19,342	19,465	—	19,636	19,752	—
Transmission category												
Male adult or adolescent												
Male-to-male sexual contact	210,161	234,520	—	216,997	242,926	—	223,553	251,193	—	230,368	259,819	—
Injection drug use	49,766	58,383	—	49,047	57,808	—	48,290	57,182	—	47,666	56,652	—
Male-to-male sexual contact and injection drug use	31,102	33,754	—	31,104	33,835	—	31,032	33,835	—	30,903	33,758	—
Heterosexual contact ^d	34,886	42,499	—	36,008	44,034	—	36,975	45,395	—	38,005	46,899	—
Perinatal	2,294	2,301	—	2,378	2,386	—	2,424	2,433	—	2,485	2,494	—
Other ^e	44,968	2,161	—	46,967	2,136	—	49,105	2,122	—	51,491	2,132	—
Subtotal	373,177	373,618	295.0	382,501	383,125	299.6	391,379	392,161	303.7	400,918	401,755	308.1

Table 19b. Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by year and selected characteristics, 2010–2013—United States and 6 dependent areas (cont)

	2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Female adult or adolescent												
Injection drug use	26,403	33,138	—	26,128	33,052	—	25,873	32,947	—	25,600	32,797	—
Heterosexual contact ^d	62,222	79,953	—	64,025	82,998	—	65,650	85,862	—	67,236	88,737	—
Perinatal	2,435	2,442	—	2,543	2,551	—	2,615	2,624	—	2,723	2,736	—
Other ^e	25,289	1,140	—	26,654	1,145	—	27,990	1,141	—	29,423	1,145	—
Subtotal	116,349	116,673	87.6	119,350	119,746	89.1	122,128	122,574	90.5	124,982	125,415	91.7
Child (<13 yrs at end of year)												
Perinatal	448	451	—	360	363	—	302	305	—	257	260	—
Other ^e	19	19	—	21	21	—	18	18	—	15	15	—
Subtotal	467	470	0.9	381	384	0.7	320	323	0.6	272	275	0.5
Region of residence												
Northeast	133,906	134,509	242.9	135,485	136,157	244.7	136,936	137,671	246.6	138,144	138,904	247.9
Midwest	52,276	52,242	78.0	54,056	54,031	80.5	55,704	55,679	82.7	57,388	57,311	84.8
South	195,618	195,588	170.3	202,223	202,331	174.3	208,565	208,760	177.9	215,877	216,042	182.3
West	97,453	97,685	135.4	99,649	99,928	137.2	101,698	102,040	138.6	103,703	104,144	140.0
U.S. dependent areas	10,740	10,736	260.8	10,819	10,808	264.9	10,924	10,907	270.3	11,060	11,043	276.9
Total^f	489,993	490,760	156.6	502,232	503,254	159.4	513,827	515,057	161.9	526,172	527,445	164.6

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 20a. Persons living with diagnosed HIV infection, by race/ethnicity and selected characteristics, year-end 2013—United States

	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c					
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^d	Rate
Age at end of year (yr)																								
<13	8	8	1.8	66	68	2.7	1,547	1,578	21.8	331	336	2.6	3	3	2.9	281	287	1.1	134	137	6.1	2,370	2,415	4.6
13–14	0	0	0.0	11	11	3.0	470	475	40.1	159	160	8.5	0	0	0.0	84	85	1.9	27	27	9.5	751	759	9.1
15–19	9	9	5.0	50	51	5.2	3,638	3,734	121.8	1,107	1,132	24.6	2	2	5.4	589	602	5.2	242	248	38.5	5,637	5,778	27.3
20–24	106	109	54.1	324	337	27.7	19,474	20,110	588.5	5,956	6,147	131.4	30	31	64.3	4,732	4,883	38.3	1,323	1,363	250.0	31,945	32,980	144.3
25–29	177	179	104.5	773	799	59.0	30,615	31,329	1,096.4	12,364	12,686	289.1	66	68	138.6	11,014	11,275	91.1	2,267	2,316	560.3	57,276	58,652	271.5
30–34	299	302	190.4	1,125	1,153	82.5	33,712	34,222	1,251.8	17,725	18,066	412.5	103	105	228.4	16,979	17,252	141.2	2,840	2,884	793.9	72,783	73,984	347.4
35–39	301	302	211.4	1,553	1,584	117.9	36,864	37,234	1,499.6	21,333	21,619	531.3	108	109	280.4	20,965	21,187	188.0	3,129	3,157	1,063.4	84,253	85,193	433.6
40–44	441	443	299.5	2,116	2,148	160.0	48,098	48,386	1,850.2	27,277	27,521	729.1	132	134	369.4	35,233	35,422	278.8	4,113	4,132	1,512.6	117,410	118,185	565.7
45–49	503	502	326.1	1,863	1,883	162.5	62,760	62,920	2,370.2	33,733	33,900	1,023.2	130	131	384.8	55,171	55,261	403.4	5,724	5,730	2,388.1	159,884	160,327	754.3
50–54	469	466	285.0	1,345	1,355	126.4	63,180	63,131	2,321.5	29,647	29,676	1,053.1	106	106	319.2	62,034	61,891	398.1	5,426	5,409	2,277.0	162,207	162,034	717.2
55–59	284	281	192.6	985	989	101.7	47,934	47,678	1,958.9	19,358	19,327	873.8	95	96	347.2	42,529	42,310	278.1	3,654	3,625	1,778.2	114,839	114,306	538.9
60–64	160	159	137.6	605	607	74.1	27,825	27,530	1,434.5	10,562	10,505	642.7	53	53	246.1	25,600	25,406	188.6	2,094	2,073	1,334.2	66,899	66,334	365.8
≥65	111	109	50.0	545	544	31.3	20,735	20,317	529.0	9,275	9,207	274.5	31	31	79.6	21,765	21,430	60.8	1,386	1,354	441.8	53,848	52,994	118.5
Transmission category																								
Male adult or adolescent																								
Male-to-male sexual contact	1,228	1,387	—	6,324	7,764	—	122,737	152,303	—	89,409	104,529	—	543	608	—	191,955	210,299	—	14,954	16,652	—	427,150	493,543	—
Injection drug use	198	223	—	262	397	—	30,731	40,570	—	17,192	21,409	—	23	25	—	13,407	14,610	—	2,177	2,475	—	63,990	79,709	—
Male-to-male sexual contact and injection drug use	273	305	—	290	347	—	13,209	16,000	—	8,697	9,919	—	34	37	—	20,304	21,426	—	2,591	2,816	—	45,398	50,849	—
Heterosexual contact ^e	147	168	—	645	813	—	35,540	47,257	—	11,802	13,953	—	36	41	—	8,270	9,382	—	1,775	2,038	—	58,215	73,652	—
Perinatal	4	4	—	24	24	—	2,554	2,554	—	1,035	1,035	—	1	1	—	543	543	—	174	174	—	4,335	4,336	—
Other ^f	247	12	—	1,730	70	—	53,555	880	—	21,883	437	—	69	3	—	23,035	1,565	—	2,513	109	—	103,032	3,075	—
Subtotal	2,097	2,099	228.4	9,275	9,416	145.5	258,326	259,563	1,727.4	150,018	151,282	728.6	706	715	329.8	257,514	257,826	308.7	24,184	24,263	1,272.6	702,120	705,165	547.4

Table 20a. Persons living with diagnosed HIV infection, by race/ethnicity and selected characteristics, year-end 2013—United States (cont)

	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c					
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^d	Rate
Female adult or adolescent																								
Injection drug use	181	238	—	89	170	—	18,702	27,435	—	6,834	9,257	—	21	26	—	10,193	12,478	—	1,885	2,372	—	37,905	51,976	—
Heterosexual contact ^e	372	504	—	1,146	1,788	—	71,477	106,276	—	20,598	28,065	—	84	121	—	19,506	25,595	—	4,076	5,423	—	117,259	167,772	—
Perinatal	7	7	—	18	18	—	2,938	2,941	—	1,058	1,058	—	2	2	—	575	575	—	194	194	—	4,792	4,795	—
Other ^f	203	14	—	767	70	—	43,862	853	—	9,988	283	—	43	2	—	8,907	528	—	1,886	67	—	65,656	1,817	—
Subtotal	763	763	79.5	2,020	2,047	28.0	136,979	137,504	813.8	38,478	38,664	190.0	150	151	70.9	39,181	39,177	44.9	8,041	8,056	391.4	225,612	226,361	167.7
Child (<13 yrs at end of year)																								
Perinatal	7	7	—	44	45	—	1,275	1,298	—	308	312	—	3	3	—	208	212	—	120	122	—	1,965	1,999	—
Other ^f	1	1	—	22	23	—	272	280	—	23	23	—	0	0	—	73	75	—	14	14	—	405	416	—
Subtotal	8	8	1.8	66	68	2.7	1,547	1,578	21.8	331	336	2.6	3	3	2.9	281	287	1.1	134	137	6.1	2,370	2,415	4.6
Region of residence																								
Northeast	165	166	131.8	2,583	2,623	76.1	95,442	95,740	1,524.1	66,955	67,265	887.5	58	59	290.8	57,917	58,130	154.1	11,598	11,630	1,348.7	234,718	235,612	420.5
Midwest	361	361	90.2	989	1,000	51.0	48,987	49,007	702.7	11,694	11,777	235.4	38	38	132.3	45,297	45,299	87.1	4,206	4,217	355.8	111,572	111,699	165.3
South	708	705	91.5	2,118	2,152	58.3	224,317	225,606	1,001.8	56,968	57,389	289.3	130	132	163.5	108,706	108,666	156.1	12,515	12,565	624.4	405,462	407,214	343.6
West	1,634	1,639	158.4	5,671	5,755	80.4	28,106	28,292	836.9	53,210	53,851	246.8	633	640	158.5	85,056	85,194	221.6	4,040	4,045	189.0	178,350	179,415	241.2
Total^g	2,868	2,870	123.2	11,361	11,530	70.9	396,852	398,645	1,018.1	188,827	190,282	350.8	859	869	162.8	296,976	297,289	150.3	32,359	32,456	523.5	930,102	933,941	295.1

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 20b. Persons living with diagnosed HIV infection, by race/ethnicity and selected characteristics, year-end 2013—United States and 6 dependent areas

	American Indian/ Alaska Native		Asian ^a		Black/ African American		Hispanic/Latino ^b		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Age at end of year (yr)																
<13	8	8	66	68	1,548	1,579	347	352	3	3	281	287	135	138	2,388	2,434
13–14	0	0	11	11	471	476	183	185	0	0	84	85	27	27	776	784
15–19	9	9	50	51	3,641	3,737	1,196	1,223	2	2	589	602	243	249	5,730	5,874
20–24	106	109	324	337	19,479	20,115	6,438	6,649	30	31	4,732	4,883	1,323	1,363	32,432	33,486
25–29	177	179	774	800	30,627	31,342	13,119	13,464	67	69	11,014	11,275	2,267	2,316	58,045	59,444
30–34	299	302	1,129	1,157	33,746	34,257	19,016	19,386	105	107	16,984	17,257	2,842	2,886	74,121	75,352
35–39	301	302	1,560	1,591	36,893	37,264	23,072	23,377	110	111	20,965	21,187	3,129	3,157	86,030	86,990
40–44	441	443	2,126	2,157	48,151	48,440	29,587	29,849	135	137	35,241	35,430	4,115	4,134	119,796	120,590
45–49	503	502	1,874	1,894	62,826	62,987	37,045	37,215	133	134	55,187	55,277	5,726	5,732	163,294	163,741
50–54	470	467	1,352	1,362	63,231	63,182	33,030	33,056	108	108	62,047	61,904	5,428	5,411	165,666	165,490
55–59	284	281	989	993	47,976	47,720	21,820	21,778	95	96	42,538	42,319	3,655	3,626	117,357	116,813
60–64	160	159	608	610	27,850	27,555	11,955	11,888	55	55	25,609	25,415	2,095	2,074	68,332	67,757
≥65	111	109	549	548	20,776	20,358	10,738	10,653	31	31	21,773	21,438	1,386	1,354	55,364	54,492
Transmission category																
Male adult or adolescent																
Male-to-male sexual contact	1,228	1,387	6,345	7,790	122,804	152,383	93,474	108,786	550	615	191,989	210,337	14,960	16,658	431,350	497,957
Injection drug use	198	223	263	405	30,749	40,615	21,835	26,188	23	26	13,411	14,616	2,177	2,475	68,656	84,547
Male-to-male sexual contact and injection drug use	273	305	291	349	13,214	16,006	9,635	10,871	34	37	20,310	21,432	2,591	2,816	46,348	51,816
Heterosexual contact ^e	148	169	647	817	35,585	47,331	14,341	16,580	37	42	8,271	9,383	1,775	2,037	60,804	76,360
Perinatal	4	4	24	24	2,558	2,558	1,186	1,186	1	1	543	543	175	175	4,491	4,492
Other ^f	247	12	1,745	70	53,624	880	22,316	489	70	3	23,040	1,565	2,513	109	103,555	3,128
Subtotal	2,098	2,100	9,315	9,456	258,534	259,773	162,787	164,101	715	724	257,564	257,876	24,191	24,270	715,204	718,300

Table 20b. Persons living with diagnosed HIV infection, by race/ethnicity and selected characteristics, year-end 2013—United States and 6 dependent areas (cont)

	American Indian/ Alaska Native		Asian ^a		Black/ African American		Hispanic/Latino ^b		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Female adult or adolescent																
Injection drug use	181	238	89	173	18,711	27,454	8,188	10,647	21	27	10,196	12,483	1,885	2,372	39,271	53,393
Heterosexual contact ^e	372	504	1,149	1,796	71,553	106,403	24,792	32,473	87	127	19,516	25,608	4,079	5,426	121,548	172,336
Perinatal	7	7	18	18	2,943	2,946	1,183	1,183	2	2	575	575	194	194	4,922	4,925
Other ^f	203	14	775	71	43,926	856	10,249	318	46	2	8,912	528	1,887	68	65,998	1,857
Subtotal	763	763	2,031	2,058	137,133	137,659	44,412	44,620	156	158	39,199	39,195	8,045	8,060	231,739	232,511
Child (<13 yrs at end of year)																
Perinatal	7	7	44	45	1,276	1,299	324	329	3	3	208	212	121	123	1,983	2,018
Other ^f	1	1	22	23	272	280	23	23	0	0	73	75	14	14	405	416
Subtotal	8	8	66	68	1,548	1,579	347	352	3	3	281	287	135	138	2,388	2,434
Region of residence																
Northeast	165	166	2,583	2,623	95,442	95,740	66,955	67,265	58	59	57,917	58,130	11,598	11,630	234,718	235,612
Midwest	361	361	989	1,000	48,987	49,007	11,694	11,777	38	38	45,297	45,299	4,206	4,217	111,572	111,699
South	708	705	2,118	2,152	224,317	225,606	56,968	57,389	130	132	108,706	108,666	12,515	12,565	405,462	407,214
West	1,634	1,639	5,671	5,755	28,106	28,292	53,210	53,851	633	640	85,056	85,194	4,040	4,045	178,350	179,415
U.S. dependent areas	1	1	51	51	363	366	18,719	18,791	15	15	68	68	12	11	19,229	19,304
Total^g	2,869	2,871	11,412	11,581	397,215	399,011	207,546	209,073	874	885	297,044	297,357	32,371	32,467	949,331	953,245

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 21a. Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and selected characteristics, year-end 2013—United States

	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c					
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^d	Rate
Age at end of year (yr)																								
<13	2	2	0.4	4	4	0.2	174	176	2.4	38	38	0.3	2	2	1.9	32	32	0.1	17	17	0.8	269	272	0.5
13–14	0	0	0.0	2	2	0.5	101	103	8.7	29	29	1.5	0	0	0.0	15	15	0.3	5	5	1.8	152	154	1.8
15–19	2	2	1.1	12	12	1.3	1,128	1,151	37.6	333	339	7.4	1	1	2.5	141	143	1.2	79	81	12.6	1,696	1,729	8.2
20–24	22	22	11.1	48	50	4.1	5,315	5,459	159.8	1,578	1,620	34.6	7	7	14.8	838	861	6.8	410	423	77.6	8,218	8,443	37.0
25–29	50	51	29.5	199	205	15.2	9,721	9,963	348.7	3,974	4,074	92.9	17	18	35.9	2,540	2,603	21.0	795	818	198.0	17,296	17,733	82.1
30–34	111	112	70.6	391	401	28.7	13,691	13,923	509.3	7,519	7,663	175.0	42	43	93.0	5,182	5,276	43.2	1,222	1,246	343.1	28,158	28,664	134.6
35–39	132	132	92.6	655	667	49.7	18,035	18,241	734.7	10,817	10,965	269.5	52	53	134.9	8,391	8,495	75.4	1,699	1,725	581.0	39,781	40,278	205.0
40–44	246	247	167.2	1,130	1,146	85.3	26,694	26,857	1,027.0	15,953	16,094	426.4	71	72	198.9	17,322	17,419	137.1	2,556	2,582	945.2	63,972	64,418	308.3
45–49	290	289	188.0	1,115	1,126	97.2	38,218	38,326	1,443.8	21,892	22,004	664.2	88	89	261.3	30,918	30,934	225.8	3,838	3,858	1,607.8	96,359	96,627	454.6
50–54	293	292	178.7	862	867	80.8	40,038	40,017	1,471.5	20,514	20,550	729.3	76	77	230.5	37,636	37,516	241.3	3,870	3,875	1,631.3	103,289	103,194	456.8
55–59	184	183	125.1	684	687	70.6	31,440	31,329	1,287.2	13,694	13,688	618.9	57	57	208.5	26,730	26,559	174.6	2,624	2,618	1,284.2	75,413	75,120	354.2
60–64	99	98	85.3	439	441	53.8	18,271	18,126	944.5	7,669	7,652	468.1	30	30	139.1	16,403	16,272	120.8	1,523	1,517	975.9	44,434	44,136	243.4
≥65	78	77	35.3	394	393	22.6	13,764	13,578	353.5	6,793	6,762	201.6	22	22	56.5	14,031	13,820	39.2	993	981	320.0	36,075	35,633	79.7
Transmission category																								
Male adult or adolescent																								
Male-to-male sexual contact	632	694	—	3,171	3,899	—	63,264	75,577	—	49,031	57,000	—	300	328	—	103,301	110,662	—	8,432	9,384	—	228,131	257,544	—
Injection drug use	119	130	—	160	249	—	21,533	26,925	—	12,394	15,036	—	14	15	—	8,820	9,434	—	1,646	1,859	—	44,686	53,649	—
Male-to-male sexual contact and injection drug use	183	198	—	188	222	—	9,311	10,716	—	6,003	6,737	—	21	23	—	12,694	13,213	—	1,837	1,985	—	30,237	33,095	—
Heterosexual contact ^e	87	99	—	443	555	—	21,793	28,345	—	7,998	9,414	—	22	25	—	4,951	5,541	—	1,183	1,363	—	36,477	45,341	—
Perinatal	2	2	—	10	10	—	1,435	1,439	—	559	562	—	1	1	—	272	273	—	107	108	—	2,386	2,395	—
Other ^f	112	10	—	962	54	—	25,805	515	—	12,517	304	—	31	2	—	10,438	1,140	—	1,489	75	—	51,354	2,100	—
Subtotal	1,135	1,135	123.5	4,934	4,989	77.1	143,141	143,517	955.1	88,502	89,053	428.9	389	394	181.6	140,476	140,262	168.0	14,694	14,775	774.9	393,271	394,124	306.0

Table 21a. Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and selected characteristics, year-end 2013—United States (cont)

	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c					
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^d	Rate
Female adult or adolescent																								
Injection drug use	91	115	—	51	91	—	12,592	17,095	—	4,714	6,041	—	13	15	—	5,910	6,893	—	1,346	1,655	—	24,717	31,905	—
Heterosexual contact ^e	190	244	—	609	865	—	39,487	54,293	—	12,124	15,573	—	43	57	—	9,998	12,140	—	2,428	3,147	—	64,879	86,319	—
Perinatal	4	4	—	6	6	—	1,656	1,664	—	577	580	—	2	2	—	279	280	—	114	115	—	2,638	2,651	—
Other ^f	87	9	—	331	46	—	19,540	504	—	4,848	193	—	16	1	—	3,484	339	—	1,032	38	—	29,338	1,130	—
Subtotal	372	372	38.8	997	1,008	13.8	73,275	73,557	435.3	22,263	22,389	110.0	74	75	34.9	19,671	19,651	22.5	4,920	4,955	240.7	121,572	122,005	90.4
Child (<13 yrs at end of year)																								
Perinatal	1	1	—	4	4	—	163	165	—	38	38	—	2	2	—	29	29	—	17	17	—	254	257	—
Other ^f	1	1	—	0	0	—	11	11	—	0	0	—	0	0	—	3	3	—	0	0	—	15	15	—
Subtotal	2	2	0.4	4	4	0.2	174	176	2.4	38	38	0.3	2	2	1.9	32	32	0.1	17	17	0.8	269	272	0.5
Region of residence																								
Northeast	85	85	67.9	1,326	1,342	38.9	56,575	56,901	905.8	40,903	41,196	543.5	24	24	119.9	31,804	31,852	84.5	7,427	7,504	870.3	138,144	138,904	247.9
Midwest	169	169	42.2	472	474	24.2	25,116	25,046	359.1	6,524	6,532	130.5	17	17	59.4	22,751	22,731	43.7	2,339	2,341	197.5	57,388	57,311	84.8
South	326	322	41.9	961	970	26.3	118,494	118,824	527.6	31,391	31,489	158.7	51	51	63.6	57,256	56,960	81.8	7,398	7,426	369.0	215,877	216,042	182.3
West	929	932	90.1	3,176	3,215	44.9	16,405	16,479	487.5	31,985	32,263	147.9	373	378	93.5	48,368	48,403	125.9	2,467	2,476	115.7	103,703	104,144	140.0
Total^g	1,509	1,508	64.7	5,935	6,001	36.9	216,590	217,250	554.8	110,803	111,480	205.5	465	470	88.1	160,179	159,945	80.9	19,631	19,747	318.5	515,112	516,401	163.2

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 21b. Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and selected characteristics, year-end 2013—United States and 6 dependent areas

	American Indian/ Alaska Native		Asian ^a		Black/ African American		Hispanic/Latino ^b		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Age at end of year																
<13	2	2	4	4	174	176	41	41	2	2	32	32	17	17	272	275
13–14	0	0	2	2	101	103	35	35	0	0	15	15	5	5	158	161
15–19	2	2	12	12	1,131	1,154	367	374	1	1	141	143	79	81	1,733	1,767
20–24	22	22	48	50	5,316	5,460	1,706	1,751	7	7	838	861	410	423	8,347	8,574
25–29	50	51	199	205	9,723	9,965	4,197	4,301	17	18	2,540	2,603	795	818	17,521	17,961
30–34	111	112	392	402	13,701	13,933	7,912	8,064	43	44	5,182	5,276	1,222	1,246	28,563	29,077
35–39	132	132	657	669	18,046	18,252	11,544	11,701	53	54	8,391	8,495	1,700	1,726	40,523	41,029
40–44	246	247	1,132	1,148	26,722	26,886	17,152	17,301	73	74	17,325	17,422	2,556	2,582	65,206	65,659
45–49	290	289	1,125	1,136	38,261	38,370	23,932	24,042	91	92	30,928	30,945	3,839	3,859	98,466	98,732
50–54	294	293	864	869	40,068	40,046	22,813	22,842	77	78	37,642	37,522	3,873	3,878	105,631	105,528
55–59	184	183	685	688	31,464	31,353	15,409	15,391	57	57	26,738	26,567	2,624	2,618	77,161	76,856
60–64	99	98	441	443	18,291	18,147	8,651	8,623	31	31	16,408	16,277	1,523	1,516	45,444	45,135
≥65	78	77	396	395	13,786	13,600	7,833	7,787	22	22	14,039	13,828	993	981	37,147	36,691
Transmission category																
Male adult or adolescent																
Male-to-male sexual contact	632	694	3,182	3,912	63,306	75,624	51,186	59,184	304	332	103,323	110,686	8,435	9,387	230,368	259,819
Injection drug use	119	130	161	252	21,546	26,948	15,358	18,011	14	15	8,822	9,437	1,646	1,859	47,666	56,652
Male-to-male sexual contact and injection drug use	183	198	189	223	9,313	10,718	6,662	7,392	21	23	12,698	13,217	1,837	1,985	30,903	33,758
Heterosexual contact ^e	88	100	444	556	21,826	28,391	9,491	10,923	22	25	4,951	5,541	1,183	1,363	38,005	46,899
Perinatal	2	2	10	10	1,437	1,441	656	659	1	1	272	273	107	108	2,485	2,494
Other ^f	112	10	967	54	25,833	515	12,617	335	32	2	10,441	1,140	1,489	75	51,491	2,132
Subtotal	1,136	1,136	4,953	5,008	143,261	143,637	95,970	96,504	394	399	140,507	140,294	14,697	14,777	400,918	401,755

Table 21b. Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and selected characteristics, year-end 2013—United States and 6 dependent areas (cont)

	American Indian/ Alaska Native		Asian ^a		Black/ African American		Hispanic/Latino ^b		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Female adult or adolescent																
Injection drug use	91	115	51	91	12,596	17,103	5,590	6,920	13	15	5,912	6,896	1,347	1,656	25,600	32,797
Heterosexual contact ^e	190	244	611	868	39,528	54,354	14,430	17,918	45	60	10,003	12,145	2,429	3,148	67,236	88,737
Perinatal	4	4	6	6	1,660	1,668	658	661	2	2	279	280	114	115	2,723	2,736
Other ^f	87	9	332	46	19,565	505	4,903	207	18	1	3,486	339	1,032	38	29,423	1,145
Subtotal	372	372	1,000	1,011	73,349	73,631	25,581	25,706	78	79	19,680	19,660	4,922	4,957	124,982	125,415
Child (<13 yrs at end of year)																
Perinatal	1	1	4	4	163	165	41	41	2	2	29	29	17	17	257	260
Other ^f	1	1	0	0	11	11	0	0	0	0	3	3	0	0	15	15
Subtotal	2	2	4	4	174	176	41	41	2	2	32	32	17	17	272	275
Region of residence																
Northeast	85	85	1,326	1,342	56,575	56,901	40,903	41,196	24	24	31,804	31,852	7,427	7,504	138,144	138,904
Midwest	169	169	472	474	25,116	25,046	6,524	6,532	17	17	22,751	22,731	2,339	2,341	57,388	57,311
South	326	322	961	970	118,494	118,824	31,391	31,489	51	51	57,256	56,960	7,398	7,426	215,877	216,042
West	929	932	3,176	3,215	16,405	16,479	31,985	32,263	373	378	48,368	48,403	2,467	2,476	103,703	104,144
U.S. dependent areas	1	1	22	22	194	194	10,789	10,772	9	9	40	40	5	5	11,060	11,043
Total^g	1,510	1,509	5,957	6,023	216,784	217,444	121,592	122,252	474	479	160,219	159,986	19,636	19,752	526,172	527,445

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 22. Diagnoses of HIV infection, by area of residence, 2014—United States and 6 dependent areas

Area of residence	Adults and adolescents			Children			Total		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate
Alabama	633	699	17.2	1	1	0.1	634	700	14.4
Alaska	39	41	6.8	1	1	0.9	40	42	5.7
Arizona	755	798	14.3	3	3	0.3	758	801	11.9
Arkansas	309	340	13.8	2	2	0.4	311	343	11.6
California	4,874	5,533	17.2	16	18	0.3	4,890	5,551	14.3
Colorado	379	401	9.0	2	2	0.2	381	403	7.5
Connecticut	287	305	10.0	0	0	0.0	287	305	8.5
Delaware	119	132	16.8	0	0	0.0	119	132	14.2
District of Columbia	327	381	66.9	0	0	0.0	327	381	57.8
Florida	5,037	5,332	31.3	14	15	0.5	5,051	5,347	26.9
Georgia	1,686	2,247	27.0	4	6	0.3	1,690	2,253	22.3
Hawaii	104	110	9.2	0	0	0.0	104	110	7.8
Idaho	25	27	2.0	1	1	0.4	26	28	1.7
Illinois	1,510	1,728	16.1	8	9	0.4	1,518	1,737	13.5
Indiana	463	490	9.0	2	2	0.2	465	492	7.5
Iowa	98	104	4.0	2	2	0.5	100	107	3.4
Kansas	130	138	5.8	2	2	0.4	132	140	4.8
Kentucky	344	365	9.9	7	7	1.0	351	372	8.4
Louisiana	1,332	1,408	36.6	3	3	0.4	1,335	1,411	30.4
Maine	57	60	5.2	1	1	0.6	58	61	4.6
Maryland	1,121	1,388	27.7	2	3	0.3	1,123	1,391	23.3
Massachusetts	543	735	12.8	2	3	0.3	545	738	10.9
Michigan	791	837	10.0	5	5	0.3	796	842	8.5
Minnesota	300	318	7.0	2	2	0.2	302	320	5.9
Mississippi	488	519	21.0	0	0	0.0	488	519	17.3
Missouri	473	500	9.9	6	6	0.6	479	506	8.3
Montana	15	16	1.9	0	0	0.0	15	16	1.6
Nebraska	88	93	6.1	0	0	0.0	88	93	5.0
Nevada	435	463	19.6	2	2	0.4	437	465	16.4
New Hampshire	38	41	3.6	0	0	0.0	38	41	3.1
New Jersey	1,316	1,536	20.4	2	2	0.2	1,318	1,538	17.2
New Mexico	133	141	8.2	1	1	0.3	134	142	6.8
New York	3,529	3,817	22.8	7	7	0.2	3,536	3,825	19.4
North Carolina	1,338	1,416	17.0	10	11	0.6	1,348	1,426	14.3
North Dakota	23	24	4.0	0	0	0.0	23	24	3.3
Ohio	948	1,005	10.3	3	3	0.2	951	1,008	8.7
Oklahoma	303	334	10.5	6	6	0.9	309	340	8.8
Oregon	231	244	7.3	0	0	0.0	231	244	6.1
Pennsylvania	1,238	1,322	12.2	3	3	0.2	1,241	1,325	10.4
Rhode Island	96	101	11.2	0	0	0.0	96	101	9.6
South Carolina	792	841	20.7	5	5	0.7	797	846	17.5
South Dakota	27	28	4.1	0	0	0.0	27	28	3.3
Tennessee	761	808	14.7	9	9	0.9	770	817	12.5
Texas	4,377	4,817	22.1	15	16	0.3	4,392	4,833	17.9
Utah	114	120	5.3	0	0	0.0	114	120	4.1
Vermont	18	19	3.5	0	0	0.0	18	19	3.1
Virginia	927	983	14.1	5	5	0.4	932	989	11.9
Washington	431	455	7.7	3	3	0.3	434	459	6.5
West Virginia	88	96	6.1	0	0	0.0	88	96	5.2
Wisconsin	217	230	4.8	1	1	0.1	218	231	4.0
Wyoming	11	12	2.4	1	1	1.0	12	13	2.2
Subtotal	39,718	43,899	16.5	159	174	0.3	39,877	44,073	13.8
U.S. dependent areas									
American Samoa	0	0	0.0	0	0	0.0	0	0	0.0
Guam	0	0	0.0	0	0	0.0	0	0	0.0
Northern Mariana Islands	0	0	0.0	0	0	0.0	0	0	0.0
Puerto Rico	594	686	22.7	1	1	0.2	595	687	19.4
Republic of Palau	0	0	0.0	0	0	0.0	0	0	0.0
U.S. Virgin Islands	21	24	27.4	0	0	0.0	21	24	23.1
Subtotal	615	710	21.3	1	1	0.2	616	711	18.0
Total^b	40,333	44,608	16.6	160	176	0.3	40,493	44,784	13.9

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^b Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 23. Stage 3 (AIDS), by area of residence, 2014 and cumulative—United States and 6 dependent areas

Area of residence	2014									Cumulative ^a		
	Adults and adolescents			Children			Total			Estimated ^b		
	Estimated ^b			Estimated ^b			Estimated ^b			Adults and adolescents	Children	Total
	No.	No.	Rate ^c	No.	No.	Rate ^c	No.	No.	Rate ^c			
Alabama	306	356	8.8	1	1	0.2	307	357	7.4	11,842	77	11,919
Alaska	14	15	2.5	0	0	0.0	14	15	2.0	819	7	826
Arizona	305	322	5.8	2	2	0.2	307	324	4.8	13,813	48	13,861
Arkansas	136	149	6.1	0	0	0.0	136	149	5.0	5,124	39	5,163
California	1,984	2,213	6.9	5	6	0.1	1,989	2,219	5.7	172,642	706	173,348
Colorado	181	189	4.3	0	0	0.0	181	189	3.5	10,875	34	10,909
Connecticut	176	191	6.2	0	0	0.0	176	191	5.3	16,944	189	17,133
Delaware	76	83	10.5	0	0	0.0	76	83	8.9	4,467	28	4,495
District of Columbia	185	219	38.5	3	4	4.1	188	223	33.9	21,295	196	21,492
Florida	2,424	2,568	15.1	9	10	0.3	2,433	2,578	13.0	132,488	1,568	134,056
Georgia	943	1,133	13.6	3	4	0.2	946	1,137	11.3	47,658	262	47,921
Hawaii	43	47	3.9	0	0	0.0	43	47	3.3	3,539	17	3,556
Idaho	17	19	1.4	0	0	0.0	17	19	1.2	828	4	832
Illinois	699	777	7.2	4	4	0.2	703	781	6.1	42,684	289	42,973
Indiana	255	270	4.9	3	3	0.3	258	274	4.1	10,895	60	10,956
Iowa	61	65	2.5	0	0	0.0	61	65	2.1	2,356	14	2,370
Kansas	69	75	3.1	0	0	0.0	69	75	2.6	3,550	17	3,567
Kentucky	167	180	4.9	1	1	0.1	168	181	4.1	6,393	39	6,432
Louisiana	609	635	16.5	2	2	0.3	611	637	13.7	23,988	146	24,134
Maine	21	25	2.2	0	0	0.0	21	25	1.9	1,395	9	1,404
Maryland	615	684	13.7	0	0	0.0	615	684	11.4	39,656	338	39,994
Massachusetts	178	201	3.5	1	1	0.1	179	202	3.0	23,684	229	23,913
Michigan	362	390	4.7	2	2	0.1	364	393	4.0	19,262	119	19,381
Minnesota	153	160	3.5	3	3	0.3	156	164	3.0	6,578	33	6,611
Mississippi	301	315	12.8	2	2	0.4	303	317	10.6	9,355	58	9,413
Missouri	242	256	5.1	1	1	0.1	243	257	4.2	13,971	66	14,037
Montana	7	7	0.8	0	0	0.0	7	7	0.7	532	3	535
Nebraska	41	43	2.8	0	0	0.0	41	43	2.3	2,005	12	2,017
Nevada	208	220	9.3	0	0	0.0	208	220	7.8	7,775	29	7,804
New Hampshire	15	16	1.4	0	0	0.0	15	16	1.2	1,329	10	1,339
New Jersey	682	723	9.6	1	1	0.1	683	724	8.1	57,734	806	58,541
New Mexico	61	64	3.7	0	0	0.0	61	64	3.1	3,411	9	3,420
New York	1,662	1,914	11.4	4	5	0.2	1,666	1,919	9.7	203,817	2,443	206,260
North Carolina	710	749	9.0	1	1	0.1	711	750	7.5	23,407	140	23,548
North Dakota	11	12	1.9	0	0	0.0	11	12	1.6	210	2	212

Table 23. Stage 3 (AIDS), by area of residence, 2014 and cumulative—United States and 6 dependent areas (cont)

Area of residence	2014									Cumulative ^a		
	Adults and adolescents			Children			Total			Estimated ^b		
	Estimated ^b			Estimated ^b			Estimated ^b			Adults and adolescents	Children	Total
	No.	No.	Rate ^c	No.	No.	Rate ^c	No.	No.	Rate ^c			
Ohio	410	442	4.5	0	0	0.0	410	442	3.8	20,145	153	20,298
Oklahoma	156	170	5.3	1	1	0.2	157	171	4.4	6,332	29	6,361
Oregon	129	139	4.2	1	1	0.2	130	141	3.5	7,482	19	7,502
Pennsylvania	629	673	6.2	0	0	0.0	629	673	5.3	41,681	373	42,054
Rhode Island	31	33	3.7	0	0	0.0	31	33	3.1	3,156	29	3,185
South Carolina	369	389	9.6	1	1	0.2	370	391	8.1	18,157	119	18,276
South Dakota	10	11	1.5	1	1	0.7	11	12	1.4	368	7	375
Tennessee	405	450	8.2	31	34	3.2	436	484	7.4	17,130	97	17,227
Texas	2,079	2,254	10.3	8	9	0.2	2,087	2,263	8.4	92,408	407	92,816
Utah	39	41	1.8	0	0	0.0	39	41	1.4	2,831	21	2,852
Vermont	9	10	1.8	0	0	0.0	9	10	1.6	556	6	562
Virginia	406	501	7.2	3	4	0.3	409	505	6.1	22,097	191	22,288
Washington	202	219	3.7	0	0	0.0	202	219	3.1	14,423	34	14,457
West Virginia	50	56	3.5	0	0	0.0	50	56	3.0	1,994	11	2,005
Wisconsin	104	110	2.3	0	0	0.0	104	110	1.9	5,836	38	5,874
Wyoming	4	4	0.9	0	0	0.0	4	4	0.7	331	2	333
Subtotal	18,951	20,792	7.8	94	104	0.2	19,045	20,896	6.6	1,201,247	9,588	1,210,835
U.S. dependent areas												
American Samoa	0	0	0.0	0	0	0.0	0	0	0.0	1	0	1
Guam	0	0	0.0	0	0	0.0	0	0	0.0	86	1	87
Northern Mariana Islands	0	0	0.0	0	0	0.0	0	0	0.0	2	0	2
Puerto Rico	305	404	13.4	0	0	0.0	305	404	11.4	34,841	404	35,245
Republic of Palau	0	0	0.0	0	0	0.0	0	0	0.0	2	0	2
U.S. Virgin Islands	14	17	19.8	0	0	0.0	14	17	16.7	789	21	810
Subtotal	319	422	12.6	0	0	0.0	319	422	10.7	35,720	426	36,146
Total^d	19,270	21,214	7.9	94	104	0.2	19,364	21,318	6.6	1,236,967	10,014	1,246,981

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a From the beginning of the epidemic through 2014.

^b Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting.

^c Rates are per 100,000 population.

^d Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 24. Adults and adolescents living with diagnosed HIV infection, by race/ethnicity and area of residence, year-end 2013—United States and 6 dependent areas

Area of residence	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	No.	Estimated ^c		No.	Estimated ^c		No.	Estimated ^c		No.	Estimated ^c		No.	Estimated ^c		No.	Estimated ^c		No.	Estimated ^c		No.	Estimated ^c	
		No.	Rate		No.	No.		Rate	No.		No.	Rate		No.	No.		Rate	No.		No.	Rate		No.	No.
Alabama	5	5	21.5	28	28	55.5	7,854	7,778	749.7	337	337	245.3	4	4	230.2	3,422	3,382	122.9	497	492	1,175.7	12,147	12,025	297.4
Alaska	145	144	180.0	15	15	43.2	74	73	335.5	67	67	187.1	3	3	47.6	303	301	76.8	24	23	77.7	631	626	104.3
Arizona	508	511	243.5	134	135	81.9	1,369	1,371	641.8	3,927	3,947	262.6	14	14	142.5	7,066	7,056	214.5	251	250	314.7	13,269	13,284	242.8
Arkansas	4	4	22.0	14	14	41.3	2,198	2,192	604.9	298	299	209.8	3	3	62.4	2,302	2,288	123.3	206	206	656.4	5,025	5,006	204.4
California	402	404	287.6	4,301	4,372	97.0	20,624	20,792	1,115.1	40,564	41,142	363.6	278	282	241.3	50,179	50,336	380.2	2,511	2,518	371.0	118,859	119,845	376.2
Colorado	63	63	219.7	102	103	80.3	1,626	1,629	967.1	2,372	2,382	288.9	10	10	185.5	7,237	7,234	230.0	204	204	287.6	11,614	11,624	266.0
Connecticut	8	8	132.5	78	79	62.6	3,354	3,332	1,144.2	3,368	3,348	827.2	4	4	391.8	3,315	3,286	149.9	199	198	542.9	10,326	10,255	335.5
Delaware	3	3	118.0	7	7	25.9	1,882	1,877	1,178.9	241	241	413.6	0	0	0.0	840	840	161.7	85	85	740.0	3,058	3,053	392.2
District of Columbia	12	12	929.9	79	81	356.8	11,010	11,158	4,247.2	1,005	1,024	1,932.0	9	9	3,781.9	2,411	2,447	1,151.0	439	443	4,265.9	14,965	15,173	2,695.8
Florida	63	63	144.2	388	392	89.2	48,599	48,631	1,978.4	21,469	21,533	566.2	31	31	306.6	28,697	28,653	292.7	2,160	2,148	1,110.8	101,407	101,452	606.1
Georgia	17	18	89.7	164	172	57.5	28,045	29,004	1,178.3	2,466	2,562	393.2	12	13	261.6	8,344	8,564	183.4	1,700	1,735	1,757.3	40,748	42,067	512.7
Hawaii	8	8	261.4	392	395	85.3	112	112	454.8	241	242	246.0	242	243	231.7	1,229	1,226	420.3	203	203	103.6	2,427	2,430	205.6
Idaho	12	12	82.3	5	5	29.1	40	40	500.0	115	115	86.1	0	0	0.0	659	659	59.5	24	24	126.5	855	856	65.7
Illinois	38	38	241.0	387	393	71.9	16,280	16,280	1,085.4	6,118	6,179	388.5	21	21	732.4	9,981	10,010	143.6	1,751	1,760	1,554.7	34,576	34,681	322.9
Indiana	6	6	47.3	70	71	69.9	3,469	3,480	726.2	775	781	262.3	0	0	0.0	4,943	4,935	110.0	323	323	520.2	9,586	9,595	176.4
Iowa	2	2	28.1	44	44	84.1	349	350	473.8	189	190	162.7	0	0	0.0	1,295	1,296	56.5	69	69	276.3	1,948	1,952	75.9
Kansas	9	9	45.4	24	24	38.0	728	730	538.2	444	446	197.0	2	2	115.5	1,461	1,457	77.3	140	140	343.0	2,808	2,807	118.4
Kentucky	5	5	62.9	31	31	68.3	1,908	1,896	669.7	354	356	351.2	1	1	55.5	3,361	3,341	104.7	219	219	551.4	5,879	5,849	159.4
Louisiana	21	21	88.3	63	64	96.1	13,087	13,109	1,108.1	828	832	495.3	4	4	259.0	4,837	4,829	206.0	342	341	935.0	19,182	19,201	502.2
Maine	7	7	98.9	4	4	30.8	135	136	1,079.9	60	60	421.0	0	0	0.0	973	973	89.6	13	13	96.5	1,192	1,193	104.1
Maryland	24	25	202.5	180	187	61.8	23,269	23,880	1,664.3	1,632	1,683	418.4	12	12	500.9	4,583	4,681	170.9	1,390	1,423	1,763.3	31,090	31,890	641.1
Massachusetts	16	16	161.9	305	321	94.5	5,336	5,513	1,498.0	4,568	4,715	873.4	2	2	95.3	7,739	7,936	180.5	326	336	475.6	18,292	18,839	329.0
Michigan	33	33	71.0	94	95	42.9	8,958	8,920	791.6	764	764	227.7	2	2	94.5	5,158	5,138	79.4	488	487	395.1	15,497	15,439	185.4
Minnesota	108	108	242.9	109	109	57.2	2,455	2,462	1,107.8	689	692	367.7	6	6	345.9	3,720	3,721	98.3	207	207	330.1	7,294	7,306	162.4
Mississippi	9	9	79.1	13	13	53.7	6,675	6,608	747.1	249	249	385.6	2	2	222.9	1,794	1,775	121.6	380	380	2,107.2	9,122	9,036	366.9
Missouri	10	10	47.1	66	67	73.0	5,234	5,241	930.1	631	633	379.8	2	2	39.1	5,742	5,727	138.8	292	289	414.6	11,977	11,968	237.3
Montana	23	23	50.4	1	1	16.3	13	13	313.2	18	18	73.3	0	0	0.0	323	323	42.7	18	18	114.2	396	396	46.4
Nebraska	21	21	183.7	23	23	75.1	493	495	747.6	257	259	201.0	2	2	227.2	990	993	78.0	46	46	268.7	1,832	1,839	120.3
Nevada	43	43	208.9	205	207	109.8	1,687	1,686	904.7	1,783	1,793	312.2	22	22	165.7	3,739	3,727	291.4	209	210	385.1	7,688	7,689	331.8
New Hampshire	0	0	0.0	15	15	58.1	150	151	1,140.6	148	147	470.7	0	0	0.0	830	825	78.5	40	40	330.5	1,183	1,178	103.6
New Jersey	14	14	131.1	326	333	49.6	17,603	17,704	1,855.5	9,805	9,921	749.7	10	10	435.7	7,598	7,649	172.3	2,185	2,201	2,745.5	37,541	37,832	505.8
New Mexico	216	217	153.0	9	9	35.2	132	132	414.0	1,314	1,319	171.5	0	0	0.0	949	946	129.7	54	54	259.6	2,674	2,677	155.7
New York	79	79	165.3	1,605	1,619	118.6	52,108	52,210	2,172.8	43,278	43,356	1,502.8	17	17	243.6	26,038	26,088	267.5	7,308	7,321	3,555.2	130,433	130,691	784.1
North Carolina	176	174	189.1	121	122	60.7	17,265	17,144	992.6	1,836	1,846	303.8	11	11	224.2	6,697	6,663	121.8	809	809	809.7	26,915	26,770	326.3
North Dakota	9	9	33.2	2	2	26.6	44	44	453.7	14	14	96.1	0	0	0.0	139	139	25.9	10	10	132.8	218	218	36.2

Table 24. Adults and adolescents living with diagnosed HIV infection, by race/ethnicity and area of residence, year-end 2013—United States and 6 dependent areas (cont)

Area of residence	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total				
	Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c				
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^d
Ohio	15	15	78.4	76	77	42.1	8,535	8,562	751.3	1,055	1,059	378.7	1	1	34.0	9,026	9,038	113.8	684	688	544.7	19,392	19,441	200.5		
Oklahoma	280	277	111.3	43	44	69.7	1,301	1,300	569.7	493	494	192.8	5	5	138.4	3,047	3,028	136.0	286	285	208.1	5,455	5,433	171.8		
Oregon	44	44	119.9	91	92	68.2	370	371	660.1	736	740	213.7	13	13	114.0	4,010	4,012	151.4	111	111	138.3	5,375	5,383	162.3		
Pennsylvania	33	33	221.1	206	208	63.2	15,929	15,863	1,427.9	5,032	5,019	838.4	23	23	834.5	10,010	9,964	114.6	1,443	1,436	1,275.4	32,676	32,547	299.6		
Rhode Island	8	8	217.7	32	32	111.1	502	500	1,045.2	566	567	522.9	1	1	183.1	1,013	1,010	144.1	51	51	391.2	2,173	2,168	240.1		
South Carolina	11	11	74.5	31	31	54.6	10,959	10,966	1,027.4	617	622	339.4	2	2	96.1	3,604	3,599	136.9	319	320	771.9	15,543	15,550	389.3		
South Dakota	79	79	154.3	5	5	61.7	100	101	861.0	21	21	105.6	0	0	0.0	279	280	47.4	8	8	83.6	492	494	71.3		
Tennessee	7	7	45.6	58	58	67.0	9,297	9,140	1,037.5	695	698	316.4	9	9	346.5	5,872	5,773	138.6	450	448	761.1	16,388	16,133	297.2		
Texas	51	51	68.7	636	646	69.8	26,807	26,918	1,074.0	22,531	22,694	297.2	19	19	114.4	21,198	21,171	211.5	2,454	2,459	1,124.5	73,696	73,959	345.8		
Utah	20	20	90.8	36	36	68.5	215	216	965.2	496	498	181.2	4	4	20.6	1,725	1,729	95.2	62	62	198.7	2,558	2,565	114.6		
Vermont	0	0	0.0	3	3	38.1	44	44	814.7	24	24	281.2	1	1	669.2	363	361	70.6	12	12	170.7	447	445	82.2		
Virginia	16	16	84.0	240	240	57.9	12,915	12,743	981.1	1,731	1,730	320.7	6	6	119.9	6,477	6,408	141.6	638	630	521.5	22,023	21,774	314.5		
Washington	140	140	188.5	354	357	77.7	1,655	1,663	794.4	1,468	1,478	250.8	45	46	132.8	7,406	7,414	173.0	343	343	201.4	11,411	11,441	196.5		
West Virginia	2	2	59.7	4	4	32.7	426	424	775.1	66	66	335.1	0	0	0.0	1,092	1,094	74.2	74	74	449.5	1,664	1,664	105.3		
Wisconsin	28	28	70.9	73	74	65.2	2,071	2,067	741.8	701	703	276.7	1	1	71.8	2,506	2,506	61.5	156	157	320.0	5,536	5,535	115.0		
Wyoming	7	7	76.4	3	3	66.2	14	14	173.6	40	40	96.5	0	0	0.0	173	173	41.9	12	12	199.9	249	249	51.6		
Subtotal	2,860	2,862	152.4	11,295	11,462	83.2	395,305	397,067	1,243.8	188,496	189,946	462.0	856	866	201.3	296,695	297,003	174.0	32,225	32,319	815.1	927,732	931,526	353.2		
U.S. dependent areas^e																										
American Samoa	0	0	—	1	1	—	0	0	—	0	0	—	0	0	—	0	0	—	0	0	—	1	1	2.4		
Guam	0	0	—	47	47	—	5	5	—	4	4	—	10	10	—	9	9	—	7	7	—	82	82	65.8		
Northern Mariana Islands	0	0	—	0	0	—	0	0	—	0	0	—	2	2	—	0	0	—	0	0	—	2	2	5.2		
Puerto Rico	1	1	—	3	3	—	7	7	—	18,479	18,548	—	1	1	—	14	14	—	0	0	—	18,505	18,574	610.0		
Republic of Palau	0	0	—	0	0	—	1	1	—	1	1	—	2	2	—	0	0	—	0	0	—	4	4	23.4		
U.S. Virgin Islands	0	0	—	0	0	—	349	352	—	219	222	—	0	0	—	45	46	—	4	3	—	617	623	705.8		
Subtotal	1	1	—	51	51	—	362	365	—	18,703	18,775	—	15	15	—	68	68	—	11	10	—	19,211	19,286	574.5		
Total^f	2,861	2,863	—	11,346	11,513	—	395,667	397,432	—	207,199	208,721	—	871	882	—	296,763	297,071	—	32,236	32,329	—	946,943	950,811	355.9		

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 25. Adults and adolescents living with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and area of residence, year-end 2013—United States and 6 dependent areas

Area of residence	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^d	Rate
Alabama	0	0	0.0	12	12	24.0	3,287	3,280	316.2	168	169	122.8	1	1	56.4	1,352	1,340	48.7	223	223	533.8	5,043	5,025	124.3
Alaska	91	92	114.9	14	14	40.5	42	42	192.5	46	46	129.3	2	2	32.4	168	168	42.9	13	12	42.0	376	377	62.8
Arizona	282	283	134.7	60	61	36.7	637	638	298.4	1,953	1,959	130.4	11	11	112.4	3,381	3,376	102.6	132	132	165.7	6,456	6,459	118.1
Arkansas	3	3	17.1	7	7	20.9	994	996	274.9	144	145	101.6	2	2	41.4	1,105	1,100	59.3	126	127	405.2	2,381	2,381	97.2
California	229	229	163.0	2,448	2,478	55.0	12,643	12,702	681.2	25,160	25,406	224.5	145	147	125.5	30,560	30,561	230.9	1,585	1,591	234.4	72,770	73,113	229.5
Colorado	28	28	98.1	49	49	38.4	743	745	442.2	1,237	1,242	150.6	4	4	74.0	2,905	2,905	92.4	114	115	161.9	5,080	5,087	116.4
Connecticut	7	7	116.9	41	42	33.0	2,301	2,287	785.3	2,328	2,319	573.0	4	4	396.0	2,153	2,133	97.3	173	172	472.0	7,007	6,965	227.8
Delaware	2	2	79.0	5	5	18.8	1,234	1,236	776.6	151	152	261.0	0	0	0.0	519	521	100.3	48	48	418.7	1,959	1,965	252.4
District of Columbia	4	4	303.1	38	38	168.2	6,745	6,803	2,589.5	592	599	1,130.1	4	4	1,676.4	1,170	1,178	554.3	299	300	2,894.2	8,852	8,927	1,586.0
Florida	20	20	45.1	174	175	39.7	28,206	28,190	1,146.8	11,367	11,360	298.7	11	11	108.3	16,156	16,071	164.2	1,387	1,380	713.5	57,321	57,205	341.8
Georgia	8	8	40.8	73	76	25.2	15,256	15,644	635.5	1,439	1,483	227.5	4	4	87.6	4,495	4,552	97.5	1,121	1,144	1,158.8	22,396	22,910	279.2
Hawaii	3	3	97.2	234	237	51.3	74	74	302.7	151	152	154.7	163	165	156.8	730	732	250.7	125	126	64.3	1,480	1,489	126.0
Idaho	4	4	27.8	2	2	11.6	17	17	209.7	62	62	46.4	0	0	0.0	318	319	28.8	15	15	79.3	418	419	32.2
Illinois	14	14	86.3	193	193	35.4	8,716	8,588	572.6	3,474	3,471	218.2	11	11	385.4	5,009	4,969	71.3	1,014	1,012	894.2	18,431	18,259	170.0
Indiana	0	0	0.0	47	47	46.8	1,735	1,738	362.6	435	437	146.7	0	0	0.0	2,635	2,631	58.7	175	175	282.0	5,027	5,028	92.4
Iowa	1	1	14.2	19	19	36.5	236	237	321.2	120	120	102.9	0	0	0.0	778	776	33.8	41	41	163.1	1,195	1,195	46.5
Kansas	6	6	30.1	11	11	17.1	374	374	275.8	272	271	120.0	0	0	0.0	859	854	45.3	79	79	194.3	1,601	1,595	67.3
Kentucky	3	3	37.7	13	13	28.4	944	936	330.7	223	224	220.9	1	1	55.8	1,812	1,803	56.5	125	125	315.5	3,121	3,105	84.6
Louisiana	11	11	45.9	22	22	33.2	6,963	6,981	590.1	440	442	263.0	2	2	128.9	2,606	2,601	111.0	181	181	495.3	10,225	10,239	267.8
Maine	4	4	58.8	4	4	33.7	63	64	509.6	33	33	231.4	0	0	0.0	556	559	51.4	10	10	73.8	670	674	58.8
Maryland	5	5	42.0	80	82	27.0	13,037	13,236	922.5	980	996	247.6	6	6	247.0	2,371	2,395	87.4	900	911	1,129.7	17,379	17,630	354.4
Massachusetts	10	10	96.6	173	173	50.8	3,298	3,271	888.8	2,753	2,730	505.8	1	1	44.8	4,387	4,328	98.4	200	199	282.0	10,822	10,712	187.1
Michigan	13	13	27.9	42	43	19.3	4,691	4,686	415.9	424	425	126.5	2	2	94.7	2,683	2,677	41.4	280	281	227.4	8,135	8,126	97.6
Minnesota	56	56	126.3	54	54	28.4	1,156	1,160	522.0	375	376	199.9	1	1	57.4	1,583	1,584	41.8	108	108	171.4	3,333	3,339	74.2
Mississippi	3	3	24.6	6	6	24.6	3,122	3,067	346.7	131	130	202.0	1	1	111.2	815	798	54.7	220	220	1,220.0	4,298	4,225	171.5
Missouri	5	5	23.4	29	29	31.9	2,774	2,777	492.8	330	331	198.7	1	1	19.5	2,982	2,972	72.0	175	174	249.2	6,296	6,289	124.7
Montana	15	15	32.7	0	0	0.0	8	8	191.4	15	15	61.7	0	0	0.0	202	202	26.7	12	12	76.5	252	252	29.5
Nebraska	11	11	97.3	13	13	42.3	277	278	419.9	162	162	126.2	2	2	226.0	511	512	40.2	30	30	173.6	1,006	1,008	65.9
Nevada	24	24	118.1	91	92	48.5	834	834	447.7	902	904	157.5	11	11	81.6	1,747	1,740	136.0	117	118	217.3	3,726	3,724	160.7
New Hampshire	0	0	0.0	7	7	27.2	80	81	610.2	85	85	271.2	0	0	0.0	427	424	40.3	29	29	239.8	628	626	55.0
New Jersey	9	9	83.1	164	165	24.6	9,204	9,166	960.7	5,150	5,152	389.3	4	4	172.7	3,857	3,835	86.4	1,328	1,329	1,657.8	19,716	19,661	262.8
New Mexico	115	115	81.4	5	5	19.7	69	69	216.7	761	762	99.1	0	0	0.0	622	621	85.1	39	39	186.8	1,611	1,611	93.7
New York	40	41	84.8	800	814	59.6	31,913	32,399	1,348.3	27,161	27,501	953.2	3	3	43.5	13,922	14,125	144.8	4,726	4,810	2,335.8	78,565	79,693	478.1
North Carolina	77	76	82.5	37	37	18.5	7,336	7,271	420.9	831	832	136.9	1	1	19.9	2,706	2,683	49.0	364	364	364.5	11,352	11,264	137.3
North Dakota	8	8	29.7	2	2	26.5	17	17	173.3	7	7	48.5	0	0	0.0	60	60	11.2	5	5	67.7	99	99	16.5

Table 25. Adults and adolescents living with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and area of residence, year-end 2013—United States and 6 dependent areas (cont)

Area of residence	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^d	Rate
Ohio	7	7	37.3	29	29	16.1	4,068	4,118	361.4	528	534	190.9	0	0	0.1	4,325	4,370	55.0	348	353	279.1	9,305	9,411	97.1
Oklahoma	147	146	58.5	22	22	35.6	594	596	261.0	232	233	90.7	3	3	82.9	1,482	1,476	66.3	129	129	94.2	2,609	2,604	82.3
Oregon	25	25	68.3	52	52	39.1	230	232	411.8	495	501	144.4	8	8	70.5	2,489	2,503	94.4	72	72	89.7	3,371	3,392	102.3
Pennsylvania	10	10	66.6	110	110	33.5	9,308	9,225	830.4	2,976	2,959	494.2	10	10	359.3	5,642	5,591	64.3	920	914	811.3	18,976	18,819	173.2
Rhode Island	5	5	136.0	24	24	83.6	354	353	737.9	391	390	360.0	1	1	183.5	645	644	91.8	32	32	246.4	1,452	1,449	160.4
South Carolina	5	5	31.5	19	19	33.5	6,084	6,094	571.0	359	360	196.8	2	2	95.9	1,845	1,846	70.2	192	193	465.8	8,506	8,519	213.3
South Dakota	35	35	68.8	2	2	24.5	42	42	361.2	8	8	40.1	0	0	0.0	102	102	17.3	5	5	52.3	194	195	28.1
Tennessee	4	4	24.3	25	24	28.0	4,502	4,352	494.1	390	390	176.6	2	2	76.9	3,060	2,960	71.1	280	279	474.3	8,263	8,012	147.6
Texas	26	26	34.7	305	308	33.3	14,200	14,215	567.1	13,021	13,045	170.8	8	8	46.8	12,086	11,993	119.8	1,433	1,433	655.0	41,079	41,027	191.8
Utah	15	15	68.0	19	19	35.7	114	114	511.8	287	288	104.7	2	2	10.5	960	961	52.9	36	36	116.0	1,433	1,435	64.1
Vermont	0	0	0.0	2	2	25.8	28	28	514.0	15	15	176.2	1	1	670.5	209	208	40.6	6	6	85.6	261	260	47.9
Virginia	8	8	42.4	123	125	30.0	5,692	5,627	433.2	888	896	166.1	3	3	60.4	3,025	2,987	66.0	310	307	254.4	10,049	9,953	143.7
Washington	92	93	124.5	199	202	43.9	971	980	468.3	875	884	150.1	25	25	73.9	4,177	4,208	98.2	193	194	114.1	6,532	6,587	113.1
West Virginia	0	0	0.0	0	0	0.1	196	197	360.6	29	29	146.5	0	0	0.0	639	643	43.6	55	55	338.2	919	925	58.5
Wisconsin	12	12	30.2	30	30	26.6	1,000	1,000	358.9	383	384	151.0	0	0	0.0	1,218	1,218	29.9	73	73	149.2	2,716	2,717	56.5
Wyoming	5	5	54.3	1	1	25.6	7	7	87.0	26	26	62.3	0	0	0.0	101	100	24.3	11	11	184.5	151	151	31.2
Subtotal	1,507	1,506	80.2	5,931	5,997	43.5	216,416	217,074	680.0	110,765	111,441	271.1	463	468	108.8	160,147	159,913	93.7	19,614	19,730	497.6	514,843	516,129	195.7
U.S. dependent areas^e																								
American Samoa	0	0	—	0	0	—	0	0	—	0	0	—	0	0	—	0	0	—	1	1	—	1	1	2.4
Guam	0	0	—	22	22	—	1	1	—	0	0	—	6	6	—	1	1	—	2	2	—	32	32	25.6
Northern Mariana Islands	0	0	—	0	0	—	0	0	—	0	0	—	2	2	—	0	0	—	0	0	—	2	2	5.3
Puerto Rico	1	1	—	0	0	—	2	2	—	10,668	10,651	—	0	0	—	10	10	—	0	0	—	10,681	10,664	350.2
Republic of Palau	0	0	—	0	0	—	0	0	—	0	0	—	1	1	—	0	0	—	0	0	—	1	1	5.7
U.S. Virgin Islands	0	0	—	0	0	—	191	191	—	118	118	—	0	0	—	29	29	—	2	2	—	340	340	385.8
Subtotal	1	1	—	22	22	—	194	194	—	10,786	10,769	—	9	9	—	40	40	—	5	5	—	11,057	11,040	328.9
Total^f	1,508	1,507	—	5,953	6,019	—	216,610	217,268	—	121,551	122,211	—	472	477	—	160,187	159,953	—	19,619	19,734	—	525,900	527,170	197.4

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 26. Diagnoses of HIV infection, 2014, and persons living with diagnosed HIV infection (prevalence), year-end 2013, by metropolitan statistical area of residence—United States and Puerto Rico

MSA of residence	Diagnosis, 2014				Prevalence of diagnosed HIV infection, year-end 2013		
	No.	Estimated ^a		Rank ^b	No.	Estimated ^a	
		No.	Rate			No.	Rate
Akron, OH	47	50	7.1	88	852	855	121.6
Albany–Schenectady–Troy, NY	54	57	6.5	92	2,099	2,102	239.2
Albuquerque, NM	67	71	7.8	83	1,339	1,341	148.4
Allentown–Bethlehem–Easton, PA–NJ	58	62	7.5	86	1,471	1,470	177.8
Atlanta–Sandy Springs–Roswell, GA	1,095	1,454	25.9	7	27,136	27,993	506.6
Augusta–Richmond County, GA–SC	77	99	17.0	26	2,168	2,216	381.9
Austin–Round Rock, TX	303	334	17.2	25	4,949	4,951	262.6
Bakersfield, CA	111	123	14.0	42	1,779	1,778	205.3
Baltimore–Columbia–Towson, MD	549	678	24.3	10	18,375	18,821	678.5
Baton Rouge, LA	350	369	44.7	1	4,584	4,594	560.0
Birmingham–Hoover, AL	152	168	14.6	37	3,842	3,805	333.9
Boise City, ID	15	16	2.4	104	466	467	71.7
Boston–Cambridge–Newton, MA–NH ^c	411	555	11.7	53	12,961	13,357	284.3
Boston Division	246	334	17.0	—	7,723	7,957	407.6
Cambridge Division	157	212	9.1	—	4,944	5,108	219.9
Bridgeport–Stamford–Norwalk, CT	95	101	10.7	60	2,716	2,702	286.8
Buffalo–Cheektowaga–Niagara Falls, NY	110	116	10.2	64	2,316	2,321	204.3
Cape Coral–Fort Myers, FL	96	102	14.9	34	1,691	1,691	255.6
Charleston–North Charleston, SC	140	148	20.3	18	2,366	2,366	332.3
Charlotte–Concord–Gastonia, NC–SC	427	451	19.0	22	7,222	7,186	307.4
Chattanooga, TN–GA	58	62	11.4	57	1,155	1,141	210.4
Chicago–Naperville–Elgin, IL–IN–WI	1,347	1,544	16.2	30	30,562	30,665	321.3
Chicago Division	1,190	1,373	18.7	—	27,851	27,949	380.8
Elgin Division	52	57	9.0	—	630	632	100.5
Gary Division	60	64	9.1	—	1,229	1,231	174.5
Lake County Division	45	50	5.7	—	852	854	98.0
Cincinnati, OH–KY–IN	209	221	10.3	62	3,602	3,611	168.9
Cleveland–Elyria, OH	232	245	11.9	51	4,951	4,960	240.2
Colorado Springs, CO	40	42	6.1	93	819	818	120.5
Columbia, SC	194	205	25.6	8	4,030	4,034	509.1
Columbus, OH	244	258	13.0	46	5,221	5,241	266.2
Dallas–Fort Worth–Arlington, TX	1,311	1,498	21.5	17	22,530	22,796	334.1
Dallas Division	1,068	1,231	26.7	—	17,883	18,155	402.4
Fort Worth Division	243	267	11.4	—	4,647	4,641	200.8
Dayton, OH	67	70	8.8	74	1,481	1,487	185.4
Deltona–Daytona Beach–Ormond Beach, FL	108	114	18.7	23	1,496	1,493	248.3
Denver–Aurora–Lakewood, CO	283	299	10.9	58	9,111	9,120	337.8
Des Moines–West Des Moines, IA	35	37	6.0	94	620	623	103.9

Table 26. Diagnoses of HIV infection, 2014, and persons living with diagnosed HIV infection (prevalence), year-end 2013, by metropolitan statistical area of residence—United States and Puerto Rico (cont)

MSA of residence	Diagnosis, 2014				Prevalence of diagnosed HIV infection, year-end 2013		
	No.	Estimated ^a		Rank ^b	No.	Estimated ^a	
		No.	Rate			No.	Rate
Detroit–Warren–Dearborn, MI	529	559	13.0	45	10,118	10,073	234.5
Detroit Division	364	385	21.8	—	7,341	7,293	410.7
Warren Division	165	174	6.9	—	2,777	2,780	110.3
Durham–Chapel Hill, NC	80	84	15.5	33	2,050	2,036	381.5
El Paso, TX	108	118	14.1	41	1,879	1,883	225.6
Fayetteville–Springdale–Rogers, AR–MO	19	21	4.1	101	537	536	108.8
Fresno, CA	110	121	12.5	48	1,637	1,634	170.9
Grand Rapids–Wyoming, MI	65	69	6.7	90	1,018	1,018	100.1
Greensboro–High Point, NC	114	120	16.1	31	2,413	2,403	324.1
Greenville–Anderson–Mauldin, SC	106	113	13.1	44	1,677	1,679	197.4
Harrisburg–Carlisle, PA	49	52	9.2	71	1,347	1,342	240.6
Hartford–West Hartford–East Hartford, CT	90	96	7.9	82	3,496	3,469	285.3
Honolulu (Urban), HI	77	81	8.2	79	1,678	1,680	170.2
Houston–The Woodlands–Sugar Land, TX	1,474	1,567	24.1	11	25,120	25,153	397.1
Indianapolis–Carmel–Anderson, IN	242	255	13.0	47	4,775	4,784	245.0
Jackson, MS	176	186	32.2	4	3,440	3,404	589.7
Jacksonville, FL	336	357	25.1	9	6,300	6,301	451.4
Kansas City, MO–KS	181	191	9.2	70	4,828	4,824	234.7
Knoxville, TN	44	47	5.5	98	1,202	1,184	139.0
Lakeland–Winter Haven, FL	89	94	14.9	35	1,805	1,804	289.5
Lancaster, PA	28	30	5.6	97	773	771	145.4
Las Vegas–Henderson–Paradise, NV	387	412	19.9	19	6,453	6,459	318.3
Little Rock–North Little Rock–Conway, AR	147	161	22.1	15	1,946	1,942	268.1
Los Angeles–Long Beach–Anaheim, CA	2,235	2,578	19.4	20	52,832	53,648	407.2
Anaheim Division	281	325	10.3	—	6,796	6,873	220.2
Los Angeles Division	1,954	2,253	22.3	—	46,036	46,775	465.2
Louisville/Jefferson County, KY–IN	174	185	14.6	38	3,092	3,071	243.3
Madison, WI	25	26	4.2	100	690	690	110.0
McAllen–Edinburg–Mission, TX	78	85	10.3	61	1,076	1,079	131.7
Memphis, TN–MS–AR	348	370	27.6	6	7,404	7,293	543.5
Miami–Fort Lauderdale–West Palm Beach, FL	2,399	2,535	42.8	2	52,663	52,709	898.9
Fort Lauderdale Division	792	836	44.7	—	17,072	17,086	925.8
Miami Division	1,257	1,328	49.9	—	27,614	27,650	1,046.6
West Palm Beach Division	350	371	26.5	—	7,977	7,974	579.4
Milwaukee–Waukesha–West Allis, WI	133	140	8.9	73	3,102	3,097	197.3
Minneapolis–St. Paul–Bloomington, MN–WI	272	288	8.2	77	6,480	6,492	187.5
Modesto, CA	16	17	3.3	103	542	540	102.6
Nashville–Davidson–Murfreesboro–Franklin, TN	228	242	13.5	43	5,144	5,069	288.3

Table 26. Diagnoses of HIV infection, 2014, and persons living with diagnosed HIV infection (prevalence), year-end 2013, by metropolitan statistical area of residence—United States and Puerto Rico (cont)

MSA of residence	Diagnosis, 2014				Prevalence of diagnosed HIV infection, year-end 2013		
	No.	Estimated ^a		Rank ^b	No.	Estimated ^a	
		No.	Rate			No.	Rate
New Haven–Milford, CT	81	86	9.9	68	3,130	3,107	360.0
New Orleans–Metairie, LA	437	462	36.9	3	8,372	8,368	673.7
New York–Newark–Jersey City, NY–NJ–PA ^c	4,077	4,496	22.4	14	144,508	144,963	724.7
Nassau County Division	266	285	10.0	—	6,009	6,037	211.2
New York Division	3,286	3,605	25.2	—	122,139	122,464	859.7
Newark Division	494	572	22.8	—	15,053	15,153	605.7
North Port–Sarasota–Bradenton, FL	102	108	14.5	39	1,705	1,705	232.7
Ogden–Clearfield, UT	13	14	2.2	105	302	303	48.6
Oklahoma City, OK	148	164	12.2	50	2,668	2,660	201.5
Omaha–Council Bluffs, NE–IA	66	70	7.7	84	1,304	1,308	146.0
Orlando–Kissimmee–Sanford, FL	631	669	28.8	5	10,450	10,462	460.7
Oxnard–Thousand Oaks–Ventura, CA	52	57	6.7	89	903	900	107.0
Palm Bay–Melbourne–Titusville, FL	57	60	10.9	59	1,250	1,246	226.0
Philadelphia–Camden–Wilmington, PA–NJ–DE–MD	942	1019	16.8	28	26,647	26,578	440.3
Camden Division	140	164	13.1	—	3,118	3,150	251.5
Montgomery County Division	94	100	5.1	—	2,291	2,285	117.2
Philadelphia Division	620	657	30.9	—	18,773	18,679	881.9
Wilmington Division	88	98	13.7	—	2,465	2,464	344.0
Phoenix–Mesa–Scottsdale, AZ	604	638	14.2	40	10,084	10,100	229.3
Pittsburgh, PA	164	174	7.4	87	3,091	3,084	130.6
Portland–South Portland, ME	40	42	8.1	80	657	658	126.5
Portland–Vancouver–Hillsboro, OR–WA	183	193	8.2	78	4,432	4,440	191.8
Providence–Warwick, RI–MA	134	153	9.5	69	3,263	3,289	204.9
Provo–Orem, UT	3	3	0.6	107	151	151	26.9
Raleigh, NC	174	184	14.8	36	3,447	3,442	283.2
Richmond, VA	191	202	16.0	32	4,484	4,418	354.3
Riverside–San Bernardino–Ontario, CA	472	545	12.3	49	8,191	8,281	188.6
Rochester, NY	78	82	7.6	85	2,979	2,981	275.0
Sacramento–Roseville–Arden–Arcade, CA	171	187	8.3	75	3,829	3,818	172.2
St. Louis, MO–IL	297	321	11.4	55	6,957	6,975	249.0
Salt Lake City, UT	90	95	8.2	76	1,920	1,925	168.6
San Antonio–New Braunfels, TX	358	393	16.9	27	5,457	5,462	239.3
San Diego–Carlsbad, CA	460	530	16.3	29	12,542	12,672	393.2
San Francisco–Oakland–Hayward, CA ^c	725	798	17.4	24	24,729	24,755	546.5
Oakland Division	310	357	13.1	—	7,359	7,426	277.2
San Francisco Division	392	416	25.8	—	16,491	16,461	1,034.2
San Jose–Sunnyvale–Santa Clara, CA	159	175	9.0	72	3,285	3,276	169.9
San Juan–Carolina–Caguas, PR	458	527	23.5	12	13,200	13,248	583.2

Table 26. Diagnoses of HIV infection, 2014, and persons living with diagnosed HIV infection (prevalence), year-end 2013, by metropolitan statistical area of residence—United States and Puerto Rico (cont)

MSA of residence	Diagnosis, 2014				Prevalence of diagnosed HIV infection, year-end 2013		
	No.	Estimated ^a		Rank ^b	No.	Estimated ^a	
		No.	Rate			No.	Rate
Santa Rosa, CA	30	33	6.5	91	1,223	1,217	245.5
Scranton–Wilkes-Barre–Hazleton, PA	24	25	4.5	99	597	594	105.7
Seattle–Tacoma–Bellevue, WA	349	369	10.0	66	8,803	8,827	244.3
Seattle Division	305	322	11.3	—	7,756	7,776	278.4
Tacoma Division	44	46	5.6	—	1,047	1,051	128.1
Spokane–Spokane Valley, WA	6	6	1.2	106	441	442	82.6
Springfield, MA	47	63	10.0	67	1,883	1,938	308.4
Stockton–Lodi, CA	74	82	11.4	56	1,231	1,227	174.0
Syracuse, NY	72	76	11.5	54	1,124	1,127	170.1
Tampa–St. Petersburg–Clearwater, FL	632	669	22.9	13	10,688	10,682	371.7
Toledo, OH	46	49	8.0	81	980	981	161.2
Tucson, AZ	97	103	10.2	63	2,054	2,052	205.6
Tulsa, OK	104	114	11.7	52	1,680	1,672	173.7
Virginia Beach–Norfolk–Newport News, VA–NC	313	331	19.3	21	6,652	6,561	384.3
Washington–Arlington–Alexandria, DC–VA–MD–WV	1,108	1,304	21.6	16	33,081	33,567	562.5
Silver Spring Division	196	244	19.1	—	4,141	4,256	337.5
Washington Division	912	1,060	22.3	—	28,940	29,311	622.8
Wichita, KS	34	36	5.6	96	801	801	125.5
Winston-Salem, NC	63	66	10.1	65	1,719	1,705	261.8
Worcester, MA–CT	27	36	3.9	102	1812	1852	199.6
Youngstown–Warren–Boardman, OH–PA	31	33	6.0	95	685	685	123.1
Subtotal for MSAs (population of ≥500,000)	33,068	36,594	16.8	—	790,818	794,576	368.1
Metropolitan areas (population of 50,000–499,999)	4,784	5,232	9.0	—	97,640	97,723	168.8
Nonmetropolitan areas	2,338	2,547	5.5	—	52,051	51,977	112.1
Total^d	40,472	44,760	13.9	—	948,623	952,532	297.6

Abbreviation: MSA, metropolitan statistical area.

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Because of the lack of U.S. census information for all U.S. dependent areas, table includes data for only the United States and Puerto Rico.

MSA definitions for this report can be found at <http://www.census.gov/population/metro/>.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^b Based on estimated rate.

^c Counts of diagnoses of HIV infection for the metropolitan divisions do not sum to the MSA total. MSA total includes data from 1 metropolitan division with population of <500,000.

^d Includes persons whose county of residence is unknown. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 27. Stage 3 (AIDS), 2014 and cumulative, and persons living with diagnosed HIV infection ever classified as stage 3 (AIDS) (prevalence), year-end 2013, by metropolitan statistical area of residence—United States and Puerto Rico

MSA of residence	Diagnosis, 2014				Diagnosis, cumulative ^a			Prevalence of stage 3 (AIDS), year-end 2013		
	No.	Estimated ^b		Rank ^d	Adults or adolescents No.	Children No.	Total No.	No.	Estimated ^b	
		No.	Rate ^c						No.	No.
	No.	No.	Rate ^c	Rank ^d	No.	No.	No.	No.	No.	Rate ^c
Akron, OH	15	17	2.3	98	924	1	925	403	407	57.8
Albany–Schenectady–Troy, NY	28	33	3.8	75	2,707	27	2,734	1,182	1,219	138.8
Albuquerque, NM	34	35	3.9	72	1,832	3	1,835	820	820	90.8
Allentown–Bethlehem–Easton, PA–NJ	30	32	3.9	73	1,753	19	1,772	872	868	104.9
Atlanta–Sandy Springs–Roswell, GA	608	731	13.0	7	32,705	151	32,856	15,327	15,665	283.5
Augusta–Richmond County, GA–SC	44	51	8.7	24	2,635	24	2,659	1,132	1,154	198.8
Austin–Round Rock, TX	117	123	6.3	40	6,185	26	6,211	2,929	2,907	154.2
Bakersfield, CA	33	36	4.1	71	2,101	8	2,109	1,207	1,207	139.4
Baltimore–Columbia–Towson, MD	313	348	12.5	10	25,692	225	25,917	10,334	10,498	378.4
Baton Rouge, LA	172	178	21.6	1	5,545	24	5,569	2,521	2,532	308.6
Birmingham–Hoover, AL	90	105	9.2	23	3,449	25	3,474	1,429	1,427	125.2
Boise City, ID	10	11	1.7	102	423	0	423	218	218	33.6
Boston–Cambridge–Newton, MA–NH ^e	127	143	3.0	86	16,645	159	16,804	7,695	7,625	162.3
Boston Division	71	80	4.1	—	10,329	96	10,425	4,592	4,547	232.9
Cambridge Division	53	60	2.6	—	5,935	62	5,997	2,952	2,928	126.1
Bridgeport–Stamford–Norwalk, CT	58	63	6.6	37	4,338	58	4,396	1,850	1,841	195.4
Buffalo–Cheektowaga–Niagara Falls, NY	54	65	5.7	49	3,074	20	3,094	1,247	1,297	114.1
Cape Coral–Fort Myers, FL	50	53	7.8	27	2,236	25	2,261	1,003	1,002	151.6
Charleston–North Charleston, SC	46	48	6.6	39	2,697	24	2,721	1,269	1,272	178.7
Charlotte–Concord–Gastonia, NC–SC	249	264	11.1	16	5,838	30	5,868	2,948	2,937	125.6
Chattanooga, TN–GA	28	32	5.8	46	1,315	3	1,318	623	607	112.0
Chicago–Naperville–Elgin, IL–IN–WI	623	694	7.3	31	38,234	267	38,501	16,441	16,263	170.4
Chicago Division	562	628	8.6	—	35,008	244	35,252	14,949	14,767	201.2
Elgin Division	19	21	3.3	—	796	9	805	377	381	60.5
Gary Division	27	28	4.0	—	1,400	9	1,409	601	599	84.9
Lake County Division	15	16	1.9	—	1,031	5	1,036	514	516	59.2
Cincinnati, OH–KY–IN	93	100	4.6	62	3,790	20	3,810	1,844	1,861	87.0
Cleveland–Elyria, OH	70	76	3.7	78	5,227	48	5,275	2,387	2,409	116.7
Colorado Springs, CO	17	18	2.6	92	727	5	732	333	333	49.1
Columbia, SC	93	98	12.2	11	4,447	27	4,474	2,303	2,306	291.1
Columbus, OH	130	140	7.0	33	4,683	25	4,709	2,203	2,234	113.5

Table 27. Stage 3 (AIDS), 2014 and cumulative, and persons living with diagnosed HIV infection ever classified as stage 3 (AIDS) (prevalence), year-end 2013, by metropolitan statistical area of residence—United States and Puerto Rico (cont)

MSA of residence	Diagnosis, 2014				Diagnosis, cumulative ^a			Prevalence of stage 3 (AIDS), year-end 2013		
	Estimated ^b				Adults or adolescents	Children	Total	Estimated ^b		
	No.	No.	Rate ^c	Rank ^d	No.	No.	No.	No.	No.	Rate ^c
Dallas–Fort Worth–Arlington, TX	625	691	9.9	20	26,847	64	26,912	12,355	12,410	181.9
Dallas Division	478	537	11.7	—	21,249	38	21,287	9,855	9,933	220.2
Fort Worth Division	147	154	6.6	—	5,598	26	5,624	2,500	2,477	107.1
Dayton, OH	37	40	5.0	59	1,591	14	1,605	761	769	95.9
Deltona–Daytona Beach–Ormond Beach, FL	58	61	10.1	19	2,067	19	2,086	891	886	147.5
Denver–Aurora–Lakewood, CO	130	136	5.0	60	8,492	24	8,516	3,888	3,893	144.2
Des Moines–West Des Moines, IA	19	20	3.2	82	748	4	752	374	374	62.4
Detroit–Warren–Dearborn, MI	244	263	6.1	43	13,328	78	13,407	5,417	5,408	125.9
Detroit Division	182	196	11.1	—	10,447	61	10,509	4,024	4,011	225.9
Warren Division	62	67	2.7	—	2,881	17	2,898	1,393	1,397	55.4
Durham–Chapel Hill, NC	64	68	12.6	9	1,591	10	1,601	666	660	123.7
El Paso, TX	47	48	5.8	47	2,072	10	2,082	1,120	1,114	133.4
Fayetteville–Springdale–Rogers, AR–MO	14	16	3.1	84	513	7	520	273	272	55.2
Fresno, CA	41	45	4.6	63	2,118	11	2,129	1,008	1,007	105.4
Grand Rapids–Wyoming, MI	24	26	2.6	93	1,197	6	1,203	591	593	58.3
Greensboro–High Point, NC	30	32	4.3	69	1,752	20	1,772	802	796	107.3
Greenville–Anderson–Mauldin, SC	44	46	5.3	55	2,009	4	2,013	934	935	109.9
Harrisburg–Carlisle, PA	25	27	4.8	61	1,610	9	1,619	773	768	137.7
Hartford–West Hartford–East Hartford, CT	56	61	5.1	58	5,924	48	5,973	2,369	2,354	193.6
Honolulu (Urban), HI	31	33	3.4	81	2,403	13	2,416	1,017	1,024	103.7
Houston–The Woodlands–Sugar Land, TX	707	776	12.0	13	34,059	180	34,238	14,269	14,299	225.8
Indianapolis–Carmel–Anderson, IN	129	137	7.0	34	5,274	25	5,299	2,516	2,519	129.0
Jackson, MS	85	88	15.2	5	3,466	30	3,496	1,658	1,631	282.6
Jacksonville, FL	175	185	13.0	8	8,114	78	8,192	3,677	3,667	262.7
Kansas City, MO–KS	108	114	5.5	52	5,966	16	5,982	2,745	2,739	133.2
Knoxville, TN	12	14	1.6	104	1,310	8	1,319	640	621	72.9
Lakeland–Winter Haven, FL	56	59	9.2	21	2,493	22	2,515	1,122	1,119	179.6
Lancaster, PA	15	16	3.0	85	890	22	912	403	400	75.5
Las Vegas–Henderson–Paradise, NV	180	191	9.2	22	6,393	28	6,421	3,161	3,161	155.8
Little Rock–North Little Rock–Conway, AR	51	56	7.7	28	1,842	14	1,856	894	895	123.6

Table 27. Stage 3 (AIDS), 2014 and cumulative, and persons living with diagnosed HIV infection ever classified as stage 3 (AIDS) (prevalence), year-end 2013, by metropolitan statistical area of residence—United States and Puerto Rico (cont)

MSA of residence	Diagnosis, cumulative ^a											
	Diagnosis, 2014				Adults or adolescents			Children			Prevalence of stage 3 (AIDS), year-end 2013	
	No.	Estimated ^b		Rank ^d	No.	Estimated ^b		No.	Estimated ^b		Rate ^c	
		No.	Rate ^c			No.	No.		No.	No.		Rate ^c
Los Angeles–Long Beach–Anaheim, CA	891	1,000	7.5	29	71,351	307	71,658	30,938	31,264	237.3		
Anaheim Division	122	137	4.4	—	8,421	45	8,467	3,931	3,941	126.2		
Los Angeles Division	769	863	8.5	—	62,929	262	63,191	27,007	27,323	271.8		
Louisville/Jefferson County, KY–IN	94	102	8.0	26	3,217	30	3,247	1,524	1,516	120.1		
Madison, WI	13	14	2.1	100	674	5	679	315	315	50.2		
McAllen–Edinburg–Mission, TX	50	52	6.2	42	1,041	13	1,054	598	595	72.7		
Memphis, TN–MS–AR	253	279	20.8	2	7,349	46	7,395	3,461	3,359	250.3		
Miami–Fort Lauderdale–West Palm Beach, FL	1,019	1,084	18.3	3	68,640	1,011	69,651	28,564	28,520	486.4		
Fort Lauderdale Division	333	354	19.0	—	21,045	266	21,311	9,284	9,270	502.3		
Miami Division	497	528	19.8	—	35,874	516	36,391	14,356	14,336	542.6		
West Palm Beach Division	189	201	14.4	—	11,720	228	11,948	4,924	4,915	357.1		
Milwaukee–Waukesha–West Allis, WI	57	60	3.8	74	3,227	20	3,247	1,498	1,498	95.4		
Minneapolis–St. Paul–Bloomington, MN–WI	120	126	3.6	80	5,773	24	5,797	2,930	2,935	84.8		
Modesto, CA	12	13	2.4	97	848	6	854	389	387	73.5		
Nashville–Davidson–Murfreesboro–Franklin, TN	98	109	6.1	44	5,133	24	5,157	2,585	2,508	142.6		
New Haven–Milford, CT	48	52	6.0	45	5,209	75	5,284	2,183	2,169	251.3		
New Orleans–Metairie, LA	204	213	17.0	4	11,129	80	11,209	4,411	4,405	354.7		
New York–Newark–Jersey City, NY–NJ–PA ^e	1,926	2,156	10.7	17	230,132	3,028	233,160	85,124	85,927	429.6		
Nassau County Division	114	137	4.8	—	9,606	119	9,725	3,633	3,748	131.1		
New York Division	1,541	1,731	12.1	—	194,255	2,527	196,782	72,924	73,617	516.8		
Newark Division	260	274	10.9	—	24,201	369	24,569	7,811	7,787	311.3		
North Port–Sarasota–Bradenton, FL	41	43	5.8	48	2,445	29	2,474	1,052	1,049	143.2		
Ogden–Clearfield, UT	10	11	1.7	103	328	4	332	170	170	27.4		
Oklahoma City, OK	77	84	6.3	41	2,983	7	2,990	1,253	1,253	94.9		
Omaha–Council Bluffs, NE–IA	32	33	3.7	77	1,387	5	1,392	696	697	77.8		
Orlando–Kissimmee–Sanford, FL	317	336	14.5	6	11,723	95	11,818	5,596	5,584	245.9		
Oxnard–Thousand Oaks–Ventura, CA	20	21	2.5	95	1,250	5	1,255	550	548	65.1		
Palm Bay–Melbourne–Titusville, FL	27	29	5.1	57	1,831	11	1,842	807	801	145.3		

Table 27. Stage 3 (AIDS), 2014 and cumulative, and persons living with diagnosed HIV infection ever classified as stage 3 (AIDS) (prevalence), year-end 2013, by metropolitan statistical area of residence—United States and Puerto Rico (cont)

MSA of residence	Diagnosis, 2014				Diagnosis, cumulative ^a			Prevalence of stage 3 (AIDS), year-end 2013		
	Estimated ^b				Adults or adolescents	Children	Total	Estimated ^b		
	No.	No.	Rate ^c	Rank ^d	No.	No.	No.	No.	No.	Rate ^c
Philadelphia–Camden–Wilmington, PA–NJ–DE–MD	480	510	8.4	25	34,858	320	35,178	15,378	15,263	252.9
Camden Division	72	76	6.1	—	3,880	43	3,923	1,568	1,566	125.1
Montgomery County Division	39	42	2.1	—	2,851	12	2,863	1,207	1,199	61.5
Philadelphia Division	312	330	15.5	—	24,429	242	24,671	11,054	10,944	516.7
Wilmington Division	57	62	8.6	—	3,698	23	3,722	1,549	1,554	216.9
Phoenix–Mesa–Scottsdale, AZ	233	246	5.5	53	10,027	31	10,058	4,760	4,765	108.2
Pittsburgh, PA	80	86	3.6	79	3,986	21	4,007	1,703	1,693	71.7
Portland–South Portland, ME	13	16	3.0	87	728	1	729	354	357	68.5
Portland–Vancouver–Hillsboro, OR–WA	94	102	4.3	67	5,971	9	5,980	2,678	2,696	116.5
Providence–Warwick, RI–MA	47	51	3.2	83	4,760	45	4,806	2,145	2,131	132.7
Provo–Orem, UT	2	2	0.4	107	151	3	154	86	86	15.3
Raleigh, NC	79	82	6.6	38	3,000	16	3,016	1,661	1,653	136.0
Richmond, VA	75	92	7.3	30	4,469	34	4,503	1,949	1,926	154.4
Riverside–San Bernardino–Ontario, CA	216	245	5.5	51	10,993	63	11,056	5,209	5,219	118.9
Rochester, NY	39	47	4.3	70	3,801	17	3,819	1,681	1,736	160.1
Sacramento–Roseville–Arden–Arcade, CA	56	61	2.7	89	4,982	29	5,011	2,200	2,195	99.0
St. Louis, MO–IL	144	153	5.4	54	7,661	44	7,705	3,475	3,472	123.9
Salt Lake City, UT	24	26	2.2	99	2,000	9	2,009	1,061	1,063	93.1
San Antonio–New Braunfels, TX	149	155	6.7	36	6,755	32	6,788	3,103	3,073	134.6
San Diego–Carlsbad, CA	199	224	6.9	35	15,605	69	15,674	7,308	7,324	227.3
San Francisco–Oakland–Hayward, CA ^e	291	324	7.0	32	46,092	102	46,194	16,251	16,282	359.5
Oakland Division	135	153	5.6	—	12,027	53	12,080	5,131	5,144	192.0
San Francisco Division	146	160	9.9	—	32,352	44	32,396	10,512	10,539	662.2
San Jose–Sunnyvale–Santa Clara, CA	68	74	3.8	76	4,787	17	4,805	2,278	2,274	117.9
San Juan–Carolina–Caguas, PR	203	266	11.9	14	24,915	277	25,192	7,522	7,513	330.7
Santa Rosa, CA	12	13	2.6	94	2,161	8	2,169	872	866	174.9
Scranton–Wilkes-Barre–Hazleton, PA	13	14	2.5	96	708	7	715	320	317	56.5
Seattle–Tacoma–Bellevue, WA	152	164	4.5	66	11,038	28	11,066	4,921	4,963	137.3
Seattle Division	132	143	5.0	—	9,753	18	9,771	4,363	4,400	157.5
Tacoma Division	20	21	2.6	—	1,285	10	1,295	558	563	68.6
Spokane–Spokane Valley, WA	4	5	0.9	106	655	1	656	273	275	51.4

Table 27. Stage 3 (AIDS), 2014 and cumulative, and persons living with diagnosed HIV infection ever classified as stage 3 (AIDS) (prevalence), year-end 2013, by metropolitan statistical area of residence—United States and Puerto Rico (cont)

MSA of residence	Diagnosis, cumulative ^a									
	Diagnosis, 2014				Adults or adolescents			Prevalence of stage 3 (AIDS), year-end 2013		
	Estimated ^b				Estimated ^b			Estimated ^b		
	No.	No.	Rate ^c	Rank ^d	No.	No.	No.	No.	No.	Rate ^c
Springfield, MA	15	17	2.7	91	2,442	28	2,470	1,007	993	158.0
Stockton–Lodi, CA	30	33	4.6	64	1,461	14	1,475	706	704	99.9
Syracuse, NY	31	37	5.6	50	1,613	10	1,623	679	703	106.0
Tampa–St. Petersburg–Clearwater, FL	337	355	12.2	12	14,461	116	14,577	6,216	6,201	215.7
Toledo, OH	10	11	1.8	101	1,038	15	1,053	483	488	80.3
Tucson, AZ	43	46	4.6	65	2,485	11	2,496	1,070	1,069	107.1
Tulsa, OK	47	51	5.3	56	1,973	10	1,983	845	843	87.6
Virginia Beach–Norfolk–Newport News, VA–NC	141	175	10.2	18	6,157	66	6,223	2,581	2,534	148.4
Washington–Arlington–Alexandria, DC–VA–MD–WV	587	683	11.3	15	40,178	336	40,514	18,450	18,592	311.6
Silver Spring Division	97	107	8.4	—	4,029	27	4,056	2,231	2,260	179.2
Washington Division	490	576	12.1	—	36,149	309	36,458	16,219	16,332	347.1
Wichita, KS	16	17	2.7	90	1,033	2	1,035	447	445	69.7
Winston-Salem, NC	18	19	2.9	88	1,307	10	1,317	639	636	97.6
Worcester, MA–CT	11	12	1.3	105	2,463	22	2,485	1,162	1,149	123.8
Youngstown–Warren–Boardman, OH–PA	22	24	4.3	68	734	0	734	317	320	57.5
Subtotal for MSAs (population of ≥500,000)	15,429	16,993	7.8	—	1,046,393	8,663	1,055,056	441,278	442,547	205.0
Metropolitan areas (population of 50,000–499,999)	2,419	2,653	4.6	—	119,268	836	120,104	52,744	52,751	91.1
Nonmetropolitan areas	1,374	1,500	3.2	—	64,254	441	64,695	28,808	28,787	62.1
Total^f	19,350	21,300	6.6	—	1,236,088	9,992	1,246,080	525,796	527,069	164.7

Abbreviation: MSA, metropolitan statistical area.

Note. Because of the lack of U.S. census information for all U.S. dependent areas, table includes data for only the United States and Puerto Rico.

MSA definitions for this report can be found at <http://www.census.gov/population/metro/>.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a From the beginning of the epidemic through 2014.

^b Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting.

^c Rates are per 100,000 population.

^d Based on estimated rate.

^e Counts of stage 3 (AIDS) classifications for the metropolitan divisions do not sum to the MSA total. MSA total includes data from 1 metropolitan division with population of <500,000.

^f Includes persons whose county of residence is unknown. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Web Addresses for Reports of State and Local HIV Surveillance

Alabama	http://www.adph.org/aids/Default.asp?id=984
Alaska	http://www.epi.hss.state.ak.us/bulletins/catlist.jsp?cattype=HIV+(Human+Immunodeficiency+Virus)
Arizona	http://www.azdhs.gov/phs/edc/odis/hiv-epidemiology/reports/index.php?pg=annual
Arkansas	http://www.healthy.arkansas.gov/programsServices/healthStatistics/Pages/stdHIVRegistry.aspx
California	http://www.cdph.ca.gov/data/statistics/Pages/OAHIVAIDSStatistics.aspx
Chicago	http://www.cityofchicago.org/city/en/depts/cdp/provdrs/sti_hiv_aids.html
Colorado	http://www.colorado.gov/pacific/cdphe/sti-and-hiv-data-and-trends
Connecticut	http://www.ct.gov/dph/cwp/view.asp?a=3135&q=393048
Delaware	http://www.dhss.delaware.gov/dhss/dph/epi/disstatshiv.html
District of Columbia	http://doh.dc.gov/service/hiv-aids-reports-and-publications
Florida	http://www.floridahealth.gov/diseases-and-conditions/aids/surveillance/index.html
Georgia	http://dph.georgia.gov/georgias-hiv-aids-epidemiology-surveillance-section
Guam	http://dphss.guam.gov/content/hiv-surveillance-program
Hawaii	http://health.hawaii.gov/std-aids/hiv-aids/data-and-statistics/
Houston	http://www.houstontx.gov/health/HIV-STD/
Idaho	http://healthandwelfare.idaho.gov/Health/FamilyPlanningSTDHIV/STDStatistics/tabid/393/Default.aspx
Illinois	http://www.idph.state.il.us/aids/stats.htm
Indiana	http://www.in.gov/isdh/23266.htm
Iowa	http://idph.iowa.gov/PublicHealthData/reports
Kansas	http://www.kdheks.gov/sti_hiv/epi_profiles.htm
Kentucky	http://chfs.ky.gov/dph/epi/hiv-aids/surveillance.htm
Los Angeles	http://publichealth.lacounty.gov/dhsp/Reports.htm
Louisiana	http://new.dhh.louisiana.gov/index.cfm/newsroom/category/63
Maine	http://www.maine.gov/dhhs/mecdc/infectious-disease/hiv-std/data/hiv.shtml
Maryland	http://phpa.dhmm.maryland.gov/OIDEOR/CHSE/SitePages/statistics.aspx
Massachusetts	http://www.mass.gov/eohhs/gov/departments/dph/programs/id/hiv-aids/epi-profile/
Michigan	http://michigan.gov/mdch/0,4612,7-132-2940_2955_2982_46000_46003--,00.html
Minnesota	http://www.health.state.mn.us/divs/idepc/diseases/hiv/hivstatistics.html
Mississippi	http://www.msdc.state.ms.us/msdcsite/index.cfm/14,0,150,134.html
Missouri	http://health.mo.gov/data/hivstdaids/data.php
Montana	http://dphhs.mt.gov/publichealth/hivstd
Nebraska	http://dhhs.ne.gov/publichealth/Pages/dpc_hiv.aspx
Nevada	http://dphh.nv.gov/Programs/HIV-OPHIE/dta/Publications/HIV/AIDS_Surveillance_Program_(HIV-OPHIE)_-_Publications/
New Hampshire	http://www.dhhs.nh.gov/data/index.htm
New Jersey	http://www.state.nj.us/health/aids/aidsqtr.shtml
New Mexico	http://nmhealth.org/data/infectious/20/
New York	http://www.health.state.ny.us/diseases/aids/statistics/index.htm
New York City	http://www.nyc.gov/html/doh/html/living/std-hiv.shtml
North Carolina	http://epi.publichealth.nc.gov/cd/stds/figures.html
North Dakota	http://www.ndhealth.gov/hiv/
Ohio	http://www.odh.ohio.gov/healthstats/disease/hivdata/hivcov.aspx
Oklahoma	http://www.ok.gov/health/Disease,_Prevention,_Preparedness/HIV_STD_Service/HIV_STD_Statistics/index.html
Oregon	http://public.health.oregon.gov/DiseasesConditions/CommunicableDisease/DiseaseSurveillanceData/HIVData/Pages/index.aspx
Pennsylvania	http://www.portal.state.pa.us/portal/server.pt?open=514&objID=557343&mode=2
Philadelphia	http://www.phila.gov/health/AACO/AACODataResearch.html
Puerto Rico	http://www.salud.gov.pr/Estadisticas-Registros-y-Publicaciones/Pages/VIH.aspx
Rhode Island	http://www.health.ri.gov/programs/hivaidssurveillance/
San Francisco	http://www.sfdph.org/dph/files/reports/default.asp
South Carolina	http://www.dhec.sc.gov/health/disease/stdhiv/surveillance.htm
South Dakota	http://doh.sd.gov/diseases/infectious/HIV-AIDS/
Tennessee	http://tn.gov/health/topic/STD-surveillance
Texas	http://www.dshs.state.tx.us/hivstd/reports/default.shtm
U.S. Virgin Islands	http://www.healthvi.com/programs/communicable-disease/index.html
Utah	http://health.utah.gov/epi/diseases/hiv-aids/surveillance/index.html
Vermont	http://healthvermont.gov/prevent/aids/quarter/reports.aspx
Virginia	http://www.vdh.virginia.gov/epidemiology/DiseasePrevention/DAta/
Washington	http://www.doh.wa.gov/DataandStatisticalReports/DiseasesandChronicConditions/HIVAIDSData.aspx
West Virginia	http://www.dhhr.wv.gov/oeps/std-hiv-hep/HIV_AIDS/Pages/HIVSurveillance.aspx
Wisconsin	http://dhs.wisconsin.gov/aids-hiv/Stats/index.htm
Wyoming	http://www.health.wyo.gov/PHSD/hivsurveillance/index.html

Note. Electronic reports are not available for the following areas: American Samoa, the Northern Mariana Islands, and the Republic of Palau.