

UCSF Clinical Staff Evaluation Questionnaire

Resize font:  | 

Form Approved
OMB No: 0920-XXXX
Exp. Date: XXXX-XXXX-XXXX

Public Reporting burden of this collection of information is estimated to average 5 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Reports Clearance Officer, 1600 Clifton Road NW, MS D-74, Atlanta, GA 30333; Attn: PRA (0920-XXXX). Thank you for participating in this important study. Your responses to this questionnaire will be kept confidential and will not be shared outside our research and quality-improvement team.

This survey asks about your impressions of the usefulness of recommendations for post-hospital management of pain and mobility for older patients found to be at increased risk for falls contained in UCSF medical center discharge summaries. This should take no more than 5 minutes to complete. If you have questions, please call Dr. Andrew Auerbach (415-502-1414) or email Steadl.Study@ucsf.edu

Tell us about yourself and your background

1) Age < 30 30 - 39 40 - 49 50 - 64 65+ reset

2) Gender Female Male Decline to say reset

3) Years in clinical practice 0-5 6-10 11-15 16-20 >20 reset

4) Professional preparation/role MD/DO NP/PA RPH/PharmD RN reset

5) I am a student, resident, or fellow Yes No reset

6) Have a DEA registration number Yes No Not Applicable reset

7) Unit where you primarily care for patients

This section asks about your recent experiences in caring for older adult (age 65+) patients

In the past week....

	0	1-3	4-6	7-10	>10	
8) ... how many times did you screen a patient for possible opioid use disorder?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	reset
9) ...how many times did you seek consultation to aid in pain management for your patient(s)?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	reset
10) ... how many times did you give instructions for how to use non-pharmacologic pain treatments during hospitalization?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	reset
11) ... how many discharge opioid prescriptions did you write?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	reset
12) ... how many times did you give instructions on how to taper opioids after discharge?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	reset
13) ... how many times did you give instructions on non-pharmacologic pain treatments at discharge?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	reset

Please indicate your level of agreement with the following statements

	Strongly Agree	Agree	Neutral	Disagree	Strongly Disagree	
14) I feel confident in my ability to manage pain in hospitalized patients.	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	reset
15) I feel confident in my ability to prescribe opioids in the hospital	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	reset
16) I feel confident in my ability to use opioid conversion calculations in caring for my patients	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	reset
17) I feel confident in my ability to identify patients at risk for opioid misuse at the time they are admitted to the hospital	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	reset
18) I feel confident in my knowledge about how to avoid use of parenteral opioids for pain in hospitalized patients	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	reset
19) I feel confident in my ability to transition patients from opioid-inclusive to non-opioid regimens during hospitalization	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	reset
20) I feel confident in my knowledge about alternatives to opioids for pain in hospitalized patients	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	reset
21) I know where to find educational or clinical guideline resources to assist me with managing my patients' pain	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	reset
22) I feel confident in my ability to instruct patients on how to taper or transition off opioid pain medications at the time of discharge	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	reset
23) I feel confident in my ability to identify patients at risk for opioid misuse at the time they are being discharged from the hospital	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	reset
24) I feel confident in my knowledge of non-pharmacologic approaches to aid with pain management in patients after discharge	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	reset
25) Encouraging non-opioid medications for management of pain is time consuming	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	reset
26) I feel confident in my ability to counsel patients who are reluctant to avoid or reduce opioids	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	reset
27) Our electronic health record helps me choose non-opioid pain regimens during hospitalization	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	reset
28) Our electronic health record helps me choose non-opioid pain regimens at discharge	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	reset
29) Our electronic health record makes it easier for me to teach patients about how to manage their pain after discharge	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	reset
30) Our electronic health record makes it easier for me to teach patients about how to taper off opioids after discharge	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	reset

31) What suggestions do you have for how UCSF Health might help you take care of your patients with pain, particularly patients >65?

Expand

Submit