Form Approved OMB No. 0920-1156 Exp. Date xx/xx/xx

**Health Center Provider Survey** 

Public reporting of this collection of information is estimated to average 20 minutes per response, including the time for reviewing instructions and completing and reviewing the collection of information. An agency many not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a current valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Reports Clearance Officer, 1600 Clifton Road NE, MS D-74, Atlanta, Georgia 30333; ATTN: PRA (0920-1156)

Thank you for agreeing to complete this survey. The purpose of the survey is to assess your attitudes and practices related to adolescent sexual and reproductive health. Please answer each of the following questions as they relate to your experiences providing family planning services to your adolescent patients. Please answer questions thinking only about services you have provided at your current health center.

Your responses will be kept private and your completed assessment will be stored on a secure, password protected server. Your responses will be combined with those from others at your health center in order to tailor training and technical assistance. Your individual responses will not be shared. Your participation in this survey is voluntary and you can discontinue participation at any time.

P	ROVIDE	R AND PATIENT CHARA	CTERISTICS				
	1.	On average, approximate(provide a whole		nduplicated <b>fe</b> n	nale patients ag	<b>ges 15-19</b> do you	ı see per week?
	2.	Approximately what perchealth services? 0-24% 25-49% 50-74% 75% or more	ent of your <b>fem</b>	ale patients aç	<b>jes 15-19</b> do yo	u provide sexual	and reproductive
TRA	NING						
3.	Have yo	ou been formally trained in	the insertion ar	nd removal of th	e following cont	raceptive method	ds?
						Y	es No
	Сорг	per Intrauterine Device (Cu	u-IUD or ParaG	ard®)?			
	Levo	norgestrel-releasing Intra	uterine Device (	LNG-IUD LILE	ΓΤΑ™, Skyla® α	or Mirena®)?	
	Cont	raceptive implant (Impland	on®)?				
		AND PRACTICES  mfortable are you recomm	ending IUDs fo	r the following g	roups of adoles	cents (ages 15 to	o 19 years old)?
			Not at all comfortable	Slightly comfortable	Somewhat comfortable	Moderately comfortable	Very comfortable
	Nullipa	rous adolescents?					
	minutes	artum adolescents (10 s after delivery of placenta than 4 weeks postpartum)?					
	Obese kg/m²)′	adolescents (BMI ≥30 ?					

5.	In the past	year, have you referred, prescribed or provided IUDs to nulliparous adolescents?
		☐ Yes ☐ No
	If no, pleas	e indicate why (Select all that apply):
		My practice/health center does not provide IUDs.
		I do not have nulliparous adolescents as patients.
		I am concerned about recommending IUDs for adolescents.
		I am concerned about the effects on future fertility.
		I am concerned about difficult insertion.
		I am not trained in IUD insertion.
		My nulliparous adolescent patients generally prefer a different method.
		Adolescents are more likely to have it removed within a year of insertion, compared to adults
		My practice/health center protocol does not allow it.
		There are financial-related issues with billing, coding, and reimbursement processes that make the process very difficult.
		Providing adolescents with IUDs makes them less likely to use a condom
		I do not feel comfortable inserting IUDs.
		I do not generally provide IUDs to adolescents for other reasons (please specify):
6.	In the past $\Box$	year, have you referred, prescribed or provided <u>hormonal implants</u> to nulliparous adolescents?  Yes   No
	If no, pleas	e indicate why (select all that apply):
		My practice does not provide implants.
		I rarely have nulliparous adolescents as patients.
		I am concerned about recommending implants for adolescents.
		I am concerned about the effects on future fertility.
		I am concerned about difficult insertion.
		I am not trained in implant insertion.
		My nulliparous adolescent patients generally prefer a different method.
		Adolescents are more likely to have it removed within a year of insertion, compared to adults
		Providing adolescents with hormonal implants makes them less likely to use a condom
		My practice/health center protocol does not allow it.
		There are financial-related issues with billing, coding, and reimbursement processes that make the process very difficult.
		I do not feel comfortable inserting implants.

7.	Before providing the following contraceptive methods, please indicate if you perform any of the following exams
	and tests for an otherwise healthy female adolescent.

Contraceptive Method	Blood pressure	Clinical breast exam	Bimanual exam and cervical inspection	Cervical cytology	Chlamydia/ gonorrhea screening	Do not provide this method
COCs/patch/ring	YES NO	YES NO	YES	YES NO	YES NO	
Progestin-only pills (POPs)	YES NO	YES NO	YES NO	YES NO	YES	
DMPA (Depo- Provera®)	YES NO	YES NO	YES NO	YES NO	YES NO	
Contraceptive implant (Implanon®)	YES NO	YES NO	YES NO	YES NO	YES NO	
Copper Intrauterine Device (Cu-IUD or ParaGard®)	YES NO	YES NO	YES NO	YES NO	YES NO	
LEVONORGESTREL- RELEASING INTRAUTERINE DEVICE (LNG-IUD LILETTA™, SKYLA® OR MIRENA®)	YES NO	YES NO	YES NO	YES NO	YES NO	

	Never	Rarely	Sometimes	Often	Always
Provided or prescribed the pill, patch, ring or Depo Provera on the day of her visit regardless of the timing of her menses (Quick Start)					
Provided a hormonal implant on the day of her visit regardless of the timing of her menses ( <b>Quick Start</b> ) if you were reasonably certain she was not pregnant					
Provided an IUD on the day of her visit regardless of the timing of her menses ( <b>Quick Start</b> ) if you were reasonably certain she was not pregnant					
Provided or prescribed a contraceptive method at the same time you provided EC					
Provided contraceptive services to an adolescent that came in for a pregnancy test that was negative					
Provided a Cu-IUD as EC					
Provided another method of birth control if selected method not available on day of visit					
Dispensed a year's supply of pills at one visit for adolescent females					
Addressed contraceptive needs on the day of service regardless of chief complaint rather than scheduling a follow-up visit					

	Never	Rarely	Sometimes	Often	Alway
Assessed the patient's reproductive life plan (i.e., asked about their intentions regarding the number and timing of pregnancies in the context of their personal values and life goals)					
Worked with the client interactively to select the most effective method that meets the client's needs and preferences					
Presented information regarding potential contraceptive methods with the most effective methods presented first (tiered approach)					
Helped the patient think about potential barriers to using their selected method correctly and develop a plan to deal with these barriers					
Used a method-specific informed consent form					
Informed adolescents that long-acting reversible contraceptives are safe and effective options					
Actively encouraged communication between adolescents and parents/guardians about sex and reproductive health					
Encouraged male/partner involvement in contraceptive and reproductive health services, if appropriate					

## SERVICES TO ADOLESCENTS NOT INITIALLY SEEKING SEXUAL OR REPRODUCTIVE HEALTH CARE

pur	e following items are about services pose other than sexual or reproduct			•	were seekir	ng health care
	Check here if you only see family pla	anning clients	. You have o	completed the s	survey. Thar	nk you.
	he past month, when your <b>female</b> a					-
-	pose other than sexual or reproduct the following?	ive neaith cai	re (e.g., sick	visit), now freqi	uentiy did yo	u or your clinic
		Never	Rarely	Sometimes	Often	Always
	Provide time alone					
	Screen to determine if in need of contraceptive services (i.e., sexually active, not desiring pregnancy)					
	Provide time alone	Never	Rarely	Sometimes	Often	Always
	Provide time alone Screen to determine if in need	Never	Rarely	Sometimes	Often	Always
	Provide time alone Screen to determine if in need of counseling around preventing pregnancy					
	Screen to determine if in need of counseling around preventing					