

Intensive Didactic Post-Course Evaluation

Public reporting burden of this collection of information is estimated to average 10 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Reports Clearance Officer; 1600 Clifton Road NE, MS D-74, Atlanta, Georgia 30333; ATTN: PRA (0920-0995).

1. How satisfied were you with your overall learning experience?

very unsatisfied very satisfied

2. How satisfied were you with the quality of the content?

very unsatisfied very satisfied

3. How satisfied were you with the teaching methods?

very unsatisfied very satisfied

4. How satisfied were you with the trainer(s)?

very unsatisfied very satisfied

5. Was this training free of commercial bias?

- Yes
 No

6. Was this training evidence-based?

- Yes
 No

7a. Were the learning objectives met?

- Yes
 No

7b. If the learning objectives were not met, please explain.

8. What could improve this training?

9a. As a result of information presented, do you intend to make changes in your practice or at your worksite setting?

- Yes
 No
 Not my job
 I already use these practices
 Other reason (please specify)

9b. If yes, please list at least one intended change

10a. How confident were you in your ability to describe how chlamydia infections present clinically, BEFORE the training?

not at all confident very confident

10b. How confident are you AFTER the training?

not at all confident very confident

11a. How confident were you in your ability to describe the current CDC screening recommendations for chlamydia, including extra-genital screening BEFORE the training?

not at all confident very confident

11b. How confident are you AFTER the training?

not at all confident very confident

12a. How confident were you in your ability to obtain or instruct patients to collect the recommended specimens from indicated sites to diagnose chlamydia BEFORE the training?

not at all confident very confident NA

12b. How confident are you AFTER the training?

not at all confident very confident NA

13a. How confident were you in your ability to treat patients diagnosed with chlamydia and related anogenital syndromes based on the most current CDC treatment recommendations BEFORE the training?

not at all confident very confident NA

13b. How confident are you AFTER the training?

not at all confident very confident NA

14. What is the CDC recommended regimen for treating asymptomatic uncomplicated chlamydia infection of the cervix, urethra, or rectum?

- Acyclovir 1 g twice a day for 7 days
- Azithromycin 1 g orally in a single dose or doxycycline 100 mg twice a day for 7 days
- Azithromycin 1 g orally in a single dose plus ceftriaxone 250 mg intramuscularly in a single dose
- Ciprofloxacin 500 mg orally in a single dose

15. What is the recommended follow-up for a non-pregnant patient after diagnosis and treatment of chlamydia?

- A test of cure at 2 weeks, and repeat test at 3 months
- A test of cure at 2 weeks, and repeat test at 12 months
- Repeat test in 3 months
- Repeat test in 12 months

16a. Approximately what % of sexually active asymptomatic female patients under age 25 did you screen annually for chlamydia BEFORE this training?

0% 1-25% 26-50% 51-75% 76-90% >91% NA

16b. What % do you intend to screen AFTER the training?

0% 1-25% 26-50% 51-75% 76-90% >91% NA

17a. Approximately what % of pregnant patients under age 25 did you screen for chlamydia BEFORE this training?

0% 1-25% 26-50% 51-75% 76-90% >91% NA

17b. What % do you intend to screen AFTER the training?

0% 1-25% 26-50% 51-75% 76-90% >91% NA

18a. How confident were you in your ability to describe how gonorrhea infections present clinically BEFORE this training?

not at all confident very confident

18b. How confident are you AFTER the training?

not at all confident very confident

19a. How confident were you in your ability to describe the current CDC screening recommendations for gonorrhea, including extra-genital screening BEFORE this training?

not at all confident very confident

19b. How confident are you AFTER the training?

not at all confident very confident

20a. How confident were you in your ability to obtain or direct patients to collect the recommended specimens from indicated sites to diagnose gonorrhea BEFORE this training?

not at all confident very confident NA

20b. How confident are you AFTER the training?

not at all confident very confident NA

21a. How confident were you in your ability to treat patients with gonorrhea according to current CDC recommendations in light of antibiotic resistance in *N. gonorrhoeae* BEFORE this training?

not at all confident very confident NA

21b. How confident are you AFTER the training?

not at all confident very confident NA

22. What is the recommended treatment for a patient diagnosed with uncomplicated urethral, cervical, or rectal gonorrhea?

- Ceftriaxone 250 mg intramuscularly only
- Azithromycin 2 g orally in a single dose only
- Ceftriaxone 250 mg intramuscularly plus azithromycin 1 g orally in a single dose
- Cefixime 400 mg orally plus doxycycline 100 mg orally BID for 7 days

23. What is the recommended follow-up for a non-pregnant patient after diagnosis and treatment of gonorrhea?

- A test of cure at 2 weeks, and repeat test at 3 months
- A test of cure at 2 weeks, and repeat test at 12 months
- Repeat test in 3 months
- Repeat test in 12 months

24a. Approximately what % of sexually active asymptomatic female patients under 25 did you screen annually for gonorrhea BEFORE this training?

0% 1-25% 26-50% 51-75% 76-90% >91% NA

24b. What % do you intend to screen AFTER the training?

0% 1-25% 26-50% 51-75% 76-90% >91% NA

25a. Approximately what % of pregnant patients under age 25 did you screen for gonorrhea BEFORE this training?

0% 1-25% 26-50% 51-75% 76-90% >91% NA

25b. What % do you intend to screen AFTER the training?

0% 1-25% 26-50% 51-75% 76-90% >91% NA

26a. Approximately what % of sexually active male patients who have sex with men did you screen annually for urogenital and extragenital gonorrhea and chlamydia BEFORE this training?

0% 1-25% 26-50% 51-75% 76-90% >91% NA

26b. What % do you intend to screen AFTER the training?

- 0% 1-25% 26-50% 51-75% 76-90% >91% NA

27. As a result of the information presented do you intend to provide Expedited Partner Therapy (EPT) to heterosexual partners of those diagnosed with gonorrhea and/or chlamydia?

- Yes
 No
 Not applicable to my practice or job
 Not allowed in my state practice
 My practice/worksites is in the planning stages to offer EPT
 My practice/worksites already offers EPT
 EPT was not discussed

28a. How confident were you in your ability to identify the stages of syphilis based on their clinical manifestations BEFORE this training?

- not at all confident very confident

28b. How confident are you AFTER the training?

- not at all confident very confident

29a. How confident were you in your ability to describe current CDC screening recommendations for syphilis BEFORE this training?

- not at all confident very confident

29b. How confident are you AFTER the training?

- not at all confident very confident

30a. How confident were you in your ability to order and interpret the CDC recommended serologic tests to diagnose syphilis BEFORE this training?

- not at all confident very confident NA

30b. How confident are you AFTER the training?

- not at all confident very confident NA

31a. How confident were you in your ability to clinically manage patients diagnosed with syphilis based on CDC treatment, follow-up, and partner management recommendations BEFORE this training?

- not at all confident very confident NA

31b. How confident are you AFTER the training?

- not at all confident very confident NA

32. What is the recommended treatment for primary and secondary syphilis in adults who are *not* HIV+ or pregnant?

- Acyclovir 1 g twice a day for 7 days
 Azithromycin 1 g orally in a single dose
 Benzathine penicillin G 2.4 million units IM in a single dose
 Benzathine-procaine penicillin 2.4 million units in a single dose

33a. Approximately what % of your male patients who have sex with men did you screen at least once a year for syphilis BEFORE this training?

- 0% 1-25% 26-50% 51-75% 76-90% >91% NA

33b. What % do you intend to screen AFTER the training?

- 0% 1-25% 26-50% 51-75% 76-90% >91% NA

34a. Approximately what % of your pregnant patients did you screen for syphilis BEFORE this training?

0% 1-25% 26-50% 51-75% 76-90% >91% NA

34b. What % do you intend to screen AFTER the training?

0% 1-25% 26-50% 51-75% 76-90% >91% NA

35a. Approximately what % of your patients recently diagnosed with syphilis did you test for HIV BEFORE this training?

0% 1-25% 26-50% 51-75% 76-90% >91% NA

35b. What % do you intend to screen AFTER the training?

0% 1-25% 26-50% 51-75% 76-90% >91% NA

36a. How confident were you in your ability to describe the pathogenesis of genital HPV and identify it clinically BEFORE this training?

not at all confident very confident

36b. How confident are you AFTER the training?

not at all confident very confident

37a. How confident were you in your ability to describe the national cervical cancer screening recommendations, including HPV testing, BEFORE this training?

not at all confident very confident

37b. How confident are you AFTER the training?

not at all confident very confident

38a. How confident were you in your ability to treat genital warts BEFORE the training?

not at all confident very confident NA

38b. How confident are you AFTER the training?

not at all confident very confident NA

39a. How confident were you in your ability to describe CDC/ACIP recommendations for HPV vaccination BEFORE this training?

not at all confident very confident

39b. How confident are you AFTER the training?

not at all confident very confident

40a. With approximately what % of your non-vaccinated female patients ages 11 through 26 did you discuss HPV vaccination during a preventive health visit BEFORE this training?

0% 1-25% 26-50% 51-75% 76-90% >91% NA

40b. With what % do you intend to discuss this AFTER the training?

0% 1-25% 26-50% 51-75% 76-90% >91% NA

41a. With approximately what % of your non-vaccinated male patients ages 11 through 21 did you discuss HPV vaccination during a preventive health visit BEFORE this training?

0% 1-25% 26-50% 51-75% 76-90% >91% NA

41b. With what % do you intend to discuss this AFTER the training?

0% 1-25% 26-50% 51-75% 76-90% >91% NA

42a. With approximately what % of your non-vaccinated male patients age 26 or under who have sex with men or are HIV+ did you discuss HPV vaccination during a preventive health visit BEFORE this training?

0% 1-25% 26-50% 51-75% 76-90% >91% NA

42b. With what % do you intend to discuss this AFTER the training?

0% 1-25% 26-50% 51-75% 76-90% >91% NA

43a. How confident were you in your ability to describe the major components (5 P's) of a routine sexual history BEFORE this training?

not at all confident very confident

43b. How confident are you AFTER the training?

not at all confident very confident

44a. How confident were you in your ability to demonstrate respectful, non-judgmental communication skills to discuss sexual practices, sexual risks, and risk reduction strategies with patients BEFORE this training?

not at all confident very confident NA

44b. How confident are you AFTER the training?

not at all confident very confident NA

45a. For approximately what % of patients older than 15 seeing you for a preventive health visit did you take a sexual history that asks about behaviors that could put them at risk of getting or transmitting an STD BEFORE this training?

0% 1-25% 26-50% 51-75% 76-90% >91% NA

45b. For what % do you intend to take ask about risk behaviors AFTER the training?

0% 1-25% 26-50% 51-75% 76-90% >91% NA

46a. For approximately what % of patients older than 15 seeing you for a preventive health visit did you take a sexual history that asked whether they had oral, vaginal, or anal sex BEFORE this training?

0% 1-25% 26-50% 51-75% 76-90% >91% NA

46b. For what % do you intend to ask whether they had oral, vaginal, or anal sex AFTER the training?

0% 1-25% 26-50% 51-75% 76-90% >91% NA

Thank you for your time and thoughtful feedback!
If you are requesting Continuing Education credit, please click on "Submit Evaluation" and follow the directions posted on the next page.