

# Evaluation Tester

Training > Evaluations > Questions > Evaluation Tester

## STD Treatment Guidelines Complete Post-Course Evaluation

*Public reporting burden of this collection of information is estimated to average 6 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Reports Clearance Officer; 1600 Clifton Road NE, MS D-74, Atlanta, Georgia 30333; ATTN: PRA (0920-0995).*

1. How satisfied were you with your overall learning experience?

very unsatisfied      very satisfied

2. How satisfied were you with the quality of the content?

very unsatisfied      very satisfied

3. How satisfied were you with the trainer(s)?

very unsatisfied      very satisfied

4. How satisfied were you with the teaching methods?

very unsatisfied      very satisfied

5. Was this training free of commercial bias?

- Yes  
 No

6. Was this training evidence-based?

- Yes  
 No

7a. Were the learning objectives met?

- Yes  
 No

7b. If the learning objectives were not met, please explain.

8. What could improve this training?





24a. How confident were you in your ability to clinically manage patients diagnosed with syphilis based on CDC treatment, follow-up, and partner management recommendations BEFORE this training?

not at all confident      very confident  NA

24b. How confident are you AFTER the training?

not at all confident      very confident  NA

25. What is the recommended treatment for primary and secondary syphilis in adults who are *not* HIV+ or pregnant?

- Acyclovir 1 g twice a day for 7 days
- Azithromycin 1 g orally in a single dose
- Benzathine penicillin G 2.4 million units IM in a single dose
- Benzathine-procaine penicillin 2.4 million units in a single dose

26a. Approximately what % of your male patients who have sex with men did you screen at least once a year for syphilis BEFORE this training?

0%  1-25%  26-50%  51-75%  76-90%  >91%  NA

26b. What % do you intend to screen AFTER the training?

0%  1-25%  26-50%  51-75%  76-90%  >91%  NA

27a. Approximately what % of your pregnant patients did you screen for syphilis BEFORE this training?

0%  1-25%  26-50%  51-75%  76-90%  >91%  NA

27b. What % do you intend to screen AFTER the training?

0%  1-25%  26-50%  51-75%  76-90%  >91%  NA

28a. Approximately what % of your patients recently diagnosed with syphilis did you test for HIV BEFORE this training?

0%  1-25%  26-50%  51-75%  76-90%  >91%  NA

28b. What % do you intend to screen AFTER the training?

0%  1-25%  26-50%  51-75%  76-90%  >91%  NA

**Thank you for your time and thoughtful feedback!**

**If you are requesting Continuing Education credit, please click on "Submit Evaluation" and follow the directions posted on the next page.**

OMB Control Number 0920-0995

Exp. Date 10/31/2016