

# 21

## Multi-Site Clinical Assessment of Chronic Fatigue Syndrome

---

### Orthostatic Grading Scale (OGS)

Subject ID Number: \_\_\_\_\_

Start Date: \_\_\_\_\_/\_\_\_\_\_/\_\_\_\_\_ & Time: \_\_\_\_\_ am/pm  
Month Day Year HH:MM

Complete Date: \_\_\_\_\_/\_\_\_\_\_/\_\_\_\_\_ & Time: \_\_\_\_\_ am/pm  
Month Day Year HH:MM

Public reporting burden of this collection of information is estimated to average 3 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB Control Number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Reports Clearance Officer, 1600 Clifton Road NE, MS D-74, Atlanta, Georgia 30333; ATTN: PRA 0920-XXXX

# OGS

Some patients tell us they feel worse when they are standing. For them, standing can worsen fatigue, pain or malaise or produce a feeling of light headedness or faintness. In the next questionnaire, we will ask you about these orthostatic symptoms. For each of the following 5 questions check the box that best indicates the frequency, severity, and impact of your orthostatic symptoms.

## 1. Frequency of orthostatic symptoms

0		I <i>never</i> or rarely experience orthostatic symptoms when I stand up
1		I <i>sometimes</i> experience orthostatic symptoms when I stand up
2		I <i>often</i> experience orthostatic symptoms when I stand up
3		I <i>usually</i> experience orthostatic symptoms when I stand up
4		I <i>always</i> experience orthostatic symptoms when I stand up

## 2. Severity of orthostatic symptoms

0		I <i>do not</i> experience orthostatic symptoms when I stand up
1		I experience <i>mild</i> orthostatic symptoms when I stand up
2		I experience <i>moderate</i> orthostatic symptoms when I stand up and <i>sometimes</i> have to sit back down for relief
3		I experience <i>severe</i> orthostatic symptoms when I stand up and <i>frequently</i> have to sit back down for relief
4		I experience <i>severe</i> orthostatic symptoms when I stand up and <i>regularly faint</i> if I do not sit back down

## 3. Conditions under which orthostatic symptoms occur

0		I <i>never or rarely</i> experience orthostatic symptoms under any circumstances
1		I <i>sometimes</i> experience orthostatic symptoms under certain conditions, such as prolonged standing, a meal, exertion (e.g., walking) or when exposed to heat (e.g. hot day, hot batch, hot shower)
2		I <i>often</i> experience orthostatic symptoms under certain conditions, such as prolonged standing, a meal, exertion (e.g., walking), or when exposed to heat (e.g., hot day, hot bath, hot shower)
3		I <i>usually</i> experience orthostatic symptoms under certain conditions, such as prolonged standing, a meal, exertion (e.g., walking), or when exposed to heat (e.g., hot day, hot bath, hot shower)
4		I <i>always</i> experience orthostatic symptoms when I stand up; the specific conditions do not matter

4. Activities of daily living

0	My orthostatic symptoms <i>do not interfere</i> with activities of daily living (e.g. work, chores, dressing, bathing)
1	My orthostatic symptoms <i>mildly interfere</i> with activities of daily living (e.g. work, chores, dressing, bathing)
2	My orthostatic symptoms <i>moderately interfere</i> with activities of daily living (e.g. work, chores, dressing, bathing)
3	My orthostatic symptoms <i>severely interfere</i> with activities of daily living (e.g. work, chores, dressing, bathing)
4	My orthostatic symptoms <i>severely interfere</i> with activities of daily living (e.g. work, chores, dressing, bathing). <i>I am bed or wheelchair bound because of my symptoms</i>

5. Standing time

0	On most occasions, I can stand as long as necessary without experiencing orthostatic symptoms
1	On most occasions, I can stand <i>more than 15 minutes</i> before experiencing orthostatic symptoms
2	On most occasions, I can stand <i>5-14 minutes</i> before experiencing orthostatic symptoms
3	On most occasions, I can stand <i>1-4 minutes</i> before experiencing orthostatic symptoms
4	On most occasions, I can stand <i>less than 1 minute</i> before experiencing orthostatic symptoms

THIS IS THE END OF THE SURVEY