Form Approved
OMB No. XXXX-XXXX
Exp. Date xx/xx/xxxx

The course you are taking will contain presentation materials developed by the Mountain Plains Practice and Implementation Center (PIC), located at the University of Nevada, Reno. These materials are based on content developed by the Centers for Disease Control and Prevention (CDC).

This survey does not ask you for any personal identifying information and the results will only be used to assist us in determining the effectiveness of the course. The course instructor will not be able to view your individual responses or know if you completed this survey. Likewise, you are not required to complete this survey and your participation in this class will not be affected regardless of your decision.

Thank you for your time and enjoy the class.

1. Please enter the personal ID code you created for the	pre-
test survey.	
First letter of your mother's first name	
First digit of your social security number	
Last digit of your social security number	
First letter of your mother's maiden name	

The public reporting burden of this collection of information is estimated to average 10 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Reports Clearance Officer; 1600 Clifton Road NE, MS D-74, Atlanta, Georgia 30333 ATTN: PRA (XXXX-XXXX)

The following items will be used only to describe the demographic makeup of the class.
2. Are you (choose one)
Male
Female
3. Are you Hispanic or Latino/a?
○ No
Yes
4. How do you do only a very mose Q (about all that analy)
4. How do you describe your race? (check all that apply)
American Indian/Alaska Native
Asian
Black/African American
Native Hawaiian/Pacific Islander
White
5. In which state do you work?
6. In what type of medical setting do you work?
OB/GYN
OB/GYN Family medicine
Family medicine

The scenarios describe the alcohol use-related behaviors of two women. Please read each scenario carefully and respond to the questions that follow.

7. Scenario 1

Over the past few months, Sally has started to drink more than her usual amount of alcohol. In fact, she has noticed that she now needs to drink twice as much in order to get the same effect. Several times she has tried to cut down or stop drinking. However, each time she tried she became very agitated and couldn't sleep, so would start drinking and the symptoms would go away. Her family has complained that she is often hung over, moody, and has become unreliable – making plans one day and cancelling them the next. In fact, they report that she frequently calls them while intoxicated and has no memory of the calls or what she promised the next day.

In your opinion, how likely is it that Sally's situation might be caused by...

	Not at all likely	Not very likely	Neither likely nor unlikely	Somewhat likely	Very likely
her own bad character.					
a chemical imbalance in her brain.					
the way she was raised.					
stressful circumstances in her life.					
a genetic or inherited problem.					

	Not at all likely	2	3	4	5	Extremely likel
feel ashamed if people	140t at all likely		J	7	, and the second	Extremely like
knew someone like Sally was a member of your family?						
be uncomfortable having a conversation with Sally?	\bigcirc	\bigcirc		\bigcirc	\bigcirc	
be angry with Sally?						
believe that Sally's drinking is controllable?						
talk with Sally about her drinking?						
feel sorry for Sally?						
think that Sally's drinking is her own fault?						
interview Sally for a job if you were an employer?						
feel irritated with Sally?						
be likely to help Sally?						
feel sympathy for Sally?						
believe that Sally is responsible for her drinking?					\bigcirc	
rent Sally an apartment if you were a landlord?				\bigcirc		
be concerned about Sally?				\bigcirc		
try to stay away from Sally?						

9. Scenario 2

During the past year, Jessica has started to drink more than her usual amount of alcohol. In fact, she has noticed that she now needs to drink twice as much in order to get the same effect. Three months ago, Jessica found out that she is pregnant and her doctor told her she should not drink alcohol during her pregnancy. Jessica has tried but so far has not been able to stop drinking. Some of Jessica's friends told her that when they were pregnant the taste and/or smell of alcohol made them sick. Jessica was hoping that she would have the same reaction, but that has not been the case. Out of concern for Jessica's health and the health of her baby, her family has tried several times to talk with her about it. However, they are telling her that she is being selfish and not behaving in the best interest of her child. This approach is only increasing how guilty and ashamed Jessica is feeling about not being able to stop. As a result, Jessica has become very secretive about her drinking, sneaking a drink whenever she has a chance.

In your opinion, how likely is it that Jessica 's situation might be caused by...

	Not at all likely	Not very likely	Neither likely nor unlikely	Somewhat likely	Very likely
her own bad character.					
a chemical imbalance in her brain.			\bigcirc		
the way she was raised.					
stressful circumstances in her life.			\bigcirc		
a genetic or inherited problem.		\bigcirc			

0. On a scale of 1 (N	ot at all) - 6 (Ext	remely), how	v likely would y	ou be to		
	Not at all likely	2	3	4	5	Extremely likely
feel ashamed if people knew someone like Jessica was a member of your family?						
be uncomfortable having a conversation with Jessica?					\bigcirc	
be angry with Jessica?						
believe that Jessica's drinking is controllable?						
talk with Jessica about her drinking?						
feel sorry for Jessica?						
think that Jessica's drinking is her own fault?		\bigcirc				
interview Jessica for a job if you were an employer?						
feel irritated with Jessica?						
be likely to help Jessica?						
feel sympathy for Jessica?						
believe that Jessica is responsible for her drinking?						
rent Jessica an apartment if you were a landlord?						
be concerned about Jessica?			\bigcirc			
try to stay away from Jessica?						

The following items deal with factors related to fetal alcohol spectrum disorders.
11. Fetal Alcohol Spectrum Disorders (FASD) is an umbrella term describing the range of effects that can occur in an individual whose mother drank alcohol during pregnancy.
True
False
12. FASD is preventable if a woman does not drink during her pregnancy.
☐ True
False
13. The effects for FASD are always visible.
True
False
14. When is it safe to drink alcohol during pregnancy?
During the first three months
During the last three months
Once in a while
Never

15. How much alcohol is safe to drink during pregnancy?
One glass of wine
Two light beers
One shot of hard alcohol
None of the above
16. Screening for excessive alcohol use during pregnancy can be an effective strategy in reducing FASD.
True
False
17. How often do you talk to your patients or their parents/caregivers about prevention of excessive alcohol
use?
Never
Occasionally
About Half the Time
Frequently
Always
18. The practice in which I work has a protocol to screen all patients about their alcohol use.
Yes
○ No
Oon't know
Not applicable to the patients in my practice setting.

19. V	William Constitution of the Constitution of th
	Vhat does initial patient SCREENING for alcohol use consist of in your practice?
<u> </u>	nformal questions (Do you drink? How often/much do you drink?, etc.)
I	Formal screening tool or evidence-based screening instruments (AUDIT, AUDIT-C, DAST, CAGE, etc.)
	don't know.
20. V	Who generally conducts the initial SCREEN for alcohol? (Check all that apply.)
	Nurse (including nurse practitioner)
	Social worker
	Behavioral health specialist (coach)
<u> </u>	Psychologist
	Physician
	Medical Assistant
	Other (please specify)
	s screening for alcohol use followed by some type of intervention in your practice? (Check all that
apply	
apply	<i>(</i> .)
apply	No, there is no patient education or intervention done following the initial screening. Yes, all patients are given educational materials/information on "safe" levels of alcohol and health risks associated with
apply	No, there is no patient education or intervention done following the initial screening. Yes, all patients are given educational materials/information on "safe" levels of alcohol and health risks associated with consuming too much alcohol.
apply	No, there is no patient education or intervention done following the initial screening. Yes, all patients are given educational materials/information on "safe" levels of alcohol and health risks associated with consuming too much alcohol. Yes, patients who screen positive for risky alcohol use are asked follow-up questions and provided brief counseling. Yes, patients who screen positive for risky alcohol use are asked follow-up questions and provided with additional resources (e.g.
apply	No, there is no patient education or intervention done following the initial screening. Yes, all patients are given educational materials/information on "safe" levels of alcohol and health risks associated with consuming too much alcohol. Yes, patients who screen positive for risky alcohol use are asked follow-up questions and provided brief counseling. Yes, patients who screen positive for risky alcohol use are asked follow-up questions and provided with additional resources (e.g.
apply	No, there is no patient education or intervention done following the initial screening. Yes, all patients are given educational materials/information on "safe" levels of alcohol and health risks associated with consuming too much alcohol. Yes, patients who screen positive for risky alcohol use are asked follow-up questions and provided brief counseling. Yes, patients who screen positive for risky alcohol use are asked follow-up questions and provided with additional resources (e.g.
apply	No, there is no patient education or intervention done following the initial screening. Yes, all patients are given educational materials/information on "safe" levels of alcohol and health risks associated with consuming too much alcohol. Yes, patients who screen positive for risky alcohol use are asked follow-up questions and provided brief counseling. Yes, patients who screen positive for risky alcohol use are asked follow-up questions and provided with additional resources (e.g.
apply	No, there is no patient education or intervention done following the initial screening. Yes, all patients are given educational materials/information on "safe" levels of alcohol and health risks associated with consuming too much alcohol. Yes, patients who screen positive for risky alcohol use are asked follow-up questions and provided brief counseling. Yes, patients who screen positive for risky alcohol use are asked follow-up questions and provided with additional resources (e.g.
apply	No, there is no patient education or intervention done following the initial screening. Yes, all patients are given educational materials/information on "safe" levels of alcohol and health risks associated with consuming too much alcohol. Yes, patients who screen positive for risky alcohol use are asked follow-up questions and provided brief counseling. Yes, patients who screen positive for risky alcohol use are asked follow-up questions and provided with additional resources (e.g.

22. Who generally does the intervention in your practice? (Check all that apply.) No one intervenes, to my knowledge Physician Physician's Assistant Psychologist Social worker Behavior health specialist (coach) Nurse (including nurse practitioner) Other (please specify) 23. Does your practice bill for screening and brief intervention services? Yes Not sure

	0 (Completely disagree)	1	2	3	4	5	6	7	8	9	10 (Completel agree)
Alcohol consumption during pregnancy is more prevalent in women with lower incomes.											0
Alcohol consumption during pregnancy is more prevalent in women with lower levels of education.										\bigcirc	\bigcirc
Alcohol consumption during pregnancy is more prevalent in Anglo-white women.											
5. Which of the following ONE box). Occasional consum	nption of one sta	andard d	lrink alcol	holic drinl	k per day						
5. Which of the follonly ONE box).	aption of one sta	andard d	Irink alcol e mother	holic drinl or the fe	k per day tus.	or less (i	.e., 1.5 o	z. hard lid	ηuor, 12 o	oz. of bee	er or 5 oz. of
5. Which of the following ONE box). Occasional consumwine) during pregna	aption of one sta	andard d	Irink alcol e mother	holic drinl or the fe	k per day tus.	or less (i	.e., 1.5 o	z. hard lid	ηuor, 12 o	oz. of bee	er or 5 oz. of
5. Which of the following ONE box). Occasional consumwine) during pregna	aption of one sta	andard d	Irink alcol e mother	holic drinl or the fe	k per day tus.	or less (i	.e., 1.5 o	z. hard lid	ηuor, 12 o	oz. of bee	er or 5 oz. of
5. Which of the following ONE box). Occasional consumwine) during pregna	aption of one sta	andard d	Irink alcol e mother	holic drinl or the fe	k per day tus.	or less (i	.e., 1.5 o	z. hard lid	ηuor, 12 o	oz. of bee	er or 5 oz. of
5. Which of the following ONE box). Occasional consumwine) during pregna	aption of one sta	andard d	Irink alcol e mother	holic drinl or the fe	k per day tus.	or less (i	.e., 1.5 o	z. hard lid	ηuor, 12 o	oz. of bee	er or 5 oz. of

26. On a scale from 0 to 4 where 0 means you completely disagree with the statement and 4 means you completely agree, to what extent do you disagree or agree with the following statements.

	Completely disagree	Disagree	Neither agree nor disagree	Agree	Completely agree
It is important to routinely screen all patients for alcohol use	0	0	0		0
It is important to screen all pregnant women for alcohol use					
It is important to screen all women of childbearing age, for alcohol use					
I am comfortable asking women of childbearing age, including pregnant women, about their alcohol use					
I am comfortable having a conversation with patients who indicate risky alcohol use.					
Screening for alcohol use confers a negative stigma to the woman being screened.					

Educate women of childbearing age, including those who are pregnant about the effects of alcohol on a developing baby Conduct brief interventions for reducing alcohol use Utilize resources to refer patients who need
Screen women for risky or hazardous drinking Educate women of childbearing age, including those who are pregnant about the effects of alcohol on a developing baby Conduct brief interventions for reducing alcohol use Utilize resources to refer patients who need formal treatment for
childbearing age, including those who are pregnant about the effects of alcohol on a developing baby Conduct brief interventions for reducing alcohol use Utilize resources to refer patients who need
interventions for reducing alcohol use Utilize resources to refer patients who need
refer patients who need
alcohol abuse

Medical Assistants Pretest Survey
Thank you for your participation!!!