

FASD Post-Training Survey

Page 1 of 11

Form Approved
OMB No. 0920-XXXX
Exp. Date xx/xx/20xx

Thank you for completing the training. You are invited to take part in a survey for health professionals. This survey is being conducted on behalf of the Centers for Disease Control and Prevention (CDC). The purpose of this survey is to understand the opinions and practices of health professionals around their patients'/clients' alcohol use and on the prevention, identification, and treatment of fetal alcohol spectrum disorders. Your feedback is important as it will help assess the efficacy of trainings and identify the needs of health professionals to better address the services they provide to patients/clients around alcohol consumption and/or the effects of alcohol use during pregnancy.

This survey will take approximately 5 minutes to complete and your responses will be kept secure. You will be assigned a unique identity code which will be used to invite you to take a similar survey after six months to assess how useful this training has been in your practice with your patients/clients regarding their alcohol use. ICF International and Westat are the contractors hired by CDC to conduct and analyze the surveys, respectively. Any information you provide will be presented in aggregate in a report and no individual identifying information will be included.

Risks to participating in this survey are minimal and include the risk of your information becoming known to individuals outside the CDC, ICF International, and Westat. There will be no costs for participating, nor will you benefit from participating. Your participation in this survey is voluntary. You may decline to answer any question and you have the right to stop the survey at any time.

If you have any comments or questions about the survey, you can contact Melanie Chansky at melaniechansky@westat.com or at 301-517-4019.

Thank you.

CDC estimates the average public reporting burden for this collection of information as 5 minutes per response, including the time for reviewing instructions, searching existing data/information sources, gathering and maintaining the data/information needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Information Collection Review Office, 1600 Clifton Road NE, MS D-74, Atlanta, Georgia 30333; ATTN: PRA (0920-XXXX).



FASD Post-Training Survey

Page 2 of 11

1. I have an increased understanding of fetal alcohol spectrum disorders (FASDs) after the training

- Yes
- No
- Not Sure

2. Fetal alcohol spectrum disorders are: (Check only one response.)

- Disorders a pregnant woman experiences when she drinks alcohol.
- Disorders that affect the ability of a pregnant woman who drinks alcohol to go full term
- Physical disorders that affect a fetus when a pregnant woman drinks alcohol.
- The range of effects that can occur in an individual who was exposed prenatally to alcohol

3. The effects of FASDs are always visible.

- True
- False
- Don't know

4. FASDs are certain to be prevented when: (Check only one response.)

- a woman quits drinking as soon as she knows she is pregnant.
- a woman who is pregnant or may become pregnant does not consume alcohol.
- a woman does not take drugs other than alcohol during her pregnancy.
- a woman stops drinking once she starts breastfeeding her baby

5. Which of the following are the primary facial dysmorphic features associated with Fetal Alcohol Syndrome? (Check all that apply.)

- Wide inner canthal distance
- Short palpebral fissures
- Full lips
- Smooth Philtrum
- Thin upper lip
- Flaring nares
- Don't know/unsure



FASD Post-Training Survey

Page 3 of 11



6. What advice would you give your patient/client about how much alcohol is safe to drink during pregnancy? (Check only one response.)

- One glass of wine per day
- One light beer per day
- One shot of hard alcohol per day
- There is no known safe amount of alcohol consumption during pregnancy

7. When is it safe to drink alcohol during pregnancy? (Check only one response.)

- During the first three months
- During the last three months
- Once in a while
- Never

8. What is the most effective strategy to reduce fetal alcohol spectrum disorders (FASDs)? (Check only one response.)

- Enact laws and other policy strategies that punish pregnant women for drinking alcohol.
- Screen all women of child bearing age for alcohol use and provide intervention as appropriate
- Tell women who you think may have a drinking problem to get help
- Conduct health fairs and other educational events for new mothers that focus on binge drinking.



FASD Post-Training Survey

Page 4 of 11



9. Are you currently a/an (Check only one response):

- Student
- Resident
- Medical or Allied Health Professional (e.g., medical assistant, nurse, physician, social worker)
- Other Professional



FASD Post-Training Survey

Page 5 of 11

10. On a scale from 1 to 5 where 1 means you strongly disagree with the statement and 5 means you strongly agree, to what extent do you disagree or agree with the following statements. (Select one number per row).

	1 Strongly disagree	2 Disagree	3 Neutral	4 Agree	5 Strongly agree
a. It is important to routinely screen all patients/clients for alcohol use	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Screening a person for alcohol use confers a stigma to the person being screened	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. It is important to screen all pregnant women for alcohol use	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. It is important to screen all women of reproductive age for alcohol use	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. It is important to educate women of reproductive age, including those who are pregnant, about the effects of alcohol on a developing fetus	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
f. It is important to inquire about and document potential prenatal exposure for all pediatric patients	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
g. Diagnosis of one of the FASDs may confer a stigma to a child and/or his or her family	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>



FASD Post-Training Survey

Page 6 of 11

11. Are you a Resident or Medical/Allied Health Professional (e.g., medical assistant, nurse, physician, social worker)?

Yes

No



Online Survey Software Powered by novisurvey.net

FASD Post-Training Survey



12. On a scale from 1 to 5 where 1 means "Not at all confident in my skills" and 5 means "Totally confident in my skills," how confident are you in your skills to do the following? (Select one number per row).

	1 Not at all confident in my skills	2 Slightly confident in my skills	3 Moderately confident in my skills	4 Very confident in my skills	5 Totally confident in my skills
a. Asking women, including pregnant women, about their alcohol use	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. Having a conversation with patients/clients who indicate risky alcohol use	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. Educating women of childbearing age, including those who are pregnant, about the effects of alcohol on a developing fetus	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. Conducting brief interventions for reducing alcohol use	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. Utilizing resources to refer patients/clients who need formal treatment for alcohol abuse	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
f. Inquiring about potential prenatal alcohol exposure for my patients/clients	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
g. Identifying persons who may have one of the FASDs	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
h. Diagnosing persons who may have one of the FASDs	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
i. Referring patients/clients for diagnosis and/or treatment services for an FASD or alcohol use disorder	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
j. Managing/coordinating the treatment and care of persons who have one of the FASDs	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>



FASD Post-Training Survey

Page 8 of 11

13. Has your practice experienced any of the following barriers to effective implementation of alcohol screening and brief intervention? (Check all that apply).

- No barriers; we screen and intervene consistently and well
- Time limitations during patient/client visits
- It is not required; lack of incentive
- Attitudes of providers and/or staff about substance use/abuse
- Workforce needs education and training on screening and brief intervention
- Not easily accessible in the Electronic Health Record
- Concerns about damaging rapport with patients/clients
- Patient reluctance to be honest about alcohol use, or resistance to treatment
- Concerns about confidentiality and reporting requirements (example: to social services agency)
- Inadequate referral sources and/or system for making referrals
- Billing for alcohol screening, assessment, and counseling/intervention is not in place
- Patient/client inability to pay for treatment
- Other (please specify)



FASD Post-Training Survey

Page 9 of 11

14. On a scale from 1 to 5 where 1 means you strongly disagree with the statement and 5 means you strongly agree, to what extent do you disagree or agree with the following statements. (Select one number per row).

	1 Strongly disagree	2 Disagree	3 Neutral	4 Agree	5 Strongly agree
a. This training increased my understanding of the effects of prenatal alcohol exposure on the developing fetus.	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. The training concepts were presented clearly.	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. The training was presented in a culturally competent and sensitive manner	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
d. The content will be useful to me professionally	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
e. I would recommend this training to others	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
f. Overall, I am satisfied with the quality of this training.	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>



FASD Post-Training Survey

Page 10 of 11

15. What about this training could be improved?

16. Additional comments



Online Survey Software Powered by novisurvey.net

FASD Post-Training Survey

Page 11 of 11

Thank you for completing this survey.



Online Survey Software Powered by novisurvey.net