Public reporting burden of this collection of information is estimated to average 6 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Reports Clearance Officer; 1600 Clifton Road NE, MS D-74, Atlanta, Georgia 30333; ATTN: PRA (0920-0995).

OMB Control Number 0920-0995 Exp. Date 06/30/2023

Intensive Long-Term Evaluation

Observation discourses and an analysis of the state of th	
Strongly disagree Strongly agree	
2. The training improved the way I do my work	
Strongly disagree	
I am using what I learned in this training in my work	
Strongly disagree	
3a. If you have not used what you learned, please explain why not.	
4. In the prior evaluation, you replied to the question, "Do you intend to make a change in your practice, worksit or job as a result of this training?"	le setting,
Your answer was:ANSWER	
Your answer was:ANSWER Were you able to make this change?	

	If No, please explain
5.	As a result of the training, did you make other changes in your practice or at your worksite?
	Yes No Not my job
	Other reason (please specify)
_	
ba.	If yes, what change(s) did you make?
6.	Did any of these factors MAKE IT HARDER for you to incorporate the STI practices recommended in the presentation? (select all that apply
	Lack of time with patients More important patient concerns
	Cost/lack of reimbursement Policies where I work
	Resistance to change by supervisor or colleaguesLack of equipment or supplies
	 No opportunity to apply practices □ I did not feel confident
	Coworkers need training Nothing interfered
	Other (please specify)
7.	Did any of these factors HELP you incorporate the STI practices recommended in the presentation? (select all that apply)
	Reimbursement or other financial incentive Support of supervisor and/or colleagues
	Standing orders Reminder in chart
	☐ Convenient supplies
	Posted patient instructions for obtaining specimensElectronic health system
	 ☐ Knowledge/Confidence gained from training ☐ Trained coworkers
	Nothing specific helped
	Other (please specify)
	The following questions ask about screening rates. Please select the range that is the best estimate of your screening
R	The following questions ask about screening rates. Please select the range that is the best estimate of your screening practice
8.	The following questions ask about screening rates. Please select the range that is the best estimate of your screening
8.	The following questions ask about screening rates. Please select the range that is the best estimate of your screening practice Now, 3 months AFTER training, approximately what % of sexually active asymptomatic female patients under age 25
8.	The following questions ask about screening rates. Please select the range that is the best estimate of your screening practice Now, 3 months AFTER training, approximately what % of sexually active asymptomatic female patients under age 25 are you screening annually for chlamydia?
	The following questions ask about screening rates. Please select the range that is the best estimate of your screening practice Now, 3 months AFTER training, approximately what % of sexually active asymptomatic female patients under age 25 are you screening annually for chlamydia?
	The following questions ask about screening rates. Please select the range that is the best estimate of your screening practice Now, 3 months AFTER training, approximately what % of sexually active asymptomatic female patients under age 25 are you screening annually for chlamydia? On 1-25% 26-50% 51-75% 76-90% >91% NA Now, 3 months AFTER training, approximately what % of sexually active asymptomatic female patients under age 25

10.	Now, 3 months AFTER training, approximately what % of pregnant patients under age 25 are you screening for chlamydia?		
	0% 1-25% 26-50% 51-75% 76-90% >91% NA		
11.	Now, 3 months AFTER training, approximately what % of pregnant patients under age 25 are you screening for gonorrhea?		
	0% 1-25% 26-50% 51-75% 76-90% >91% NA		
12.	Now, 3 months AFTER training, approximately what % of pregnant patients are you screening for syphilis?		
	0% 1-25% 26-50% 51-75% 76-90% >91% NA		
13.	Now, 3 months AFTER training, approximately what % of your male patients who have sex with men are you screening annually for gonorrhea and chlamydia?		
	0% 1-25% 26-50% 51-75% 76-90% >91% NA		
14.	Now, 3 months AFTER training, approximately what % of your male patients who have sex with men are you screening at least once a year for syphilis?		
	0 0 0 0 0 0		

15.	5. Now, 3 months AFTER training, approximately what % of your patients recently diagnosed with syphilis are you t for HIV?				
	0% 1-25% 26-50% 51-75% 76-90% >91% NA				
16.	Now, 3 months AFTER training with approximately what % of your non-vaccinated 11 to 26 year-old female patients are you discussing HPV vaccination during a preventive health visit?				
	0% 1-25% 26-50% 51-75% 76-90% >91% NA				
17.	Now, 3 months AFTER training, with approximately what % of your non-vaccinated 11 to 21 year-old male patients are you discussing HPV vaccination during a preventive health visit?				
	0% 1-25% 26-50% 51-75% 76-90% >91% NA				
18.	Now, 3 months AFTER training, with approximately what % of your non-vaccinated male patients age 26 or under who have sex with men or are HIV+ are you discussing HPV vaccination during a preventive health visit?				
	0% 1-25% 26-50% 51-75% 76-90% >91% NA				
19.	Now, 3 months AFTER training, for approximately what % of patients older than 15 seeing you for a preventive health visit do you take a sexual history that asks about behaviors that would put them at risk of getting or transmitting an STI?				
	0% 1-25% 26-50% 51-75% 76-90% >91% NA				
20.	Now, 3 months AFTER training, for approximately what % of patients older than 15 seeing you for a preventive health visit do you take a sexual history that asks whether they have had oral, vaginal, or anal sex?				
	0% 1-25% 26-50% 51-75% 76-90% >91% NA				
	EXPEDITED PARTNER THERAPY				
21.	As a result of the information presented did you provide Expedited Partner Therapy (EPT) to heterosexual partners of those diagnosed with gonorrhea and/or chlamydia?				
	Yes No Not applicable to my practice or job My practice/worksite is in the planning stages to offer EPT My practice/worksite already offers EPT EPT was not discussed				
	As a result of the information presented did you				

23.	Use the CDC STI Treatment Guidelines in your practice?
	Yes No I was already do this
24.	Download the CDC STI Treatment Guidelines app?
	Yes No I was already do this
25.	Use the STI Treatment Guidelines wall chart or pocket guide?
	Yes No I was already do this
26.	Send a consult to the STD Clinical Consultation Network? www.stdccn.org
	Yes No I was already do this
27.	As a result of the information presented did you(please select 'Not Applicable if the training did not cover the content listed)
	vrogenital chlamydia and gonorrhea? Yes No I was already doing this Not applicable
29.	Increase the proportion of your male patients who have sex with men screened for syphilis, gonorrhea, and chlamydia at least annually?
	Yes No I was already doing this Not applicable
30.	Use CDC-recommended antibiotic therapy to treat uncomplicated gonorrhea?
	Yes No I was already doing this Not applicable
31.	Recommend rescreening in 3 months following a gonorrhea, chlamydia or trichomonas diagnosis?
	Yes No I was already doing this Not applicable
	THANK YOU!