

PRAMS Prescription and Illicit Opioid Use Supplemental Module

[PRAMS Opioid Supplemental Module – English Web](#)

 Department of Health and Human Services
Centers for Disease Control and Prevention

  

South Carolina
MomID: 2022TT333011  75%

The next questions are about the use of pain relievers *during pregnancy*.

 





South Carolina
MomID: 2022TT333011

75%

Q1. During your most recent pregnancy, did you use any of the following over-the-counter pain relievers? Over-the-counter pain relievers are those usually available without a prescription. For each one, check **No** if you did not use it *during* your pregnancy or **Yes** if you did.

	No	Yes
a. Acetaminophen (like regular Tylenol®, Tylenol Extra Strength®, or Tylenol PM®)	<input type="radio"/>	<input type="radio"/>
b. Ibuprofen (like Motrin® or Advil®), including high dose pills that may be prescribed	<input type="radio"/>	<input type="radio"/>
c. Aspirin (like Bayer® or Ecotrin®)	<input type="radio"/>	<input type="radio"/>
d. Naproxen (like Aleve® or Midol®)	<input type="radio"/>	<input type="radio"/>

Back

Next



South Carolina
MomID: 2022TT333011

76%

O2. During your most recent pregnancy, did you use any of the following *prescription* pain relievers?

For each one, check **No** if you did not use it *during* your pregnancy or **Yes** if you did. Do not include pain relievers you used *only* during labor and delivery.

	No	Yes
a. Hydrocodone (like Vicodin®, Norco®, or Lortab®)	<input type="radio"/>	<input type="radio"/>
b. Codeine (like Tylenol® #3 or #4, not regular Tylenol®)	<input type="radio"/>	<input type="radio"/>
c. Oxycodone (like Percocet®, Percodan®, OxyContin®, or Roxicodone®)	<input type="radio"/>	<input type="radio"/>
d. Tramadol (like Ultram® or Ultracet®)	<input type="radio"/>	<input type="radio"/>
e. Hydromorphone or meperidine (like Demerol®, Exalgo®, or Dilaudid®)	<input type="radio"/>	<input type="radio"/>
f. Oxymorphone (like Opana®)	<input type="radio"/>	<input type="radio"/>
g. Morphine (like MS Contin®, Avinza®, or Kadian ®)	<input type="radio"/>	<input type="radio"/>
h. Fentanyl (like Duragesic®, Fentora®, or Actiq®)	<input type="radio"/>	<input type="radio"/>

Back

Next





South Carolina

MomID: 2022TT333011

76%

The next questions are only about the use of *prescription* pain relievers listed in question Q2.

Back

Next





South Carolina

MomID: 2022TT333011

76%

O3. Where did you get the *prescription* pain relievers that you used *during* your most recent pregnancy?

	No	Yes
OB-GYN, midwife, or prenatal care provider	<input type="radio"/>	<input type="radio"/>
Family doctor or primary care provider	<input type="radio"/>	<input type="radio"/>
Dentist or oral health care provider	<input type="radio"/>	<input type="radio"/>
Doctor in the emergency room	<input type="radio"/>	<input type="radio"/>
I had pain relievers left over from an old prescription	<input type="radio"/>	<input type="radio"/>
Friend or family member gave them to me	<input type="radio"/>	<input type="radio"/>
I got the pain relievers <u>without a prescription</u> some other way	<input type="radio"/>	<input type="radio"/>
Other	<input type="radio"/>	<input checked="" type="radio"/>

Please tell us:

[Back](#)[Next](#)



South Carolina

MomID: 2022TT333011

77%

O4. What were your reasons for using *prescription pain relievers* during your most recent pregnancy?

	No	Yes
a. To relieve pain from an injury, condition, or surgery I had before pregnancy	<input type="radio"/>	<input type="radio"/>
b. To relieve pain from an injury, condition, or surgery that happened during my pregnancy	<input type="radio"/>	<input type="radio"/>
c. To relax or relieve tension or stress	<input type="radio"/>	<input type="radio"/>
d. To help me with my feelings or emotions	<input type="radio"/>	<input type="radio"/>
e. To help me sleep	<input type="radio"/>	<input type="radio"/>
f. To feel good or get high	<input type="radio"/>	<input type="radio"/>
g. Because I was “hooked” or I had to have them	<input type="radio"/>	<input type="radio"/>
h. Other	<input type="radio"/>	<input checked="" type="radio"/>

Please tell us:

[Back](#)[Next](#)



South Carolina

MomID: 2022TT333011

78%

05. In each of the following time periods *during your pregnancy*, for how many weeks or months did you use *prescription* pain relievers? Please write the total number of weeks OR months in each time period.

	Weeks	Months	Less than a week	Never
a. In the first 3 months of pregnancy	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
b. In the second 3 months of pregnancy	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
c. In the last 3 months of pregnancy	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

[Back](#)[Next](#)



Department of Health and Human Services

Centers for Disease Control and Prevention



South Carolina

MomID: 2022TT333011

78%

06. During your most recent pregnancy, did you want or need to cut down or stop using prescription pain relievers?

No

Yes

Back

Next





Department of Health and Human Services

Centers for Disease Control and Prevention



South Carolina

MomID: 2022TT333011

79%

07. During your most recent pregnancy, did you have trouble cutting down or stopping use of the prescription pain relievers?

No

Yes

Back

Next





Department of Health and Human Services

Centers for Disease Control and Prevention



South Carolina

MomID: 2022TT333011

80%

08. During your most recent pregnancy, did you get help from a doctor, nurse, or other health care worker to cut down or stop using prescription pain relievers?

No

Yes

Back

Next





Department of Health and Human Services

Centers for Disease Control and Prevention



South Carolina

MomID: 2022TT333011

81%

09. During your most recent pregnancy, did you receive medication-assisted treatment to help you stop using prescription pain relievers? This is when a doctor prescribes medicines such as methadone, buprenorphine, Suboxone®, Subutex® or naltrexone (Vivitrol®).

No

Yes

Back

Next





Department of Health and Human Services

Centers for Disease Control and Prevention



South Carolina

MomID: 2022TT333011

81%

O10. Do you think the use of *prescription pain relievers during pregnancy* could be harmful to a baby's health?

Check ONE answer

- Not harmful at all
- Not harmful, if taken as prescribed
- Harmful, even if taken as prescribed

Back

Next



prams-stg.cdc.gov

Solved: location n... Login Screen Acorn TV | Watch... Google Maps Fulton Water LOG IN

CDC Department of Health and Human Services Centers for Disease Control and Prevention

✉️ ☎️ ➔

South Carolina
MomID: 2022TT333011

82%

011. Do you think the use of *prescription* pain relievers could be harmful to a woman's own health?

Check ONE answer

Not harmful at all

Not harmful, if taken as prescribed

Harmful, even if taken as prescribed

Back Next





Department of Health and Human Services

Centers for Disease Control and Prevention



South Carolina

MomID: 2022TT333011

83%

O12. At any time *during your most recent pregnancy*, did a doctor, nurse, or other health care worker talk with you about how using prescription pain relievers during pregnancy could affect a baby?

No

Yes

Back

Next





Department of Health and Human Services

Centers for Disease Control and Prevention



South Carolina

MomID: 2022TT333011

84%

The next question is about the use of other medications or drugs during pregnancy.

Back

Next





Department of Health and Human Services

Centers for Disease Control and Prevention



South Carolina

MomID: 2022TT333011

84%

013. During your most recent pregnancy, did you take or use any of the following medications or drugs for any reason? For each item, check No if you did not take or use it or Yes if you did.

	No	Yes
a. Medication for depression (like Prozac®, Zoloft®, Lexapro®, Paxil®, or Celexa®)	<input type="radio"/>	<input type="radio"/>
b. Medication for anxiety (like Valium®, Xanax®, Ativan®, Klonopin®, or other "benzos" (benzodiazepines))	<input type="radio"/>	<input type="radio"/>
c. Methadone, Subutex®, Suboxone®, or buprenorphine	<input type="radio"/>	<input type="radio"/>
d. Naloxone	<input type="radio"/>	<input type="radio"/>
e. Cannabidiol (CBD) products	<input type="radio"/>	<input type="radio"/>
f. Adderall®, Ritalin®, or another stimulant	<input type="radio"/>	<input type="radio"/>
g. Marijuana or hash	<input type="radio"/>	<input type="radio"/>
h. Synthetic marijuana (K2 or Spice)	<input type="radio"/>	<input type="radio"/>
i. Heroin (smack, junk, Black Tar, or Chiva)	<input type="radio"/>	<input type="radio"/>
j. Amphetamines (uppers, speed, crystal meth, crank, ice, or agua)	<input type="radio"/>	<input type="radio"/>
k. Cocaine (crack, rock, coke, blow, snow, or nieve)	<input type="radio"/>	<input type="radio"/>
l. Tranquilizers (downers or ludes)	<input type="radio"/>	<input type="radio"/>
m. Hallucinogens (LSD/acid, PCP/angel dust, Ecstasy, Molly, mushrooms, or bath salts)	<input type="radio"/>	<input type="radio"/>
n. Sniffing gasoline, glue, aerosol spray cans, or paint to get high (huffing)	<input type="radio"/>	<input type="radio"/>

[Back](#)[Next](#)

PRAMS Opioid Supplemental Module – Spanish Web



Department of Health and Human Services

Centers for Disease Control and Prevention



South Carolina

MomID: 2022TT333012

77%

Las próximas preguntas son acerca del uso de medicamentos para el dolor durante el embarazo.

Anterior

Siguiente





Department of Health and Human Services

Centers for Disease Control and Prevention



South Carolina

MomID: 2022TT333012

77%

01. Durante su embarazo más reciente, ¿usó algunos de los siguientes medicamentos de venta libre ("over-the-counter") para aliviar el dolor? Los medicamentos de venta libre son aquellos que usualmente están disponibles sin receta. Para cada uno, marque **No**, si no lo usó *durante su embarazo*, o **Sí**, si lo usó.

	No	Sí
Acetaminofén (como Tylenol® regular, Tylenol Extra Strength® o Tylenol PM®)	<input type="radio"/>	<input type="radio"/>
Ibuprofeno (como Motrin® o Advil®) incluyendo aquellos de dosis más alta que podrían ser recetados.	<input type="radio"/>	<input type="radio"/>
Aspirina (como Bayer® o Ecotrin®)	<input type="radio"/>	<input type="radio"/>
Naproxen (como Aleve® o Midol®)	<input type="radio"/>	<input type="radio"/>

Anterior

Siguiente





Department of Health and Human Services

Centers for Disease Control and Prevention



South Carolina

MomID: 2022TT333012

77%

02. Durante su embarazo más reciente, ¿usó algunos de los siguientes medicamentos recetados para el dolor? Para cada uno, marque **No, si no lo usó durante su embarazo, o **Sí**, si lo usó. No incluya los medicamentos que utilizó solamente durante el parto y el nacimiento.**

	No	Sí
Hidrocodona (como Vicodin®, Norco® o Lortab®)	<input type="radio"/>	<input type="radio"/>
Codeína (como Tylenol® #3 o #4, <u>no</u> es el Tylenol regular)	<input type="radio"/>	<input type="radio"/>
Oxicodona (como Percocet®, Percodan®, OxyContin® o Roxicodone®)	<input type="radio"/>	<input type="radio"/>
Tramadol (como Ultram® o Ultracet®)	<input type="radio"/>	<input type="radio"/>
Hidromorfona o meperidina (como Demorol®, Exalgo® o Dilaudid®)	<input type="radio"/>	<input type="radio"/>
Oximorfona (como Opana®)	<input type="radio"/>	<input type="radio"/>
Morfina (como MS Contin®, Avinza® o Kadian®)	<input type="radio"/>	<input type="radio"/>
Fentanilo (como Duragesic®, Fentora® o Actiq®)	<input type="radio"/>	<input type="radio"/>

[Anterior](#)[Siguiente](#)



Department of Health and Human Services

Centers for Disease Control and Prevention



South Carolina

MomID: 2022TT333012

78%

Las siguientes preguntas son sólomente sobre el uso de medicamentos *recetados* para el dolor en la pregunta 02.

Anterior

Siguiente





Department of Health and Human Services

Centers for Disease Control and Prevention



South Carolina

MomID: 2022TT333012

78%

03. ¿Dónde obtuvo los medicamentos *recetados* para el dolor que usó durante su embarazo más reciente?

	No	Sí
Ginecólogo-obstetra, partera/comadrona o proveedor de cuidado prenatal	<input type="radio"/>	<input type="radio"/>
Médico de familia o un proveedor de atención primaria	<input type="radio"/>	<input type="radio"/>
Dentista o proveedor de atención de salud bucal (oral)	<input type="radio"/>	<input type="radio"/>
Doctor en la sala de emergencias	<input type="radio"/>	<input type="radio"/>
Tenía medicamentos para el dolor que me sobraron de una receta anterior	<input type="radio"/>	<input type="radio"/>
Un amigo o pariente me los dio	<input type="radio"/>	<input type="radio"/>
Obtuve los medicamentos para el dolor <u>sin receta</u> de otra forma	<input type="radio"/>	<input type="radio"/>
Otro	<input type="radio"/>	<input type="radio"/>

Anterior

Siguiente





Department of Health and Human Services

Centers for Disease Control and Prevention



South Carolina

MomID: 2022TT333012

79%

04. ¿Cuáles fueron sus razones para usar medicamentos recetados para el dolor durante su embarazo más reciente?

	No	Sí
Para tratar el dolor de una lesión, condición o cirugía que tuve antes del embarazo	<input type="radio"/>	<input type="radio"/>
Para tratar el dolor de una lesión, condición o cirugía que ocurrió durante mi embarazo	<input type="radio"/>	<input type="radio"/>
Para relajarme o aliviar la tensión o el estrés	<input type="radio"/>	<input type="radio"/>
Para ayudarme con mis sentimientos o emociones	<input type="radio"/>	<input type="radio"/>
Para ayudarme a dormir	<input type="radio"/>	<input type="radio"/>
Para sentirme bien o drogarme	<input type="radio"/>	<input type="radio"/>
Porque estaba adicta o tenía que usarlos	<input type="radio"/>	<input type="radio"/>
Otra	<input type="radio"/>	<input checked="" type="radio"/>

Por favor, escríbala:

Anterior

Siguiente





Department of Health and Human Services

Centers for Disease Control and Prevention



South Carolina

MomID: 2022TT333012

80%

05. En cada uno de los siguientes periodos de tiempo durante su embarazo, ¿cuántas semanas o meses usó medicamentos recetados para el dolor? Por favor escriba el número total de semanas o meses en cada periodo de tiempo.

	Semanas	Meses	Menos de una semana	Nunca
En los primeros 3 meses del embarazo	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
En los segundos 3 meses del embarazo	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
En los últimos 3 meses del embarazo	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Anterior

Siguiente



Attachment 11h – PRAMS Opioid Use Supplement_English and Spanish Web



Department of Health and Human Services

Centers for Disease Control and Prevention



South Carolina

MomID: 2022TT333012

80%

06. Durante su embarazo más reciente, ¿quería o necesitaba reducir o dejar de usar medicamentos recetados para el dolor?

No

Sí

[Anterior](#)

[Siguiente](#)



Attachment 11h – PRAMS Opioid Use Supplement_English and Spanish Web



South Carolina

MomID: 2022TT333012

81%

07. Durante su embarazo más reciente, ¿tuvo problemas reduciendo o dejando de usar los medicamentos recetados para el dolor?

No

Sí

[Anterior](#)

[Siguiente](#)



Attachment 11h – PRAMS Opioid Use Supplement_English and Spanish Web



Department of Health and Human Services

Centers for Disease Control and Prevention



South Carolina

MomID: 2022TT333012

82%

08. Durante su embarazo más reciente, ¿recibió ayuda de un doctor, enfermera u otro profesional de la salud para reducir o dejar de usar medicamentos recetados para el dolor?

No

Sí

[Anterior](#)

[Siguiente](#)





Department of Health and Human Services

Centers for Disease Control and Prevention



South Carolina

MomID: 2022TT333012

82%

09. Durante su embarazo más reciente, ¿recibió tratamiento asistido por medicación para ayudarle a dejar de usar medicamentos recetados para el dolor? Esto es cuando un doctor receta medicinas como metadona, buprenorfina, Suboxone®, Subutex® o naltrexone Vivitrol®.

No

Sí

[Anterior](#)

[Siguiente](#)





Department of Health and Human Services

Centers for Disease Control and Prevention



South Carolina

MomID: 2022TT333012

83%

Q10. ¿Cree que el uso de medicamentos *recetados* para el dolor durante el embarazo podría ser dañino para la salud de un bebé?

Marque UNA respuesta

- No es dañino para nada
- No es dañino, siempre y cuando sean tomados según indica la receta
- Sí es dañino, aunque sean tomados según indica la receta

Anterior

Siguiente





Department of Health and Human Services

Centers for Disease Control and Prevention



South Carolina

MomID: 2022TT333012

84%

011. ¿Cree que el uso de medicamentos *recetados* para el dolor podría ser dañino para la salud de una mujer?

Marque UNA respuesta

- No es dañino para nada
- No es dañino, siempre y cuando sean tomados según indica la receta
- Sí es dañino, aunque sean tomados según indica la receta

Anterior

Siguiente





Department of Health and Human Services

Centers for Disease Control and Prevention



South Carolina

MomID: 2022TT333012

85%

012. En cualquier momento durante su embarazo más reciente, ¿algún doctor, enfermera u otro profesional de la salud habló con usted sobre cómo el uso de medicamentos recetados para el dolor durante el embarazo podría afectar a un bebé?

No

Sí

[Anterior](#)

[Siguiente](#)



Attachment 11h – PRAMS Opioid Use Supplement_English and Spanish Web

 Department of Health and Human Services
Centers for Disease Control and Prevention

South Carolina
MomID: 2022TT333011 85%

O13. Durante su embarazo más reciente, ¿tomó o usó algunas de los siguientes medicamentos o drogas por alguna razón? Para cada una, marque No, si no la tomó o usó, o Sí, si la tomó o usó.

	No	Sí
Medicamentos para la depresión (como Prozac®, Zoloft®, Lexapro®, Paxil® o Celexa®)	<input type="radio"/>	<input type="radio"/>
Medicamentos para la ansiedad (como Valium®, Xanax®, Ativan®, Klonopin®, u otro tipos de "benzos" (benzodiacepinas))	<input type="radio"/>	<input type="radio"/>
Metadona, Subutex®, Suboxone® o buprenorfina	<input type="radio"/>	<input type="radio"/>
Naloxona	<input type="radio"/>	<input type="radio"/>
Productos de Cannabidiol (CBD)	<input type="radio"/>	<input type="radio"/>
Adderall®, Ritalin® u otro estimulante	<input type="radio"/>	<input type="radio"/>
Marihuana o hachís	<input type="radio"/>	<input type="radio"/>
Marihuana sintética (K2, Spice)	<input type="radio"/>	<input type="radio"/>
Heroína (<i>smack, junk, Black Tar o chiva</i>)	<input type="radio"/>	<input type="radio"/>
Metanfetaminas (<i>uppers, speed, crystal meth, crank, ice o agua</i>)	<input type="radio"/>	<input type="radio"/>
Cocaína (<i>crack, rock, coke, blow, snow o nieve</i>)	<input type="radio"/>	<input type="radio"/>
Tranquilizantes (<i>calmantes, downers o ludes</i>)	<input type="radio"/>	<input type="radio"/>
Alucinógenos (LSD/ácido, PCP/polvo de ángel, éxtasis, <i>Molly</i> , hongos, o sales de baño)	<input type="radio"/>	<input type="radio"/>
Inhalación de gasolina, pegamento, aerosoles en latas o pintura para drogarse (<i>huffing</i>)	<input type="radio"/>	<input type="radio"/>

[Anterior](#) [Siguiente](#)

Centers for Disease Control and Prevention
<https://prams-stg.cdc.gov/#> U.S.A.

Department of Health and Human Services
[Vulnerability Disclosure Policy | Español](#)