# **Radiologic Interpretation**

## **U.S. DEPARTMENT OF LABOR**

OFFICE OF WORKERS' COMPENSATION PROGRAMS DIVISION OF COAL MINE WORKERS' COMPENSATION



Note: This report is authorized by law (30 USC 901 et. seq. and 20 CFR 718.102). The results of this interpretation will aid in determining the claimant's eligibility for black lung benefits. This method of collecting information complies with the Freedom of Information Act, the Privacy Act of 1974, and OMB Circular No. 108.

OMB No. 1240-0023 Expires 12/31/2023

Please record your interpretation of a single image by placing "X" in the appropriate boxes on the form and return it promptly to the office that requested the interpretation. The form must be completed as per instructions, signed by a physician, and contain the miner's name and DOL's Case ID Number. The Department of Labor will pay only for images of acceptable quality (1, 2 and Images of inferior quality (U/R) must be retaken without cost to the Department. 1B. DOL's Case ID Number 1. Miner's Name (Print) 1A. Date of X-Ray 1C. Image Quality (if not Grade 1. Give Reason): 2 3 U/R 1 MC DAY 1D. Is Image Completely Negative? 2A. Any Parenchymal Abnormalities Consistent with Pneumoconiosis? YES Proceed to Section 5 NO Complete Section 2A YES Complete 2B and 2C NO Proceed to Section 3 Small Opacities Consistent With Pneumoconiosis 2C. Large Opacities Consistent With Pneumoconiosis c. PROFUSION a. SHAPE/SIZE SECONDARY PRIMARY b. ZONES 0/-0/0 0/1 р s р s 1/0 1/1 1/2 Proceed to SI С 2/1 2/2 0 q t q t 2/3 7F 3/+ u u 3/2 3/3 3A. ANY PLEURAL ABNORMALITIES Complete Sections Proceed to CONSISTENT WITH PNEUMOCONIOSIS? YES NO 3B, 3C Section 4A 3B. PLEURAL PLAQUES (mark site, calcification, extent and width) Extent (chest wall; combined for Width (in profile only) in profile and face on) (3mm minimum width required) Chest Wall Calcification Up to 1/4 of lateral chest wall = 1 3 to 5 mm = aSite In Profile О R L О R L 1/4 to 1/2 of lateral chest wall = 2 5 to 10 mm = bR R > 10 mm = c Face On 0 ī Ω > 1/2 of lateral chest wall = 3 1 Diaphragm 0 R L 0 R L 0 R 0 0 R 0 Other site(s) 0 R О R L 2 3 2 3 b b а С Proceed to Proceed to 3C. COSTOPHRENIC ANGLE OBLITERATION R L NO 3D. DIFFUSE PLEURAL THICKENING (mark site, calcification, extent, and width) Extent (chest wall, combined for Width (in profile only) in profile and face on) (3mm minimum width required) Up to 1/4 of lateral chest wall = 1 3 to 5 mm = a1/4 to 1/2 of lateral chest wall = 2 5 to 10 mm = bCalcification > 10 mm = c Chest wall > 1/2 of lateral chest wall = 3 R L R 0 R R In Profile 0 O 0 0 0 L R 2 3 3 b b Face On а С С 4A. ANY OTHER ABNORMALITIES? Complete Proceed to YES 4B and 4C Section 5 4B. OTHER SYMBOLS (OBLIGATORY pb tb aa at ax ca cg cn СО ср cv ho pa pi рх ra rp REPORT ITEMS WHICH MAY BE OF PRESENT (Specify od.) Date Personal Physician notified? OD D D CLINICAL SIGNIFICANCE IN THIS SECTION OTHER COMMENTS SHOULD WORKER SEE PERSONAL PHYSICIAN BECAUSE OF COMMENTS IN SECTION 4C? YES NO Proceed to Section 5 FACILITY PROVIDING RADIOLOGIC EXAMINATION: DOL Medical Provider Number (if applicable): Was image taken by a registered radiographer/radiographic technologist? ☐ Yes ☐ No State Name Registration No. 5B. Physician Interpreting Image (Print Name): Are you: Board-certified radiologist? Yes No Board-eligible radiologist? Yes No B-reader? Yes No Date Current B-reader Certification Expires: I certify that this image has been interpreted in accordance with the instructions provided on Form CM-954a and/or 20 CFR 718, Subpart B, 718.102 and Appendix A. I also certify that the information furnished is correct and am aware that my signature attests to the accuracy of the results reported. I am aware that any person who willfully makes any false or misleading statement or representation in support of an application for benefits shall be guilty of a misdemeanor under 30 USC 941 and, on conviction, subject to a fine of up to \$1,000, or to imprisonment for up to one year, or both. PHYSICIAN'S SIGNATURE DATE OF READING (Mo., Day, Yr.)

### **PUBLIC BURDEN STATEMENT**

We estimate that it will take an average of 5 minutes to complete this information collection, including time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the information. If you have any comments regarding these estimates or any other aspect of this information collection, including suggestions for reducing this burden, send them to the Division of Coal Mine Workers' Compensation, U. S. Department of Labor, Room N-3464, 200 Constitution Avenue, N.W., Washington, D.C. 20210. **DO NOT SEND THE COMPLETED FORM TO THIS OFFICE**.

#### PRIVACY ACT NOTICE

In accordance with the Privacy Act of 1974, as amended (5 U.S.C. 552a), you are hereby notified that: (1) the Black Lung Benefits Act (BLBA) (30 U.S.C. 901 et seq.), as amended, is administered by the Office of Workers' Compensation Programs (OWCP) of the U.S. Department of Labor, which receives and maintains personal information, relative to this application, on claimants and their immediate families; (2) information obtained by OWCP will be used to determine eligibility for benefits payable under the BLBA; (3) information may be given to other government agencies, coal mine operators potentially liable for

payment of the claim or to the insurance carrier or other entity which secured the operator's compensation liability, contractors providing automated data processing services to the Department of Labor; and representatives of the parties to the claim; (4) information may be given to physicians or other medical service providers for use in providing treatment, making evaluations and for other purposes relating to the medical management of the claim; (5) information may be given to the Department of Labor's Office of Administrative Law Judges, or other person, board or organization, which is authorized or required to render decisions with respect to the claim or other matters arising in connection with the claim; (6) information may be given to Federal, state or local agencies

for law enforcement purposes, to obtain information relevant to a decision under the BLBA, to determine whether benefits are being or have been paid properly.

and where appropriate, to pursue administrative offset and/or debt collection actions required or permitted by law; (7) disclosure of the claimant's or deceased miner's Social Security Number (SSN) or tax identifying number (TIN) on this form is voluntary, and the SSN and/or TIN and other information maintained by the OWCP may be used for identification and for other purposes authorized by law; (8) failure to disclose all requested information, may delay the processing of this claim or the payment of benefits, or may result in an unfavorable decision or reduced level of benefits; and (9) this information is included in a System of

Records, DOL/OWCP-2 published at 81 Federal Register 25765, 25858 (April 29, 2016) or as updated and republished.

### NOTICE

If you have a substantially limiting physical or mental impairment, Federal disability nondiscrimination law gives you the right to receive help from OWCP in the form of communication assistance, accommodation and modification to aid you in the claims process. For example, we will provide you with copies of documents in alternate formats, communication services such as sign language interpretation, or other kinds of adjustments or changes to account for the limitations of your disability. Please contact our office or the claims examiner to ask about this assistance.

NOTE: Persons are not required to respond to this collection of information unless it displays a currently valid OMB control number

# TWO FILING OPTIONS:

- 1. To file electronically, submit completed form to the COAL Mine Portal: <a href="https://eclaimant.dol.gov/portal/?program\_name=BL">https://eclaimant.dol.gov/portal/?program\_name=BL</a>
- To file by mail, send completed form to: US Department of Labor OWCP/DCMWC PO Box 8307 London, KY 40742-8307

For Purpose of Coding for the Department of Labor, the following codes will be used ILO 2011 INTERNATIONAL CLASSIFICATION OF RADIOGRAPHS OF THE PNEUMOCONIOSES

FEATURE		CODES	ON OF RADIOGRAPHS OF THE PNEUMOCONIOSES  DEFINITIONS
Technical Quality	:5	1	Good
-		2	Acceptable, with no technical defect likely to impair classification of the radiograph for pneumoconiosis.  Acceptable, with some technical defect but still acceptable for classification purposes.
Parenchymal Abnormalities Small Opacities		3	Unacceptable.
Small Opacities		U/R	The category of profusion is based on the assessment of concentration of opacities by comparison with the standard radiographs.  Category 0 – small opacities absent or less profuse than the lower limit of Category 1.
	Profusion	0/- 0/0 0/1 1/0 1/1 1/2	Categories 1, 2 and 3 - represent increasing profusion of small opacities as defined by the corresponding standard radiographs.
		2/1 2/2 2/3 3/2 3/3 3/+	The zones in which the opacities are seen are recorded. The right (R) and
			left (L) thorax are both divided into three zones – upper (U), middle (M) and lower (L).
	Zones	RU RM RL	The category of profusion is determined by considering the profusion as a whole over the affected zones of the lung and by comparing this with the standard radiographs.
		LU LM LL	The letters p, q, and r denote the presence of small rounded opacities.  Three sizes are defined by the appearances on standard radiographs.
			p = diameter up to about 1.5 mm.
			q = diameter exceeding about 1.5 mm and up to about 3 mm. r = diameter exceeding about 3 mm and up to about 10 mm.
	Shape and Size rounded	n/n a/a r/r	The letters s, t and u denote the presence of small irregular opacities.
		p/p q/q r/r	Three sizes are defined by the appearance on standard radiographs.
			s = width up to about 1.5 mm. t = width exceeding about 1.5 mm and up to about 3 mm.
	irregular	s/s t/t u/u	u = width exceeding 3 mm and up to about 10 mm.  For mixed shapes (or sizes) of small opacities the predominant shape
		p/s p/t p/u p/g p/r	And size is recorded first. The presence of a significant number or another shape and size is recorded after the oblique stroke.
	mixed	g/s g/t g/u g/p g/r r/s r/t r/u r/p r/g	recorded direct the oblique stroke.
		p/s p/t p/u p/q p/r q/s q/t q/u q/p q/r r/s r/t r/u r/p r/q s/p s/q s/r s/t s/u t/p t/q t/r t/s t/u	
Large Opacities		u/p u/q u/r u/s u/t ABC	The categories are defined in terms of dimensions of the opacities.
			Category A – an opacity having a greatest diameter exceeding about 10 mm and up to and
			including 50 mm, or several opacities each greater than about 10 mm, the sum of whose greatest diameters does not exceed 50 mm.
			Category B – one or more opacities larger or more numerous than those in category A whose combined area does not exceed the equivalent of the right upper zone.
			Category C – one or more opacities whose combined area exceed the equivalent of the right upper
Diaural Abnormalities			zone.
Pleural Abnormalities			Two types of pleural thickening of the chest wall are recognized: circumscribed (plaques) and diffuse. Both types may occur together.
Pleural Thickening Chest Wall	Туре		ordanistrised (staques) and amase. Both types may occur together.
Chest Wall	Site	R L	Pleural thickening of the chest wall is recorded separately for the
			right (R) and left (L) thorax.
	Width	ABC	For pleural thickening seen along the lateral chest wall the measurement of maximum width is made from the inner line of the chest wall to the inner margin of the shadow seen most sharply at the parenchymal-pleural boundary. The maximum width usually occurs at the inner margin of the rib shadow at its outermost point.
			a = maximum width up to about 5 mm. b = maximum width over about 5 mm and up to about 10 mm. c = maximum width over about 10 mm.
	Face On	Y N	<ul> <li>c = maximum width over about 10 mm.</li> <li>The presence of pleural thickening seen face-on is recorded even if it can be seen also in profile. If pleural thickening is seen face-on only,</li> </ul>
			width cannot usually be measured.
	Extent	1 2 3	Extent of pleural thickening is defined in terms of the maximum length of pleural involvement, or as the sum of maximum lengths, whether seen in profile or face-on.
			1 = total length equivalent up to one quarter of the projection of the lateral chest wall.
			2 = total length exceed one quarter but not one half of the projection of the lateral chest wall
			3 = total length exceeding one half of the projection of the lateral chest lateral chest wall
Diaphragm	Presence	Y N	A plaque involving the diaphragmatic playra is recorded as proceed (V)
-13			A plaque involving the diaphragmatic pleura is recorded as present (Y) or absent (N) separately for the right (R) or left (L) thorax.  The presence (Y) or absence (N) costophragic angle obligeration is
Costophrenic Angle	Site Presence	R L Y N	The presence (Y) or absence (N) costophrenic angle obliteration is recorded separately from thickening over other areas for the right (R) and left (L) thorax. The lower limit for the obliteration is defined by a standard radiograph showing profusion subcategory 1/1 t/t.
			standard radiograph showing profusion subcategory 1/1 Vt.  If the thickening extends up the chest wall then both costophrenic
Pleural Calcification	Site	R L	If the thickening extends up the chest wall then both costophrenic angle obliteration and pleural thickening should be recorded.
i icarai CaiciiiCatiOff	Sito		The site and extent of pleural calcification are recorded separately for the two lungs, and the extent defined in terms of dimensions.
	Site		
	chest wall diaphragm	R L	"Other" includes calcification of the mediastinal and pericardial pleura.
	chest wall diaphragm other	R L R L R L	1 = an area of calcified pleura with greatest diameter up to about 20 mm
	chest wall diaphragm	R L	1 = an area of calcified pleura with greatest diameter up to about 20 mm or a number of such areas the sum of whose greatest diameters does not exceed about 20 mm.
	chest wall diaphragm other	R L R L	1 = an area of calcified pleura with greatest diameter up to about 20 mm or a number of such areas the sum of whose greatest diameters does not exceed about 20 mm.
	chest wall diaphragm other	R L R L	1 = an area of calcified pleura with greatest diameter up to about 20 mm or a number of such areas the sum of whose greatest diameters does not exceed about 20 mm.  2 = an area of calcified pleura with greatest diameter exceeding about 20 mm and up to about 100 mm, or a number of such areas the sum of whose greatest diameters exceed about 20 mm but dies not exceed about 100 mm.
	chest wall diaphragm other	R L R L	1 = an area of calcified pleura with greatest diameter up to about 20 mm or a number of such areas the sum of whose greatest diameters does not exceed about 20 mm.  2 = an area of calcified pleura with greatest diameter exceeding about 20 mm and up to about 100 mm, or a number of such areas the sum of whose greatest diameters exceed about 20 mm but dies not exceed about 100 mm.  3 = an area of calcified pleura with greatest diameter exceeding about 100 mm or a number of such areas with greatest diameters.
	chest wall diaphragm other	R L R L	1 = an area of calcified pleura with greatest diameter up to about 20 mm or a number of such areas the sum of whose greatest diameters does not exceed about 20 mm.  2 = an area of calcified pleura with greatest diameter exceeding about 20 mm and up to about 100 mm, or a number of such areas the sum of whose greatest diameters exceed about 20 mm but dies not exceed about 100 mm.
	chest wall diaphragm other	R L R L	1 = an area of calcified pleura with greatest diameter up to about 20 mm or a number of such areas the sum of whose greatest diameters does not exceed about 20 mm.  2 = an area of calcified pleura with greatest diameter exceeding about 20 mm and up to about 100 mm. or a number of such areas the sum of whose greatest diameters exceed about 20 mm but dies not exceed about 100 mm.  3 = an area of calcified pleura with greatest diameter exceeding about 100 mm or a number of such area whose sum of greatest diameters exceeds about 100 mm.  It is to be taken that the definition of such of the Symbols is preceded by an appropriate word or phrase such as "suspect", "pneumoconiotic
	chest wall diaphragm other	R L R L	1 = an area of calcified pleura with greatest diameter up to about 20 mm or a number of such areas the sum of whose greatest diameters does not exceed about 20 mm.  2 = an area of calcified pleura with greatest diameter exceeding about 20 mm and up to about 100 mm, or a number of such areas the sum of whose greatest diameters exceed about 20 mm but dies not exceed about 100 mm.  3 = an area of calcified pleura with greatest diameter exceeding about 100 mm or a number of such area whose sum of greatest diameters exceeds about 100 mm.  It is to be taken that the definition of such of the Symbols is preceded
	chest wall diaphragm other	R L R L	1 = an area of calcified pleura with greatest diameter up to about 20 mm or a number of such areas the sum of whose greatest diameters does not exceed about 20 mm.  2 = an area of calcified pleura with greatest diameter exceeding about 20 mm and up to about 100 mm, or a number of such areas the sum of whose greatest diameters exceed about 20 mm but dies not exceed about 100 mm.  3 = an area of calcified pleura with greatest diameter exceeding about 100 mm or a number of such area whose sum of greatest diameters exceeds about 100 mm.  It is to be taken that the definition of such of the Symbols is preceded by an appropriate word or phrase such as "suspect", "pneumoconiotic
Symbols  aa - atherosclerotic	chest wall diaphragm other	R L R L	1 = an area of calcified pleura with greatest diameter up to about 20 mm or a number of such areas the sum of whose greatest diameters does not exceed about 20 mm.  2 = an area of calcified pleura with greatest diameter exceeding about 20 mm and up to about 100 mm. or a number of such areas the sum of whose greatest diameters exceed about 20 mm but dies not exceed about 100 mm.  3 = an area of calcified pleura with greatest diameter exceeding about 100 mm or a number of such area whose sum of greatest diameters exceeds about 100 mm.  It is to be taken that the definition of such of the Symbols is preceded by an appropriate word or phrase such as "suspect", "pneumoconiotic changes suggestive of", or "opacities suggestive of ", etc.
Symbols aa - atherosclerotic at - significant apical pleural thick	chest wall diaphragm other extent	R L R L	1 = an area of calcified pleura with greatest diameter up to about 20 mm or a number of such areas the sum of whose greatest diameters does not exceed about 20 mm.  2 = an area of calcified pleura with greatest diameter exceeding about 20 mm and up to about 100 mm, or a number of such areas the sum of whose greatest diameters exceed about 20 mm but dies not exceed about 100 mm.  3 = an area of calcified pleura with greatest diameter exceeding about 100 mm or a number of such area whose sum of greatest diameters exceeds about 100 mm.  It is to be taken that the definition of such of the Symbols is preceded by an appropriate word or phrase such as "suspect", "pneumoconiotic
aa - atherosclerotic at - significant apical pleural thicl ax - coalescence of small pneum	chest wall diaphragm other extent	R L R L	1 = an area of calcified pleura with greatest diameter up to about 20 mm or a number of such areas the sum of whose greatest diameters does not exceed about 20 mm.  2 = an area of calcified pleura with greatest diameter exceeding about 20 mm and up to about 1000 mm, or a number of such areas the sum of whose greatest diameters exceed about 20 mm but dies not exceed about 100 mm.  3 = an area of calcified pleura with greatest diameter exceeding about 100 mm or a number of such area whose sum of greatest diameters exceeds about 100 mm.  It is to be taken that the definition of such of the Symbols is preceded by an appropriate word or phrase such as "suspect", "pneumoconiotic changes suggestive of", or "opacities suggestive of ", etc.  hi - enlargement of non-calcified hilar or mediastinal lymph nodes ho - honeycomb lung id - ill-defined diaphragm border
aa - atherosclerotic at - significant apical pleural thici ax - coalescence of small pneum bu - bulla(e)	chest wall diaphragm other  extent  extent  kening noconiotic opacities	R L R L	1 = an area of calcified pleura with greatest diameter up to about 20 mm or a number of such areas the sum of whose greatest diameters does not exceed about 20 mm.  2 = an area of calcified pleura with greatest diameter exceeding about 20 mm and up to about 100 mm, or a number of such areas the sum of whose greatest diameters exceed about 20 mm but dies not exceed about 100 mm.  3 = an area of calcified pleura with greatest diameter exceeding about 100 mm exceed about 100 mm.  It is to be taken that the definition of such of the Symbols is preceded by an appropriate word or phrase such as "suspect", "pneumoconiotic changes suggestive of", or "opacities suggestive of ", etc.  hi - enlargement of non-calcified hilar or mediastinal lymph nodes  ho - honeycomb lung  id - ill-defined diaphragm border  ih - ill-defined heart border
aa - atherosclerotic at - significant apical pleural thick ax - coalescence of small pneum bu - bulla(e) ca - cancer thoracic malignancies	chest wall diaphragm other  extent  extent  kening noconiotic opacities	R L R L 1 2 3	1 = an area of calcified pleura with greatest diameter up to about 20 mm or a number of such areas the sum of whose greatest diameters does not exceed about 20 mm.  2 = an area of calcified pleura with greatest diameter exceeding about 20 mm and up to about 100 mm, or a number of such areas the sum of whose greatest diameters exceed about 20 mm but dies not exceed about 100 mm.  3 = an area of calcified pleura with greatest diameter exceeding about 100 mm or a number of such area whose sum of greatest diameters exceeds about 100 mm.  It is to be taken that the definition of such of the Symbols is preceded by an appropriate word or phrase such as "suspect", "pneumoconiotic changes suggestive of", or "opacities suggestive of ", etc.  hi - enlargement of non-calcified hilar or mediastinal lymph nodes ho - honeycomb lung id - ill-defined diaphragm border
aa - atherosclerotic at - significant apical pleural thic ax - coalescence of small pneum bu - bulla(e) ca - cancer thoracic malignancies cg - calcified non-pneumoconiotic cn - calcification in small pneumo	chest wall diaphragm other extent  extent  kening noconiotic opacities s excluding mesothelioma c nodules (e.g. granuloma) or noconiotic opacities	R L R L 1 2 3	1 = an area of calcified pleura with greatest diameter up to about 20 mm or a number of such areas the sum of whose greatest diameters does not exceed about 20 mm.  2 = an area of calcified pleura with greatest diameter exceeding about 20 mm and up to about 100 mm.  3 = an area of calcified pleura with greatest diameter exceeding about 20 mm but dies not exceed about 100 mm.  3 = an area of calcified pleura with greatest diameter exceeding about 100 mm or a number of such area whose sum of greatest diameters exceeds about 100 mm.  It is to be taken that the definition of such of the Symbols is preceded by an appropriate word or phrase such as "suspect", "pneumoconiotic changes suggestive of", or "opacities suggestive of ", etc.  hi - enlargement of non-calcified hilar or mediastinal lymph nodes ho - honeycomb lung id - ill-defined diaphragm border ih - ill-defined heart border kl - septal (Kerley) lines me - mesothelioma pa - plate atelectasis
aa - atherosclerotic at - significant apical pleural thick ax - coalescence of small pneum bu - bulla(e) ca - cancer thoracic malignancies cg - calcified non-pneumoconiotic	chest wall diaphragm other extent  extent  kening noconiotic opacities s excluding mesothelioma c nodules (e.g. granuloma) or noconiotic opacities	R L R L 1 2 3	1 = an area of calcified pleura with greatest diameter up to about 20 mm or a number of such areas the sum of whose greatest diameters does not exceed about 20 mm.  2 = an area of calcified pleura with greatest diameter exceeding about 20 mm and up to about 100 mm.  3 = an area of calcified pleura with greatest diameter exceeding about exceed about 100 mm.  3 = an area of calcified pleura with greatest diameter exceeding about 100 mm or a number of such area whose sum of greatest diameters exceed about 100 mm or a number of such area whose sum of greatest diameters exceeds about 100 mm.  It is to be taken that the definition of such of the Symbols is preceded by an appropriate word or phrase such as "suspect", "pneumoconiotic changes suggestive of", or "opacities suggestive of ", etc.  hi - enlargement of non-calcified hilar or mediastinal lymph nodes  ho - honeycomb lung  id - ill-defined diaphragm border  ih - ill-defined heart border  kl - septal (Kerley) lines  me - mesothelioma

di	di - marked distortion of an intrathoracic structure				ra	- rounded atelectasis	
ef	ef - pleural effusion				rp	- rheumatoid pneumoconiosis	
em	m - emphysema				tb	- tuberculosis	
es	es - eggshell calcification of hilar or mediastinal lymph nodes				od	- other disease or significant abnormality	
fr	fr - fractured rib(s) (acute or healed)						
Comments		Presence	Y N			Comments should be recorded pertaining to the classification of the radiograph particularly if some other cause is thought to be responsible for a shadow.	

CM-933 (Rev. August 2020)