SCID-5-MICS

STRUCTURED CLINICAL INTERVIEW FOR DSM-5® DISORDERS

MODIFIED FOR THE MENTAL ILLNESS CALIBRATION STUDY

Michael B. First, M.D., Janet B. W. Williams, Ph.D., Rhonda S. Karg, Ph.D., and Robert L. Spitzer, M.D.

Copyright © 2015 Michael B. First, M.D., Janet B. W. Williams, Ph.D., and Robert L. Spitzer, M.D.

For citation: First MB, Williams JBW, Karg RS, Spitzer RL: *User's Guide for the Structured Clinical Interview for DSM-5 Disorders, Research Version* (SCID-5-RV). Arlington, VA, American Psychiatric Association, 2015

The Structured Clinical Interview for DSM-5®, Research Version (SCID-5-RV), includes the User's Guide and score sheets. Use of any component of the SCID-5-RV requires permission or licensing through American Psychiatric Publishing before use. Inquiries should be directed to SCID Permissions & Licensing, American Psychiatric Publishing, 1000 Wilson Boulevard, Suite 1825, Arlington, VA 22209-3901, or online at: http://www.appi.org/CustomerService/Pages/Permissions.aspx. For more information, please visit the SCID products page on www.appi.org.

DSM and DSM-5 are registered trademarks of the American Psychiatric Association. Use of these terms is prohibited without permission of the American Psychiatric Association.

DSM-5® diagnostic criteria are reprinted or adapted with permission from American Psychiatric Association: *Diagnostic and Statistical Manual of Mental Disorders*, Fifth Edition. Arlington VA, American Psychiatric Association, 2013. Copyright © 2013 American Psychiatric Association. Used with permission.

ALL RIGHTS RESERVED. Unless authorized in writing by the American Psychiatric Association (APA), no part of the DSM-5® criteria may be reproduced or used in a manner inconsistent with the APA's copyright. This prohibition applies to unauthorized uses or reproductions in any form, including electronic applications. Correspondence regarding copyright permission for DSM-5 criteria should be directed to DSM Permissions, American Psychiatric Publishing, 1000 Wilson Boulevard, Suite 1825, Arlington, VA 22209-3901, www.appi.org.

All versions of the SCID-5 represent the findings, conclusions, and views of the individual authors and do not necessarily represent the policies and opinions of American Psychiatric Publishing or the American Psychiatric Association.

Respondent:	Date of Interview:
Clinical Interviewer:	Month Day Year

Nonpatient Overview

I'm going to be asking you about problems or difficulties you may have had, and I'll be making some notes as we go along. Do you have any questions before we begin?

NOTE: Any current suicidal thoughts, plans, or actions should be thoroughly assessed by the clinician and action taken if necessary.

Demographic Data

What is your current gender?	GENDER:	
What's your date of birth?	DOB: AGE: month day year	ONP2 ONP3
Are you married?	MARITAL STATUS (most recent):	
IF NO: Do you live with someone as if you are married?	1 Married or living with someone as if married2 Widowed	ONP4
IF NO: Were you ever married?	3 Divorced or annulled4 Separated	
How long have you been (MARITAL STATUS)?	5 Never married	
IF EVER MARRIED: How many times have you been married?		
Do you have any children?		
IF YES: How many? (What are their ages?)		
With whom do you live? (How many children under the age of 18 live in your household?)		
In what city, town, or neighborhood do you live?		
In what kind of place do you live? (A house, an apartment, a shelter, a halfway house, or some other living arrangement? Are you homeless?)		
I would like to now ask you about your cultural background or identity. By background or identity, I mean, for example, the communities you belong to, the languages you speak, where you or your family are from, your race or ethnic background, your gender or sexual orientation, or your faith or religion.	1	
For you, what are the most important aspects of your background or identity?		

Education and Work History

How far did you go in school?	EDUCATION: 1 Grade 6 or less 2 Grades 7 to 12 (without graduating high school) 3 Graduated high school or high school equivalent 4 Part college/trade school 5 Graduated 2-year college or trade school 6 Graduated 4-year college 7 Part graduate/professional school	ONPS
IF FAILED TO COMPLETE A PROGRAM IN WHICH THEY WERE ENROLLED: Why did you leave?	8 Completed graduate/professional school	
What kind of work do you do? (Do you work outside of your home?)	·	
Education and Work History (continu	ed)	
Have you always done that kind of work?		
$\it IF~NO:~$ What other kind of work have you done in the past?		
What's the longest you've worked at one place?		
Are you currently employed (getting paid)?	PRIMARY EMPLOYMENT STATUS:	ONP
▶ F YES: Do you work part-time or full-time?	1 Full-time job2 Part-time job	ON
IF PART-TIME: How many hours do you typically work each week? (Why do you work part-time instead of full-time?)	 3 Keeping house or care giving full-time 4 In school/training 5 Retired 6 Unemployed, looking for work 	
→ F NO: Why is that? When was the last time you worked? How are you supporting yourself now?	7 Unemployed, not looking for work 8 Disabled	
IF DISABLED: Are you currently receiving disability payments? What are you receiving disability for?		
IF EMPLOYED: How long have you worked at your current job?		
IF LESS THAN 6 MONTHS: Why did you leave your last job?		
IF UNKNOWN: Has there ever been a period of time when you were unable to work or go to school?		
IF YES: Why was that?		
Have you ever been arrested, involved in a lawsuit, or had other legal trouble?		
	,	

Current and Past Periods of Psychopathology

NOTE: FOR A COMPLICATED HX, USE THE LIFE CHART ON PAGE 7. Have you ever seen anybody for emotional or psychiatric problems? ► IF YES: What was that for? (What treatment did you get? Any medications? When was that? When was the first time you ever saw someone for emotional or psychiatric problems?) ► IF NO: Was there ever a time when you, or someone else, thought you should see someone because of the way you were feeling or acting? (Tell me more.) Have you ever seen anybody for problems with alcohol or drugs? IF YES: What was that for? (What treatment[s] did you get? Any medications? When was that?) Have you ever attended a self-help group, like Alcoholics Anonymous, Gamblers Anonymous, or **Overeaters Anonymous?** IF YES: What was that for? When was that? **Hospitalization History** Have you ever been a patient in a psychiatric hospital? Number of previous hospitalizations (Do not include transfers): IF YES: What was that for? (How many times?) ONP7 IF AN INADEOUATE ANSWER IS GIVEN, CHALLENGE GENTLY: e.g., Wasn't there something else? People don't usually go to psychiatric hospitals just because they are tired or nervous. Have you ever been in a hospital for treatment of a medical problem? IF YES: What was that for? Thinking back over your whole life, when were you the most upset? (Why? What was that like? How were you feeling?)

Suicidal Ideation and Behavior

CHECK FOR THOUGHTS: In the past year, have you wished you were dead or wished you could go to sleep and not wake up? (Tell me about that.)	RECORD ANY HISTORY OF SUICIDAL THOUGHTS OR BEHAVIORS, INCLUDING IN THE PAST WEEK:	
► IF NO: SKIP TO NEXT PAGE, *SUICIDE ATTEMPT*		
IF YES: Did you have any of these thoughts in the past week (including today)?		
►IF NO: SKIP TO NEXT PAGE, *SUICIDE ATTEMPT*		
→ IF YES: CHECK FOR INTENT: Have you had a strong urge to kill yourself at any point during the past week? (Tell me about that.) In the past week, did you have any intention of attempting suicide? (Tell me about that.)		
CHECK FOR PLAN AND METHOD: In the past week, have you thought about how you might actually do it? (Tell me about what you were thinking of doing.) Have you thought about what you would need to do to carry this out? (Tell me about that. Do you have the means to do this?)	d	
Do you have the means to do this?)	Check if:	ONDO
	Suicidal Ideation past week with suicide intent	ONP9
	with suicide plan	ONP10 ONP11
	with suicide plan with access to chosen method	ONP11 ONP12

Suicide Attempt

CHECK FOR ATTEMPT: In the past year, have you tried to kill yourself?		
$\it IF~NO:~$ In the past year, have you done anything to harm yourself?		
<pre>IF NO: GO TO *OTHER CURRENT PROBLEMS,* BELOW.</pre>		
IF YES TO EITHER OF ABOVE: What did you do? (Tell me what happened.) Were you trying to end your life?		
IF MORE THAN ONE ATTEMPT: Which attempt had the most severe medical consequences (going to emergency department, needing hospitalization, requiring ICU)?		
Have you made any suicide attempts in the past week (including today)?	Check if: Suicide attempt past week	ONP1
Other Current Problems		
Have you had any other problems in the past year? (How are things going at work, at home, and with other people?)		
What has your mood been like?		
How has your physical health been? (Have you had any medical problems?)		
Do you take any medication, vitamins, nutritional supplements, or natural health remedies (other than those you've already told me about?)		
IF YES: How much and how often do you take (MEDICATION)? (Has there been any change in the amount you have been taking?)		
How have you been spending your free time? Who do you spend time with?		

GO TO NEXT MODULE

THE LIFE CHART (BELOW) MAY BE USED AT ANY POINT IN THE OVERVIEW TO RECORD THE DETAILS OF A COMPLICATED HISTORY.

LIFE CHART

Age (or date)	Description (symptoms, triggering events)	Treatment
		_

RETURN TO OVERVIEW PAGE 3, *HOSPITALIZATION HISTORY* TO CONTINUE WITH OVERVIEW QUESTIONS.

NO

CIRCLE

"NO" ON

F.1

NO

CIRCLE

"NO" ON

F.8

SCID Screening Module (including optional disorders)

Now I want to ask you some more specific questions about problems you may have had. We'll go into more detail about them later.

1. In the past year, since (1 YEAR AGO), have you had an intense rush of anxiety, or what someone might call a "panic attack," when you <u>suddenly</u> felt very frightened, or anxious or <u>suddenly</u> developed a lot of physical symptoms?

(screening for panic attacks)

2. In the past year, since (1 YEAR AGO), have you been very anxious about or afraid of situations like going out of the house alone, being in crowds, going to stores, standing in lines, or traveling on buses or trains?

(screening for Agoraphobia)

3. In the past year, since (1 YEAR AGO), have you been especially nervous or anxious in social situations like having a conversation or meeting unfamiliar people?

(screening for Social Anxiety Disorder)

NO YES S3

CIRCLE
"NO" ON 1st ITEM, F.14

S3

CIRCLE
"YES" ON 1st ITEM, F.14

S1

S2

YES

CIRCLE

"YES" ON

F.1

YES

CIRCLE

"YES" ON

F.8

4. In the past year, since (1 YEAR AGO), was there anything that you have been afraid to do or felt very uncomfortable doing in front of other people, like speaking, eating, writing, or using a public bathroom?

(screening for Social Anxiety Disorder)

CIRCLE
"NO" ON
2nd ITEM,
F.14

S4

CIRCLE
"YES" ON
2nd ITEM,
F.14

5. Are there any other things since (1 YEAR AGO) that have made you especially anxious or afraid, like flying, seeing blood, getting a shot, heights, closed places, or certain kinds of animals or insects?

(screening for Specific Phobia)

NO YES 55

| CIRCLE "NO" ON F.19 F.19

NO

YES

S6

6. Over the last 12 months have you been feeling anxious and worried for a lot of the time?


(screening for current Generalized Anxiety Disorder)

8. In the past year, since (1 YEAR AGO), have you been bothered with thoughts that kept coming back to you even when you didn't want them to, like being exposed to germs or dirt or needing everything to be lined up in a certain way?

(screening for obsessions in Obsessive-Compulsive Disorder)

9. How about having images pop into your head that you didn't want like violent or horrible scenes or something of a sexual nature?

(screening for obsessions in Obsessive-Compulsive Disorder)



10. How about having urges to do something that kept coming back to you even though you didn't want them to, like an urge to harm a loved one?

(screening for obsessions in Obsessive-Compulsive Disorder)

NO YES S10

CIRCLE "NO" ON 3rd ITEM, G.1 G.1

 $11.\,$ In the past year, since $(1\ \text{YEAR AGO})$, was there anything that you had to do over and over again and was hard to resist doing, like washing your hands again and again, repeating something over and over again until it "felt right," counting up to a certain number, or checking something many times to make sure that you'd done it right?

(screening for compulsions in Obsessive-Compulsive Disorder)


NO YES S11

CIRCLE CIRCLE "YES" ON G.2

G.2


12. In the past year, since (1 YEAR AGO), have you had a time when you weighed much less than other people thought you ought to weigh?

(screening for Anorexia Nervosa)




13. Have you had eating binges in the past year, that is, times when you couldn't resist eating a lot of food or stop eating once you started?

(screening for binge eating in Bulimia Nervosa and Binge Eating Disorder)




15a. In the past year, since (1 YEAR AGO), have you frequently lost control of your temper and ended up yelling or getting into arguments with others?

(screening for current Intermittent Explosive Disorder)



15b. In the past year, since (1 YEAR AGO), have you lost your temper so that you shoved, hit, kicked, or threw something at a person or an animal, or damaged someone's property?

(screening for current Intermittent Explosive Disorder)



A. MOOD EPISODES

*CURRENT MAJOR DEPRESSIVE MAJOR DEPRESSIVE EPISODE **EPISODE* CRITERIA**

Now I am going to ask you some more questions about your mood.

A. Five (or more) of the following symptoms have been present during the same 2-week period and represent a change from previous functioning; at least one of the symptoms is either (1) depressed mood, or (2) loss of interest or pleasure.

Since (1 YEAR AGO), has there been a period of time when you were feeling depressed or down most of the day nearly every day? (Has anyone said that you look sad, down, or depressed?)

IF NO: What about feeling empty or hopeless most of the day nearly every day?


IF YES TO EITHER OF ABOVE: What has that been like? How long has it lasted? (As long as 2 weeks?)

- ►IF PREVIOUS ITEM CODED "3:" During that time, did you lose interest or pleasure in things you usually enjoyed? (What has that been like? Give me some examples.)
 - ► IF PREVIOUS ITEM NOT CODED "3:" What about a time since (1 YEAR AGO) when you lost interest or pleasure in things you usually enjoyed? (What has that been like? Give me some examples.)

IF YES: Has it been nearly every day? How long has it lasted? (As long as 2 weeks?)

1. Depressed mood most of the day, nearly every day, as indicated either by subjective report (e.g., feels sad, empty, hopeless) or observation made by others (e.g., appears tearful). NOTE: in children or adolescents, can be irritable mood.

2. Markedly diminished interest or pleasure in all, or almost all, activities most of the day, nearly every day (as indicated either by subjective account or observation).



FOR THE FOLLOWING QUESTIONS, FOCUS (OR ELSE THE PAST 2 WEEKS IF EQUALLY DEPRESSED FOR ENTIRE YEAR).

NOTE: When rating the following items, code "1" if ON THE WORST 2 WEEKS IN THE PAST YEAR the symptoms are clearly due to a general medical condition (e.g., insomnia due to severe back pain).

IF UNKNOWN: Since (1 YEAR AGO), during which 2-week period would you say you have been doing the worst?

SCID-RV (for DSM-5®) (Version 1.0.0)	Current MDE Mood E	pisodes w/o Specifie	ers		
During (2-WEEK PERIOD)					
how has your appetite been? (What about compared to your usual appetite? Have you had to force yourself to eat? Eat [less/more] than usual? Has that been nearly every day? Have you lost or gained any weight? How much?	 Significant weight loss when not dietir weight gain (e.g., a change of more the of body weight in a month) or decrease increase in appetite nearly every day. in children, consider failure to make e weight gains. Check if:	han 5% se or NOTE:	2	3	A3
IF YES: Have you been trying to [lose/gain] weight?)	weight loss or decreased appetite weight gain or increased appetite				A5
how have you been sleeping? (Trouble falling asleep, waking frequently, trouble staying asleep, waking too early, OR sleeping too	Insomnia or hypersomnia nearly every Check if: insomnia	y day. ? 1	2	3	A6
much? How many hours of sleep [including naps] have you been getting? How many hours of sleep did you typically get before you got [depressed/OWN WORDS]? Has it been nearly every night?)	hypersomnia				A7 A8
have you been so fidgety or restless that you were unable to sit still? What about the opposite—talking more slowly, or moving more slowly than is normal for you, as if you're moving through molasses or mud? (In either	 Psychomotor agitation or retardation is every day (observable by others, not subjective feelings of restlessness or is slowed down) NOTE: Consider behavior during the intercolor of the considering the subjective feelings of restlessness or is slowed down. 	merely being	2	3	A9
instance, has it been so bad that other people have noticed it? What have they noticed? Has that been nearly every day?)	Check if: psychomotor agitation psychomotor retardation				A10 A11
what has your energy level been like? (Tired all the time? Nearly every day?)	6. Fatigue or loss of energy nearly every	day. ? 1	2	3	A12
have you been feeling worthless? What about feeling guilty about things you have done or not done?	 Feelings of worthlessness or excessive inappropriate guilt (which may be delu- nearly every day (not merely self-repr- guilt about being sick) 	usional)	2	3	A13
IF YES: What things? (Is this only because you can't take care of things since you have been sick?)	Check if: worthlessness inappropriate guilt				A14 A15
IF YES TO EITHER OF ABOVE: Nearly every day?					
have you had trouble thinking or concentrating? Has it been hard to make decisions about everyday things? (What kinds of things has it been interfering with? Nearly every day?)	Diminished ability to think or concentred indecisiveness, nearly every day (either subjective account or as observed by a subjective account or as observed by a subjective account or as observed.	er by	2	3	A16

...have things been so bad that you thought a lot about death or that you would be better off dead? Have you thought about taking your own life?

IF YES: Have you done something about it? (What have you done? Have you made a specific plan? Have you taken any action to prepare for it? Have you actually made a suicide attempt?)

9. Recurrent thoughts of death (not just fear of dving), recurrent suicidal ideation without a specific plan, or a suicide attempt or a specific plan for committing suicide.

NOTE: Code "1" for self-mutilation without suicidal intent.

Check if:

suicide attempt

___ thoughts of own death A19 A20 suicidal ideation A21 _ specific plan

NOTE: Any current suicidal thoughts, plans, or actions should be thoroughly assessed by the clinician and action taken if necessary.

AT LEAST FIVE OF THE ABOVE SXS (A.1-A.9) ARE CODED "3" AND AT LEAST ONE OF THESE IS ITEM A.1 OR A.2.

B. The symptoms cause clinically significant distress

or impairment in social, occupational, or other

important areas of functioning.

A22 3 GO TO *CURRENT **MANIC EPISODE*** A23 ? 2 3 GO TO *CURRENT MANIC EPISODE*

1

2

3

A17

A18

IF UNKNOWN: What effect have (DEPRESSIVE SXS) had on your life?

ASK THE FOLLOWING QUESTIONS AS <u>NEEDED</u> TO RATE CRITERION B:

How have (DEPRESSIVE SXS) affected your relationships or your interactions with other people? (Has this caused you any problems in your relationships with your family, romantic partner or friends?)

How have (DEPRESSIVE SXS) affected your work/school? (How about your attendance at work or school? Did [DEPRESSIVE SXS] make it more difficult to do your work/schoolwork? How have [DEPRESSIVE SXS] affected the quality of your work/schoolwork?)

How have (DEPRESSIVE SXS) affected your ability to take care of things at home? How about doing simple everyday things like getting dressed, bathing, or brushing your teeth? What about doing other things that are important to you like religious activities, physical exercise, or hobbies? Have you avoided doing anything because you felt like you weren't up to

Have (DEPRESSIVE SXS) affected any other important part of your life?

IF DOES NOT INTERFERE WITH LIFE: How much have you been bothered or upset by having (DEPRESSIVE SXS)?

IF UNKNOWN: When did (EPISODE OF DEPRESSION) begin?

Just before this began, were you physically ill?

IF YES: What did the doctor say?

Just before this began, were you using any medications?

IF YES: Any change in the amount you were using?

Just before this began, were you drinking or using any drugs?


C. [Primary Depressive Episode:] The episode is not attributable to the physiological effects of a substance (e.g., a drug of abuse, medication) or to another medical condition.


IF THERE IS ANY INDICATION THAT THE DEPRESSION MAY BE SECONDARY (I.E., A DIRECT PHYSIOLOGICAL CONSEQUENCE OF GMC OR SUBSTANCE/MEDICATION), GO TO *GMC/SUBSTANCE* A.45, AND RETURN HERE TO MAKE A RATING OF "1" OR "3."

<u>Etiological medical conditions include:</u> stroke, Huntington's disease, Parkinson's disease, traumatic brain injury, Cushing's disease, hypothyroidism, multiple sclerosis, systemic lupus erythematosus.

Etiological substances/medications include: alcohol (I/W), phencyclidine (I), hallucinogens (I), inhalants (I), opioids (I/W), sedative, hypnotics or anxiolytics (I/W), amphetamine and other stimulants (I/W), cocaine (I/W), antiviral agents (etavirenz), cardiovascular agents (clonodine, guanethidine, methyldopa, reserpine), retinoic acid derivatives (isotretinoin), antidepressants, anticonvulsants, anti-migraine agents (triptans), antipsychotics, hormonal agents (corticosteroids, oral contraceptives, gonadotropin-releasing hormone agonists, tamoxifen), smoking cessation agents (varenicline) and immunological agents (interferon).

MAJOR DEPRESSIVE EPISODE CRITERIA A, B, AND C ARE CODED "3."





1

GO TO *PAST

GO TO *PAST

MANIC

EPISODE*

MANIC

EPISODE*

2

3

3

CURRENT MANIC EPISODE

MANIC EPISODE CRITERIA

Since (1 YEAR AGO), has there been a period of time when you were feeling so good, "high," excited, or "on top of the world" that other people thought you were not your normal self?

► IF YES: What has it been like? (More than just feeling good?)

Have you also been feeling like you were "hyper" or "wired" and had an unusual amount of energy? Have you been much more active than is typical for you? (Have other people commented on how much you have been doing?)

► IF NO: Since (1 YEAR AGO), have you had a period of time when you were feeling irritable, angry, or shorttempered most of the day, nearly every day, for at least several days? What has it been like? (Is that different from the way you usually are?)

> IF YES: Have you also been feeling like you were "hyper" and had an unusual amount of energy? Have you been much more active than is typical for you? (Have other people commented on how much you have been doing?)

How long has this lasted? (As long as 1 week?)

IF LESS THAN 1 WEEK: Did you need to go into the hospital to protect you from hurting yourself or someone else, or from doing something that could have caused serious financial or legal problems?

Have you been feeling (high/irritable/OWN WORDS) for most of the day, nearly every day during this time?

FOCUS ON THE MOST SEVERE WEEK IN THE PAST YEAR OF THE CURRENT EPISODE FOR THE FOLLOWING QUESTIONS.

IF UNCLEAR: During (EPISODE), when were you the most (high/irritable/OWN WORDS)?

During that time...

...how did you feel about yourself?

(More self-confident than usual? Did you feel much smarter or better than everyone else? Did you feel like you had any special powers or abilities?)

...did you need less sleep than usual? (How much sleep did you get?)

IF YES: Did you still feel rested?

A. A distinct period [lasting at least several days] of abnormally and persistently elevated, expansive, or irritable mood and abnormally and persistently increased [...] activity or energy.

Check if:

elevated, expansive mood

irritable mood

...lasting at least 1 week and present most of the day, nearly every day (or any duration if hospitalization is necessary).

NOTE: If elevated mood lasts less than 1 week, check whether irritable mood lasts at least 1 week before skipping to A.14.

B. During the period of mood disturbance and increased energy or activity, three (or more) of the following symptoms have persisted (four if the mood is only irritable) and have been present to a significant degree and represent a noticeable change from usual

1. Inflated self-esteem or grandiosity.

2. Decreased need for sleep (e.g., feels rested after only 3 hours of sleep).

2

2

3

A59

A58

A54

A55

A56

A57

?=inadequate information

1=absent or false

hehavior:

2=subthreshold

3=threshold or true

...were you doing anything that could have caused trouble for you or your family?

(Spending money on things you didn't need or couldn't afford? How about giving away money or valuable things? Gambling with money you couldn't afford to lose?)

(Anything sexual that was likely to get you in trouble? Driving recklessly?)

(Did you make any risky or impulsive business investments or get involved in a business scheme that you wouldn't normally have done?) Excessive involvement in activities that have a high potential for painful consequences (e.g., engaging in unrestrained buying sprees, sexual indiscretions, or foolish business investments). A66

GO TO *PAST

MANIC

EPISODE*

AT LEAST THREE "B" SXS ARE CODED "3" (FOUR IF MOOD ONLY IRRITABLE).

1 3 A67
GO TO *PAST
MANIC
EPISODE*

? 1 2 3 A68

IF UNKNOWN: What effect have these (MANIC SXS) had on your life?

IF UNKNOWN: Have you needed to go into the hospital to protect you from hurting yourself or someone else, or from doing something that could have caused serious financial or legal problems?

ASK THE FOLLOWING QUESTIONS <u>AS NEEDED</u> TO RATE CRITERION C.

How have (MANIC SXS) affected your relationships or your interactions with other people? (Have (MANIC SXS) caused you any problems in your relationships with your family, romantic partner or friends?)

How have (MANIC SXS) affected your work/ school? (How about your attendance at work or school? Did [MANIC SXS] make it more difficult to do your work/ schoolwork? How have [MANIC SXS] affected the quality of your work/ schoolwork?)

How have (MANIC SXS) affected your ability to take care of things at home?

C. The mood disturbance is sufficiently severe to cause marked impairment in social or occupational functioning or to necessitate hospitalization to prevent harm to self or others, or there are psychotic features.

NOTE: Code "3" if psychotic symptoms have been present. You may need to return here to recode after screening for psychotic symptoms in Module B.

DESCRIBE:

IF UNKNOWN: When did this period of being (high/irritable/OWN WORDS) begin?

Just before this began, were you physically ill?

IF YES: What did the doctor say?

Just before this began, were you taking any medications?

IF YES: Any change in the amount you were taking?

Just before this began, were you drinking or using any drugs?

D. [Primary Manic Episode:] The episode is not attributable to the physiological effects of a substance (i.e., a drug of abuse, medication) or to another medical condition.

IF THERE IS ANY INDICATION THAT MANIA MAY BE SECONDARY (I.E., A DIRECT PHYSIOLOGICAL CONSEQUENCE OF A GMC OR SUBSTANCE), GO TO *GMC/SUBSTANCE* A.41 AND RETURN HERE TO MAKE A RATING OF "1" OR "3."

NOTE: A full Manic Episode that emerges during antidepressant treatment (e.g., medication, electroconvulsive therapy) but persists at a fully syndromal level beyond the physiological effect of that treatment is sufficient evidence for a Manic Episode and, therefore, a Bipolar I diagnosis.

Etiological medical conditions include: Alzheimer's disease, vascular dementia, HIV-induced dementia, Huntington's disease, Lewy body disease, Wernicke-Korsakoff, Cushing's disease, multiple sclerosis, ALS, Parkinson's disease, Pick's disease, Creutzfelt-Jakob disease, stroke, traumatic brain injuries, hyperthyroidism

Etiological substances/medications include: alcohol (I/W), phencyclidine (I), hallucinogens (I), sedatives, hypnotics, anxiolytics (I/W), amphetamines (I/W), cocaine (I/W), corticosteroids, androgens, isoniazid, levodopa, interferon alpha, varenicline, procarbazine, clarithromycin, ciprofloxacin

MANIC EPISODE CRITERIA A, B, C, AND D ARE CODED "3."

PRIMARY
MANIC
EPISODE

DUE TO
SUBSTANCE
USE OR GMC,
GO TO *PAST
MANIC
EPISODE*

CONTINUE
WITH NEXT
ITEM

A69

CURRENT MANIC EPISODE

1

A70

PAST MANIC EPISODE

NOTE: IF CURRENTLY ELEVATED OR IRRITABLE MOOD BUT FULL CRITERIA ARE NOT MET FOR A MANIC EPISODE, SUBSTITUTE THE PHRASE "Has there ever been <u>another</u> time ..." IN EACH OF THE SCREENING QUESTIONS BELOW.

Have you <u>ever</u> had a period of time when you were feeling so good, "high," excited, or "on top of the world" that other people thought you were not your normal self?

► IF YES: What was it like? (Was that more than just feeling good?) Did you also feel like you were "hyper" or "wired" and had an unusual amount of energy? Were you much more active than is typical for you? (Did other people comment on how much you were doing?)

▶ IF NO: Have you ever had a period of time when you were feeling irritable, angry, or short-tempered for most of the day, every day, for at least several days? What was that like? (Was that different from the way you usually are?)

IF YES: Did you also feel like you were "hyper" or "wired" and had an unusual amount of energy? Were you much more active than is typical for you? (Did other people comment on how much you were doing?)

MANIC EPISODE CRITERIA


IF CURRENT MAJOR DEPRESSIVE
EPISODE HAS BEEN CONFIRMED, GO TO
*CURRENT PERSISTENT DEPRESSIVE
DISORDER*

A. A distinct period [lasting at least several days] of abnormally and persistently elevated, expansive or irritable mood and abnormally and persistently increased [...] activity or energy.

Check if:

____ elevated, expansive mood

___ irritable mood



When was that?

How long did that last? (As long as 1 week?)

IF LESS THAN 1 WEEK: Did you need to go into the hospital to protect you from hurting yourself or someone else, or from doing something that could have caused serious financial or legal problems?)

Did you feel (high/irritable/OWN WORDS) for most of the day, nearly every day during this time?


Have you had more than one time like that? (Which time was the most extreme?)

IF UNCLEAR: Have you had any times like that in the past year, since (1 YEAR AGO)?

...lasting at least 1 week and present most of the day, nearly every day (or any duration if hospitalization is necessary).

NOTE: If elevated mood lasts less than 1 week, check whether irritable mood lasts at least 1 week before skipping to A.23.

NOTE: If there is evidence for more than one past episode, select the worst episode that occurred in the prior year; if none of the past episodes occurred in the prior year, select the worst episode that occurred regardless of the time it occurred.



EPISODE THAT YOU ARE INQUIRING ABOUT.	i t	During the period of mood disturbance and ncreased energy or activity, three (or more) of the following symptoms have persisted (four if the mood is only irritable) and have been					
<pre>IF UNCLEAR: During (EPISODE), when were you the most (high/irritable/OWN WORDS)?</pre>	ŗ	present to a significant degree and represent a noticeable change from usual behavior:					
During that time							
how did you feel about yourself? (More self-confident than usual? Did you feel much smarter or better than everyone else? Did you feel like you had any special powers or abilities?)	1	. Inflated self-esteem or grandiosity.	?	1	2	3	A96
did you need less sleep than usual? (How much sleep did you get?)	2	 Decreased need for sleep (e.g., feels rested after only 3 hours of sleep). 	?	1	2	3	A97
IF YES: Did you still feel rested?							
were you much more talkative than usual? (Did people have trouble stopping you or understanding you? Did people have trouble getting a word in edgewise?)	3	 More talkative than usual or pressure to keep talking. 	?	1	2	3	A98
did you have thoughts racing through your head? (What was that like?)	4	 Flight of ideas or subjective experience that thoughts are racing. 	?	1	2	3	A99
were you so easily distracted by things around you that you had trouble concentrating or staying on one track? (Give me an example of that.)	5	 Distractibility (i.e., attention too easily drawn to unimportant or irrelevant external stimuli) as reported or observed. 	?	1	2	3	A100
how did you spend your time? (Work, friends, hobbies? Were you especially busy during that time?)	6	 Increase in goal-directed activity (either socially, at work or school, or sexually) or psychomotor agitation (i.e., purposeless non-goal-directed activity). 	?	1	2	3	A101
(Did you find yourself more enthusiastic at work or working harder at your job?	c	Check if:					
Did you find yourself more engaged in		increase in activity					A102 A103
school activities or studying harder?)		psychomotor agitation					
(Were you more sociable during that time, such as calling on friends or going out with them more than you usually do or making a lot of new friends?)							
(Were you spending more time thinking about sex or involved in doing something sexual, by yourself or with others? Was that a big change for you?)							
Were you physically restless during this time, doing things like pacing a lot, or being unable to sit still? (How bad was it?)							
•							

A104


During that time...

...did you do anything that could have caused trouble for you or your family?

(Spending money on things you didn't need or couldn't afford? How about giving away money or valuable things? Gambling with money you couldn't afford to lose?)

(Anything sexual that was likely to get you in trouble? Driving recklessly?)

(Did you make any risky or impulsive business investments or get involved in a business scheme that you wouldn't normally have done?) 7. Excessive involvement in activities which have ? 1 2 3 a high potential for painful consequences (e.g., engaging in unrestrained buying sprees, sexual indiscretions, or foolish business investments)



IF UNKNOWN: What effect did these (MANIC SXS) have on your life?

IF UNKNOWN: Did you need to go into the hospital to protect you from hurting yourself or someone else, or from doing something that could have caused serious financial or legal problems?

ASK THE FOLLOWING QUESTIONS <u>AS</u> <u>NEEDED</u> TO RATE CRITERION C.

How did (MANIC SXS) affect your relationships or your interactions with other people? (Did (MANIC SXS) cause you any problems in your relationships with your family, romantic partner or friends?)

How did (MANIC SXS) affect your work/school? (How about your attendance at work or school? Did [MANIC SXS] make it more difficult to do your work/schoolwork? How did [MANIC SXS] affect the quality of your work/schoolwork?)

How did (MANIC SXS) affect your ability to take care of things at home?

C. The mood disturbance is sufficiently severe to cause marked impairment in social or occupational functioning or to necessitate hospitalization to prevent harm to self or others or there are psychotic features. ? 1 2

A106

3

CONTINUE ON NEXT PAGE

GO TO *CURRENT PERSISTENT
DEPRESSIVE DISORDER*

1

IF UNKNOWN: When did this period of being (high/irritable/OWN WORDS) begin?

Just before this began, were you physically ill?

IF YES: What did the doctor say?

Just before this began, were you taking any medications?

IF YES: Any change in the amount you were taking?

Just before this began, were you drinking or using any drugs?

D. [Primary Manic Episode:] The episode is not attributable to the physiological effects of a substance (e.g., a drug of abuse, medication) or to another medical condition.

IF THERE IS ANY INDICATION THAT THE MANIA MAY BE SECONDARY (I.E., A DIRECT PHYSIOLOGICAL CONSEQUENCE OF GMC OR SUBSTANCE), GO TO *GMC/SUBSTANCE* A.41, AND RETURN HERE TO MAKE A RATING OF "1" OR "3."

NOTE: A full Manic Episode that emerges during antidepressant treatment (e.g., medication, electroconvulsive therapy) but persists at a fully syndromal level beyond the physiological effect of that treatment is sufficient evidence for a Manic Episode and, therefore a Bipolar I diagnosis.

NOTE: Refer to lists of etiological medical conditions and substances/medications on page A.13.

DUE TO SUBSTANCE USE OR GMC PRIMARY MANIC EPISODE

G

CONTINUE WITH NEXT ITEM

E

1

3

A108

A107

GO TO *CURRENT PERSISTENT
DEPRESSIVE DISORDER*

MANIC EPISODE CRITERIA A, B, C, AND D ARE CODED "3."

GO TO *CURRENT
PERSISTENT
DEPRESSIVE
DISORDER*

PAST MANIC EPISODE

SCID-RV (for DSM-5®) (Version 1.0.0)Current Persistent Depressive Disorder Mood Episodes A				
?=inadequate information	1=absent or false	2=subthreshold	3=threshold or true	

CURRENT PERSISTENT DEPRESSIVE DISORDER

CURRENT PERSISTENT DEPRESSIVE DISORDER CRITERIA

IF THERE HAS EVER BEEN A MANIC OR HYPOMANIC EPISODE, CHECK HERE AND GO TO THE NEXT MODULE A139 Since (1 YEAR AGO), have you been A. Depressed mood for most of the day, for more ? 2 3 A140 1 bothered by depressed mood most of days than not, as indicated either by subjective the day, more days than not? (More account or observation by others, for at least 2 than half of the time?) years. NOTE: in adolescents, mood can be GO TO THE irritable and duration must be at least 1 year. **NEXT MODULE** IF YES: What has that been like? During these periods of (OWN WORDS B. Presence, while depressed, of two (or more) of FOR CHRONIC DEPRESSION) did you the following: often... A141 ...lose your appetite? (What about 2 3 1 1. Poor appetite or overeating. overeating?) A142 ...have trouble sleeping or sleep too 2 3 1 2. Insomnia or hypersomnia. much? A143 ...have little energy to do things or feel 3. Low energy or fatigue. 2 3 tired a lot? A144 ...feel down on yourself? (Feel 3 2 4. Low self-esteem. 1 worthless, or a failure?) A145 3 ...have trouble concentrating or making 2 5. Poor concentration or difficulty making 1 decisions? decisions. A146 ...feel hopeless? 1 2 3 6. Feelings of hopelessness. A147 AT LEAST TWO "B" SYMPTOMS ARE CODED "3." 1 2 3 GO TO THE NEXT MODULE Since (1 YEAR AGO), what was the A148 C. During the 1-year period of the disturbance, the 3 1 longest period of time that you felt OK individual has never been without the symptoms (NO DYSTHYMIC SYMPTOMS)? in Criteria A and B for more than 2 months at a GO TO THE time NEXT MODULE NOTE: Code "1" if normal mood for more than 2 months at a time. A149 E. There has never been a Manic Episode or a 3 1 Hypomanic Episode, and criteria have never been met for Cyclothymic disorder. GO TO THE **NEXT MODULE**

IF NOT ALREADY CLEAR, RETURN TO THIS ITEM AFTER COMPLETING THE PSYCHOTIC DISORDERS SECTION.

F. The disturbance is not better explained by a persistent Schizoaffective Disorder, Schizophrenia, Delusional Disorder, or Other Specified or Unspecified Schizophrenia Spectrum or Other Psychotic Disorder. 1 3 A150

GO TO THE
NEXT MODULE

NOTE: Code "3" if NO chronic psychotic disorder has been present or if NOT better explained by a chronic psychotic disorder.

IF UNKNOWN: When did this begin?

Just before this began, were you physically ill?

IF YES: What did the doctor say?

Just before this began, were you using any medications?

IF YES: Any change in the amount you
were using?

Just before this began, were you drinking or using any drugs?

G. [Primary Persistent Depressive Disorder:] The symptoms are not attributable to the physiological effects of a substance (e.g., a drug of abuse, medication) or to another medical condition (e.g., hypothyroidism).

IF THERE IS ANY INDICATION THAT THE DEPRESSION MAY BE SECONDARY (I.E., A DIRECT PHYSIOLOGICAL CONSEQUENCE OF GMC OR SUBSTANCE), GO TO *GMC/SUBSTANCE/MEDICATION* A.45, AND RETURN HERE TO MAKE A RATING OF "1" OR "3."

NOTE: Refer to lists of etiological medical conditions and substances/medications on page A.4.

? 1 3 A151
PRIMARY
DEPRESSIVE
DISORDER

GO TO THE NEXT
MODULE

CONTINUE
WITH NEXT
ITEM

2

3

?

1

A152

IF UNKNOWN: What effect have these (DEPRESSIVE SXS) had on your life?

ASK THE FOLLOWING QUESTIONS <u>AS</u>
<u>NEEDED</u> TO RATE CRITERION H:

How have (DEPRESSIVE SXS) affected your relationships or your interactions with other people? (Has it caused you any problems in your relationships with your family, romantic partner or friends?)

How have these (DEPRESSIVE SXS) affected your work/school? (How about your attendance at work or school? Have [DEPRESSIVE SXS] made it more difficult to do your work/schoolwork? How did [DEPRESSIVE SXS] affect the quality of your work/schoolwork?)

How have (DEPRESSIVE SXS) affected your ability to take care of things at home? How about doing simple everyday things like getting dressed, bathing, or brushing your teeth? How about doing other things that are important to you like religious activities, physical exercise, or hobbies? Did you avoid doing anything because you felt like you weren't up to it?

H. The symptoms cause clinically significant distress or impairment in social, occupational, or other important areas of functioning.

Have these (DEPRESSIVE SXS) affected any other important part of your life?

IF DOES NOT INTERFERE WITH LIFE: How much you been bothered or upset by having (DEPRESSIVE SXS)?

	PRESSIVE DISORDER CRITERIA A, AND H ARE CODED "3."	1	3 	A153
			CURRENT PERSISTEI DEPRESSI DISORDER	√E
1 - Ea	ate onset specifier: arly onset: onset before age 21 ate onset: onset age 21 or older			A154
Specify if (for mo	ost recent 2 years of Persistent De	pressive Disorder):		A155
	l information about onset and offse 2 years may be needed to evaluate		sive Episode	s
	dysthymic syndrome: Full criter we not been met in at least the pre		essive	
	stent Major Depressive Episode Episode have been met throughou			
Full criteria been perioc	mittent Major Depressive Episo for a Major Depressive Episode ar ds of at least 8 weeks in at least th below the threshold for a full Major	e currently met, bue preceding 2 year	ut there have s with	е
episode: F	mittent Major Depressive Episo full criteria for a Major Depressive I as been one or more Major Depres 2 years.	Episode are not cur	rently met,	
Specify if:				A156
		GO TO THE MODULE	NEXT	

IF UNKNOWN: Have there been any panic attacks in the past month?

GMC/SUBSTANCE CAUSING BIPOLAR AND RELATED SYMPTOMS


BIPOLAR AND RELATED DISORDER DUE BIPOLAR AND RELATED DISORDER DUE TO ANOTHER MEDICAL CONDITION ANOTHER MEDICAL CONDITION CRITERIA

IF SYMPTOMS NOT TEMPORALLY ASSOCIATED WITH A GENERAL MEDICAL CONDITION, CHECK HERE ___ AND GO TO *SUBSTANCE-INDUCED BIPOLAR AND RELATED DISORDER* A.43.

A193

CODE BASED ON INFORMATION ALREADY OBTAINED.

- A. A prominent and persistent period of abnormally elevated, expansive, or irritable mood and abnormally increased activity or energy that predominates in the clinical picture.
- ? 1 2 3 A194
- B/C. There is evidence from the history, physical examination, or laboratory findings that the disturbance is the direct physiological consequence of another medical condition and the disturbance is not better accounted for by another mental disorder.



Did the (BIPOLAR SXS) change after (GMC) began? Did (BIPOLAR SXS) start or get much worse only after (GMC) began? How long after (GMC) began did (BIPOLAR SXS) start or get much worse?

IF GMC HAS RESOLVED: Did the (BIPOLAR SXS) get better once the (GMC) got better?

NOTE: The following factors should be considered and, if present, support the conclusion that a general medical condition is etiologic to the bipolar symptoms.

- There is evidence from the literature of a wellestablished association between the general medical condition and the bipolar symptoms. (Refer to list of etiological medical conditions on page A.13.)
- 2) There is a close temporal relationship between the course of the bipolar symptoms and the course of the general medical condition.
- The bipolar symptoms are characterized by unusual presenting features (e.g., late age-atonset).
- The absence of alternative explanations (e.g., bipolar symptoms as a psychological reaction to the stress of being diagnosed with a general medical condition).

IF UNKNOWN: What effect have (BIPOLAR SXS) had on your life?

ASK THE FOLLOWING QUESTIONS <u>AS</u> <u>NEEDED</u> TO RATE CRITERION E:

How have (BIPOLAR SXS) affected your relationships or your interactions with other people? (Have they caused you any problems in your relationships with your family, romantic partner or friends?)


How have they affected your work/ school? (How about your attendance at work or school? Have they affected the quality of your work/schoolwork?)

How did (BIPOLAR SXS) affect your ability to take care of things at home? Did you need to go into the hospital to protect you from hurting yourself or someone else, or from doing something that could have caused serious financial or legal problems?


Have (BIPOLAR SXS) affected any other important part of your life?

IF HAVE NOT INTERFERED WITH LIFE: How much have (BIPOLAR SXS) bothered or upset you?

E. The disturbance causes clinically significant distress or impairment in social, occupational, or other important areas of functioning or necessitates hospitalization to prevent harm to self or others, or there are psychotic features.



NOTE: The D criterion (delirium rule-out) has been omitted.



*SUBSTANCE-/MEDICATION-INDUCED **BIPOLAR DISORDER***

SUBSTANCE-/MEDICATION-INDUCED **BIPOLAR DISORDER CRITERIA**

IF SYMPTOMS ARE NOT TEMPORALLY ASSOCIATED WITH SUBSTANCE/MEDICATION USE, AND RETURN TO EPISODE BEING EVALUATED, CONTINUING WITH THE ITEM FOLLOWING "SYMPTOMS ARE NOT ATTRIBUTABLE TO THE PHYSIOLOGICAL EFFECTS OF A SUBSTANCE OR ANOTHER MEDICAL CONDITION" (SEE PAGE NUMBERS IN BOX TO THE RIGHT).

PAGE TO RETURN TO IN **EPISODE BEING EVALUATED:**

Current Manic A.13 Past Manic A.22 A200

OBTAINED.

- CODE BASED ON INFORMATION ALREADY A. A prominent and persistent disturbance in mood that predominates in the clinical picture and is characterized by elevated, expansive, or irritable mood, with or without depressed mood, or markedly diminished interest or pleasure in all, or almost all activities.
- A201 2 3 1

IF UNKNOWN: When did the (BIPOLAR SXS) begin? Were you already using (SUBSTANCE/MEDICATION) or had you just stopped or cut down your use?

B. There is evidence from the history, physical examination, or laboratory findings of both (1) and (2):

A202 1 2 3 NOT SUBSTANCE-INDUCED, RETURN TO EPISODE BEING **EVALUATED**

IF UNKNOWN: How much (SUBSTANCE/ MEDICATION) were you using when you began to have (BIPOLAR SXS)?

- 1. The symptoms in criterion A developed during or soon after substance intoxication or withdrawal or exposure to a medication.
- capable of producing the symptoms in Criterion A. NOTE: Refer to list of etiological substances/medications on page A.13. ASK ANY OF THE FOLLOWING QUESTIONS C. The disturbance is NOT better accounted for by

AS NEEDED TO RULE OUT A NON-SUBSTANCE-INDUCED ETIOLOGY.

a bipolar or related disorder that is not substance-induced. Such evidence of an independent bipolar or related disorder could include the following:

2. The involved substance/medication is

NOTE: The following three statements constitute evidence that the bipolar symptoms are not substance-induced. Code "1" if any are true. Code

IF UNKNOWN: Which came first, the (SUBSTANCE/MEDICATION USE) or the (BIPOLAR SXS)?

IF UNKNOWN: Have you had a period of time when you stopped using (SUBSTANCE/MEDICATION)?

IF YES: After you stopped using (SUBSTANCE/MEDICATION) did the (BIPOLAR SXS) go away or get

IF YES: How long did it take for them to get better? Did they go away within a month of stopping?


IF UNKNOWN: Have you had any other episodes of (BIPOLAR SXS)?

IF YES: How many? Were you using (SUBSTANCE/MEDICATION) at those times?

1) The symptoms precede the onset of the substance/medication use;

"3" only if none are true.

- 2) The symptoms persist for a substantial period of time (e.g., about 1 month) after the cessation of acute withdrawal or severe intoxication; or
- 3) There is other evidence suggesting the existence of an independent non-substance/ medication-induced bipolar and related disorder (e.g., a history of recurrent non-substance/ medication-related episodes).



IF UNKNOWN: What effect have (BIPOLAR SXS) had on your life?

ASK THE FOLLOWING QUESTIONS <u>AS</u> <u>NEEDED</u> TO RATE CRITERION E:

friends?)

How have (BIPOLAR SXS) affected your relationships or your interactions with other people? (Have they caused you any problems in your relationships with your family, romantic partner, or

distress or impairment in social, occupational, or other important areas of functioning.

RETURN TO EPISODE BEING EVALUATED

A204

3

NOTE: The D criterion (delirium rule-out) has been omitted.


E. The symptoms cause clinically significant

How have (BIPOLAR SXS) affected your work/school? (How about your attendance at work or school? Have they affected the quality of your work/schoolwork?)

How did (BIPOLAR SXS) affect your ability to take care of things at home? Have you needed to go into the hospital to protect you from hurting yourself or someone else, or from doing something that could have caused serious financial or legal problems?

Have (BIPOLAR SXS) affected any other important part of your life?

IF HAVE NOT INTERFERED WITH LIFE: How much have (BIPOLAR SX) bothered or upset you?



GMC/SUBSTANCE CAUSING DEPRESSIVE SYMPTOMS

*DEPRESSIVE DISORDER DUE TO **ANOTHER MEDICAL CONDITION***

DEPRESSIVE DISORDER DUE TO ANOTHER MEDICAL CONDITION CRITERIA

IF SYMPTOMS NOT TEMPORALLY ASSOCIATED WITH A GENERAL MEDICAL CONDITION, CHECK HERE ____ AND GO TO *SUBSTANCE-INDUCED DEPRESSIVE DISORDER* A.48


A208

CODE BASED ON INFORMATION ALREADY A. OBTAINED.

A prominent and persistent period of depressed mood or markedly diminished interest or pleasure in all, or almost all, activities that predominates in the clinical picture.

A209 2 3 1

B./C. There is evidence from the history, physical examination, or laboratory findings that the disturbance is the direct physiological consequence of another medical condition and the disturbance is not better accounted for by another mental disorder.



(GMC) began? Did (DEPRESSIVE SXS) start or get much worse only after (GMC) began? How long after (GMC) began did (DEPRESSIVE SXS) start or get much worse?

IF GMC HAS RESOLVED: Did the (DEPRESSIVE SXS) get better once the (GMC) got better?

Did the (DEPRESSIVE SXS) change after NOTE: The following factors should be considered and, if present, support the conclusion that a general medical condition is etiologic to the depressive symptoms.

- 1) There is evidence from the literature of a wellestablished association between the general medical condition and the depressive symptoms. (Refer to list of etiological general medical conditions on page A.4.)
- 2) There is a close temporal relationship between the course of the depressive symptoms and the course of the general medical condition.
- 3) The depressive symptoms are characterized by unusual presenting features (e.g., late age-atonset).
- 4) The absence of alternative explanations (e.g., depressive symptoms as a psychological reaction to the stress of being diagnosed with a general medical condition).

IF UNKNOWN: What effect have (DEPRESSIVE SX) had on your life?

ASK THE FOLLOWING QUESTIONS <u>AS</u> <u>NEEDED</u> TO RATE CRITERION E:

How have (DEPRESSIVE SXS) affected your relationships or your interactions with other people? (Have they caused you any problems in your relationships with your family, romantic partner, or friends?)


How have (DEPRESSIVE SXS) affected your work/school? (How about your attendance at work or school? Have they affected the quality of your work/schoolwork?)

How have (DEPRESSIVE SXS) affected your ability to take care of things at home? How about doing simple everyday things like getting dressed, bathing, or brushing your teeth? How about doing other things that are important to you like religious activities, physical exercise, or hobbies? Did you avoid doing anything because you felt like you weren't up to it?


Have (DEPRESSIVE SXS) affected any other important part of your life?

IF HAVE NOT INTERFERED WITH LIFE: How much have (DEPRESSIVE SXS) bothered or upset you?

E. The disturbance causes clinically significant distress or impairment in social, occupational, or other important areas of functioning.



NOTE: The D criterion (delirium rule-out) has been omitted.



*SUBSTANCE-/MEDICATION-INDUCED **DEPRESSIVE DISORDER***

SUBSTANCE-/MEDICATION-INDUCED **DEPRESSIVE DISORDER CRITERIA**

IF SYMPTOMS NOT TEMPORALLY ASSOCIATED WITH SUBSTANCE/MEDICATION USE, CHECK AND RETURN TO EPISODE BEING EVALUATED, CONTINUING WITH THE ITEM FOLLOWING "SYMPTOMS ARE NOT ATTRIBUTABLE TO THE PHYSIOLOGICAL EFFECTS OF A SUBSTANCE OR ANOTHER MEDICAL CONDITION" (SEE PAGE NUMBERS IN BOX TO THE RIGHT).

A215 PAGE TO RETURN TO IN EPISODE BEING EVALUATED: Current MDE A.4 Current Persistent Depressive Disorder A.31


OBTAINED.

CODE BASED ON INFORMATION ALREADY A. A prominent and persistent disturbance in mood that predominates in the clinical picture and is characterized by depressed mood or markedly diminished interest or pleasure in all, or almost all, activities

2 3 A216 1

IF UNKNOWN: When did the (DEPRESSIVE SXS) begin? Were you already using (SUBSTANCE/MEDICATION) or had you just stopped or cut down your use?

B. There is evidence from the history, physical examination, or laboratory findings of both (1) and (2):



IF UNKNOWN: How much (SUBSTANCE/ MEDICATION) were you using when you began to have (DEPRESSIVE SXS)?

- 1. The symptoms in criterion A developed during or soon after substance intoxication or withdrawal or exposure to a medication
- 2. The involved substance/medication is capable of producing the symptoms in Criterion A. NOTE: refer to list of etiological substances/medications on page A.4.

NEEDED TO RULE OUT A NON-SUBSTANCE-INDUCED ETIOLOGY.

IF UNKNOWN: Which came first, the (SUBSTANCE/MEDICATION USE) or the (DEPRESSIVE SXS)?

IF UNKNOWN: Have you had a period of time when you stopped using (SUBSTANCE/MEDICATION)?

IF YES: After you stopped using (SUBSTANCE/MEDICATION) did the (DEPRESSIVE SXS) go away or get better?

IF YES: How long did it take for them to get better? Did they go away within a month of stopping?


IF UNKNOWN: Have you had any other episodes of (DEPRESSIVE SXS)?

IF YES: How many? Were you using (SUBSTANCE/MEDICATION) at those times?

ASK ANY OF THE FOLLOWING QUESTIONS AS C. The disturbance is NOT better accounted for by a depressive disorder that is not substanceinduced. Such evidence of an independent depressive disorder could include the following:

> NOTE: The following three statements constitute evidence that the depressive symptoms are not substance-induced. Code "1" if any are true. Code "3" only if none are true.

- 1) The symptoms precede the onset of the substance/medication use;
- 2) The symptoms persist for a substantial period of time (e.g., about 1 month) after the cessation of acute withdrawal or severe intoxication; or
- 3) There is other evidence suggesting the existence of an independent non-substance/ medication-induced depressive disorder (e.g., a history of recurrent non-substance/ medication-related episodes).




IF UNKNOWN: What effect have (DEPRESSIVE SXS) had on your life?

E. The symptoms cause clinically significant distress or impairment in social, occupational, or other important areas of functioning.

ASK THE FOLLOWING QUESTIONS <u>AS NEEDED</u> TO RATE CRITERION E:

NOTE: the D criterion (delirium rule-out) has been omitted.




How have (DEPRESSIVE SXS) affected your relationships or your interactions with other people? (Have they caused you any problems in your relationships with your family, romantic partner or friends?)

How have (DEPRESSIVE SXS) affected your work/school? (How about your attendance at work or school? Have they affected the quality of your work/schoolwork?)

How have (DEPRESSIVE SXS) affected your ability to take care of things at home? How about doing simple everyday things like getting dressed, bathing, or brushing your teeth? How about doing other things that are important to you like religious activities, physical exercise, or hobbies? Did you avoid doing anything because you felt like you weren't up to it?

Have (DEPRESSIVE SXS) affected any other important part of your life?

IF HAVE NOT INTERFERED WITH LIFE: How much have (DEPRESSIVE SXS) bothered or upset you?



B/C. PSYCHOTIC SCREENING MODULE

NOTE: This module is for coding psychotic and associated symptoms that have been present in the past year. It can be used for settings in which cases with primary psychotic symptoms are to be excluded i.e., psychotic symptoms that are not due to substance/medication use or to a general medical condition) and/or psychotic symptoms that occur outside the context of a Major Depressive or Manic Episode.

For each psychotic symptom coded "3," describe the actual content and indicate the period of time during which the symptom was present. Moreover, for any psychotic symptom coded "3." determine whether the symptom is definitely "primary" or whether there is a possible or definite etiological substance (including medication) or general medical condition. Refer to page B/C.6 for a list of possible etiological general medical conditions and substances/medications.

The following questions may be useful if the Overview has not already provided the information.

Just before (PSYCHOTIC SXS) began, were you using drugs? ...were you taking any medications? ...did you drink much more than usual or stop drinking after you had been drinking a lot for a while? ...were you physically ill?

IF YES TO ANY: Has there been a time in the past year when you had (PSYCHOTIC SXS) and were not (USING DRUGS/TAKING MEDICATION/CHANGING YOUR DRINKING HABITS/ILL)?

DELUSIONS

Now I'd like to ask you about unusual experiences that people sometimes have.

A false belief based on incorrect inference about external reality that is firmly held despite what almost everyone else believes and despite what constitutes incontrovertible and obvious proof or evidence to the contrary. The belief is not one ordinarily accepted by other members of the person's culture or subculture. When a false belief involves a value judgment, it is regarded as a delusion only when the judgment is so extreme as to defy credibility. Code overvalued ideas (unreasonable and sustained beliefs that are maintained with less than delusional intensity) as "2."

In the past year, since (1 YEAR AGO), has it Delusion of reference, i.e., events, objects, or seemed like people were talking about you or taking special notice of you? (What do you think they were saying about you?)

other persons in the individual's immediate environment are seen as having a particular and unusual significance.


IF YES: Were you convinced they were DESCRIBE: talking about you or did you think it might have been your imagination?

Did you ever have the feeling that something on the radio, TV, or in a movie was meant especially for you? (...not just that it was particularly relevant to you, but that it was specifically meant for you.)

Did you have the feeling that the words in a popular song were meant to send you a special message? (...not just that they were particularly relevant to you, but that they were specifically meant for you.)

Did you have the feeling that what people were wearing was intended to send you a special message?

Did you have the feeling that street signs or billboards had a special meaning for you?



What about anyone going out of their way to give you a hard time, or trying to hurt you? (Tell me about that.)

Persecutory delusion, i.e., the central theme is that one (or someone to whom one is close to) is being attacked, harassed, cheated, persecuted, or conspired against.

BC2 2 3 1 BC2a 3 POSS/DEF **PRIMARY** SUBST/GMC

Have you had the feeling in the past year that you were being followed, spied on, manipulated, or plotted against?

DESCRIBE:

Did you have the feeling that you were being poisoned or that your food had been tampered with?

In the past year, since (1 YEAR AGO), Have Grandiose delusion, i.e., content involves you thought that you were especially important in some way, or that you had special powers or knowledge? (Tell me about that.)


inflated worth, power, knowledge identity, or a special relationship to a deity or famous person.

DESCRIBE:

BC3 1 2 BC3a 1 3 POSS/DEF **PRIMARY** SUBST/GMC

In the past year, since (1 YEAR AGO), did you believe that you had a special or close relationship with a celebrity or someone else famous?

In the past year, since (1 YEAR AGO), have Somatic delusion, i.e., main content pertains to **you been convinced that something was** the appearance or functioning of one's body. very wrong with your physical health even though your doctor said nothing was wrong...like you had cancer or some other DESCRIBE: disease? (Tell me about that.)



Did you feel that something strange was happening to parts of your body?


you felt that you had committed a crime or done something terrible for which you should be punished? (Tell me about that.)

Have you felt that something you did, or should have done but did not do, caused serious harm to your parents, children, other family members, or friends?

What about feeling responsible for a disaster such as a fire, flood, or earthquake?

In the past year, since (1 YEAR AGO), have Delusion of guilt, i.e., a belief that a minor error in the past will lead to disaster, or that he or she has committed a horrible crime and should be punished severely, or that he or she is responsible for a disaster (e.g., an earthquake or fire) with which there can be no possible connection

DESCRIBE:




you been convinced that your spouse or partner was being unfaithful

IF YES: How did you know they were being unfaithful? (What clued you into this?)

In the past year, since (1 YEAR AGO), have Jealous delusion, i.e., that one's sexual partner is unfaithful

DESCRIBE:



In the past year, since (1 YEAR AGO), did you have a "secret admirer" who, when you tried to contact them, denied that they were in love with you? (Tell me about that.)

Erotomanic delusion, i.e., that another person, usually of higher status, is in love with the individual.

BC7 ? 2 1 3 1 BC7a POSS/DEF **PRIMARY** SUBST/GMC

Were you romantically involved with someone famous? (Tell me about that.) DESCRIBE:

DESCRIBE:

Are you a religious or spiritual person?

Religious delusion, i.e., a delusion with a religious or spiritual content.

3 BC8 1 3 BC8a POSS/DEF **PRIMARY**

SUBST/GMC

1

1

1

POSS/DEF

?

SUBST/GMC

2

2

3

3

PRIMARY

3

▶ IF YES: In the past year, since (1 YEAR AGO), have you had any religious or spiritual experiences that the other people in your religious or spiritual community have not experienced?

> IF YES: Tell me about your experiences. (What did they think about these experiences of yours?)

IF NO: Have you felt that God, the devil, or some other spiritual being or higher power has communicated directly with you? (Tell me about that. Do others in your religious or spiritual community also have such experiences?)

►IF NO: In the past year, since (1 YEAR AGO), have you felt that God, or the devil or some other spiritual being or higher power has communicated directly with you? (Tell me about that. Do others in your religious or spiritual community also have such experiences?)

In the past year, since (1 YEAR AGO), did

you feel that someone or something

outside yourself was controlling your

thoughts or actions against your will?

(Tell me about that.

Delusion of being controlled, i.e., feelings, impulses, thoughts, or actions are experienced as being under the control of some external force rather than under one's own control.

DESCRIBE:

In the past year, since (1 YEAR AGO), did you feel that certain thoughts that were not your own were put into your head? (Tell me about that.)


Thought insertion, i.e., that certain thoughts are not one's own, but rather are inserted into one's mind.

DESCRIBE:

What about thoughts being taken out of your head? (Tell me about that.)

Thought withdrawal, i.e., that one's thoughts have been "removed" by some outside force.

DESCRIBE:




BC9

BC9A

BC10

In the past year, since (1 YEAR AGO), did you feel as if your thoughts were being broadcast out loud so that other people could actually hear what you were thinking? (Tell me about that.) **Thought broadcasting,** i.e., the delusion that one's thoughts are being broadcast out loud so that others can perceive them.

DESCRIBE:



In the past year, since (1 YEAR AGO), did you believe that someone could read your mind? (Tell me about that.)

Other delusions (e.g., that others can read the person's mind, a delusion that one has died several years ago).

DESCRIBE:



HALLUCINATIONS

A perception-like experience with the clarity and impact of a true perception, but without the external stimulation of the relevant sensory organ. The person may or may not have insight into the nonveridical nature of the hallucination (i.e., one hallucinating person may recognize the false sensory experience, whereas another may be convinced that the experience is grounded in reality).

NOTE: Code "2" for hallucinations that are so transient as to be without diagnostic significance. Code "1" for hypnagogic or hypnopompic hallucinations.

In the past year, since (1 YEAR AGO), did you hear things that other people couldn't, such as noises, or the voices of people whispering or talking? (Were you awake at the time?)


IF YES: What did you hear? How often DESCRIBE: did you hear it?

In the past year, since (1 YEAR AGO), did you have visions or see things that other people couldn't see? (Tell me about that. Were you awake at the time?)

NOTE: DISTINGUISH FROM AN ILLUSION, I.E., DESCRIBE: A MISPERCEPTION OF A REAL EXTERNAL STIMULUS.

Auditory hallucinations, i.e., involving the perception of sound, most commonly of voice) when fully awake, heard either inside or outside of one's head.

Visual hallucinations, i.e., a hallucination involving sight, which may consist of formed images, such as of people or of unformed images, such as flashes of light.



What about strange sensations on your skin, like feeling like something is creeping or crawling on or under your skin? How about the feeling of being touched or stroked? (Tell me about that.)

Tactile hallucinations, i.e., a hallucination involving the perception of being touched or of something being under one's skin.

DESCRIBE

What about having unusual sensations inside a part of your body, like a feeling of electricity? (Tell me about that.)

Somatic hallucination, i.e., a hallucination involving the perception of physical experience localized within the body (e.g., a feeling of electricity).

DESCRIBE:

How about eating or drinking something in the past year that you thought tasted bad or strange even though everyone else unpleasant) who tasted it thought it was fine? (Tell me about that.)

Gustatory hallucinations, i.e., a hallucination involving the perception of taste (usually


DESCRIBE:


What about smelling unpleasant things that other people couldn't smell, like decaying food or dead bodies? (Tell me about that.)

Olfactory hallucinations, i.e., a hallucination involving the perception of odor

DESCRIBE:

ANY ITEM CODED "3" IN "PRIMARY" SECTION





3

PRIMARY

BC19a

1


POSS/DEF

SUBST/GMC

IF A MAJOR DEPRESSIVE OR MANIC EPISODE Psychotic symptoms occur at times other than HAS EVER BEEN PRESENT: Has there ever been a time in the past year when you had (PSYCHOTIC SXS) and you were not (depressed/high/ irritable/OWN WORDS)?

during mood episodes.

NOTE: Code "3" if psychotic symptoms have been present and either: 1) there have never been any Major Depressive or Manic Episodes, or 2) psychotic symptoms occurred outside of Major Depressive or Manic Episodes. Code '1" if psychotic symptoms have occurred only during Major Depressive or Manic Episodes.



EXPLORE DETAILS AND DESCRIBE DIAGNOSTIC SIGNIFICANCE:

BC22

Etiological general medical conditions include:

Neurological conditions (e.g., neoplasms, cerebrovascular disease, Huntington's disease, multiple sclerosis, epilepsy, auditory or visual nerve injury or impairment, deafness, migraine, central nervous system infections), endocrine conditions (e.g., hyper- and hypothyroidism, hyper- and hypoparathyroidism, hyper- and hypoadrenocorticism), metabolic conditions (e.g., hypoxia, hypercarbia, hypoglycemia), fluid or electrolyte imbalances, hepatic or renal diseases, and autoimmune disorders with central nervous system involvement (e.g., systemic lupus erythematosus).

Etiological substances/medications include:

Alcohol (during intoxication or withdrawal); cannabis (during intoxication); hallucinogens (during intoxication), phencyclidine (and related substances (during intoxication); inhalants (during intoxication); sedatives, hypnotics, and anxiolytics (during intoxication or withdrawal); and stimulants (including cocaine) (during intoxication);

Other substances and medications that can cause psychotic symptoms include anesthetics and analgesics, anticholinergic agents, anticonvulsants, antihistamines, antihypertensive and cardiovascular medications, antimicrobial medications, antiparkinsonian medications, chemotherapeutic agents (e.g., cyclosporine, procarbazine), corticosteroids, gastrointestinal medications, muscle relaxants, nonsteroidal anti-inflammatory medications, other over-the-counter medications (e.g., phenylephrine, pseudoephedrine), antidepressant medication, and disulfiram. Toxins include anticholinesterase, organophosphate insecticides, sarin and other nerve gases, carbon monoxide, carbon dioxide, and volatile substances such as fuel or paint.

L. TRAUMA- AND STRESSOR-RELATED DISORDERS

TRAUMA HISTORY

I'd now like to ask about some things that may have happened to you that may have been extremely upsetting. People often find that talking about these experiences can be helpful. I'll start by asking if these experiences apply to you, and if so, I'll ask you to briefly describe what happened and how you felt at the time.

SCREEN FOR EACH TYPE OF TRAUMA USING QUESTIONS BELOW; THEN, ON PAGES L.2-L.5 REVIEW AND INQUIRE IN DETAIL FIRST FOR ANY EVENTS OCCURRING IN THE PAST MONTH AND THEN FOR UP TO THREE PAST EVENTS (E.G., THREE WORST EVENTS, THREE MOST RECENT EVENTS, ETC.)

Have you ever been in a life-threatening situation like a major disaster or fire, combat, or a serious car or work- related accident?	L1
What about being physically or sexually assaulted or abused, or threatened with physical or sexual assault?	L2
How about seeing another person being physically or sexually assaulted or abused, or threatened with physical or sexual assault?	L3
Have you ever seen another person killed or dead, or badly hurt?	L4
How about learning that one of these things happened to someone you are close to?	L5
IF UNKNOWN: Have you ever been the victim of a serious crime?	L6
IF NO EVENTS ENDORSED: What would you say has been the most stressful or traumatic experience you have had over your life?	L7
IF NO EVENTS ACKNOWLEDGED, CHECK HERE AND GO TO *ADJUSTMENT DISORDER* L.20. OTHERWISE CONTINUE ON NEXT PAGE.	L8

Did any of these happen in the past month, since (1 MONTH AGO)?

FIF YES: ASSESS THE TRAUMATIC EVENT IN PAST MONTH USING THE QUESTIONS BELOW.

► IF NO: CONTINUE ON TOP OF PAGE L.3.

DETAILS FOR EVENT IN PAST MONTH

	Description of traumatic event:	L9
→ IF DIRECT EXPOSURE TO TRAUMA:		
What happened? Were you afraid of dying or being seriously hurt? Were you seriously hurt?		
TE WITHEGGED TRAUMATIC EVENT	Indicate type of traumatic event: (check all that apply)	
→IF WITNESSED TRAUMATIC EVENT HAPPENING TO OTHERS:	Death, actual	L10
What happened? What did you see? How close were you to (TRAUMATIC	Death, threatened	L11
EVENT)? Were you concerned about your own safety?	Serious Injury, actual	L12
→ IF LEARNED ABOUT TRAUMATIC EVENT:	Serious injury, threatened	L13
What happened? Who did it involve?	Sexual violence, actual	L14
(How close [emotionally] were you to them? Did it involve violence, suicide or a bad accident?)	Sexual violence, threatened	L15
,	Indicate mode of exposure to traumatic event: (check all that apply)	
	Directly experienced	L16
	Witnessed happening to others in person	L17
	Learning about actual or threatened violence or accidental death of a close family member or friend	L18
	 Repeated or extreme exposure to aversive details of traumatic events (e.g., police officers repeatedly exposed to details of child abuse) 	; L19
IF UNKNOWN: How old were you at the time?	Age at time of event:	L20
IF UNKNOWN: Did this happen more than once?	Indicate type of exposure: (circle the appropriate number)	
	1 - Single event	L21
	Prolonged or repeated exposure to same trauma (e.g., witnessing repeated episodes of parental domestic violence over years	

→ IF NO EVENTS PRIOR TO PAST MONTH, GO TO *ADJUSTMENT DISORDER*.

→ IF EVENTS PRIOR TO PAST MONTH, REVIEW THE TYPES OF TRAUMA INDICATED ON SCREENING AND CHOOSE THE THREE MOST SEVERE EVENTS TO ASSESS, USING THE FOLLOWING QUESTIONS:

DETAILS FOR PAST EVENT #1

	Description of traumatic event:	L22
→ IF DIRECT EXPOSURE TO TRAUMA:		
What happened? Were you afraid of dying or being seriously hurt? Were you seriously hurt?		
	Indicate type of traumatic event: (check all that apply)	
F WITNESSED TRAUMATIC EVENT HAPPENING TO OTHERS:	Death, actual	L23
What happened? What did you see? How close were you to (TRAUMATIC EVENT)? Were you concerned about your own safety?	Death, threatened	L24
→ IF LEARNED ABOUT TRAUMATIC EVENT:	Serious Injury, actual	L25
What happened? Who did it involve? (How close [emotionally] were you to	Serious injury, threatened	L26
them? Did it involve violence, suicide or a bad accident?)	Sexual violence, actual	L27
	Sexual violence, threatened	L28
	Indicate mode of exposure to traumatic event: (check all that apply)	
	Directly experienced	L29
	Witnessed happening to others in person	L30
	Learning about actual or threatened violence or accidental death of a close family member or friend	L31
	Repeated or extreme exposure to aversive details of traumatic events (e.g., police officers repeatedly exposed to details of child abuse)	L32
IF UNKNOWN: How old were you at the time?	Age at time of event:	L33
IF UNKNOWN: Did this happen more than once?	Indicate type of exposure: (circle the appropriate number)	L34
onec:	1 – Single event	
	 2 - Prolonged or repeated exposure to same trauma (e.g., witnessing repeated episodes of parental domestic violence over years) 	

DETAILS FOR PAST EVENT #2

→ IF DIRECT EXPOSURE TO TRAUMA:	Description of traumatic event:	
What happened? Were you afraid of dying or being seriously hurt? Were you seriously hurt?		L35
→ IF WITNESSED TRAUMATIC EVENT HAPPENING TO OTHERS:	Indicate type of traumatic event: (check all that apply):	
What happened? What did you see? How close were you to (TRAUMATIC EVENT)? Were you concerned about	Death, actual	L36
your own safety?	Death, threatened	L37
► IF LEARNED ABOUT TRAUMATIC EVENT: What happened? Who did it involve?	Serious Injury, actual	L38 L39
(How close [emotionally] were you to them? Did it involve violence, suicide or a bad accident?)	Serious injury, threatened	
or a para accident.	Sexual violence, actual	L40
	Sexual violence, threatened	L41
	Indicate mode of exposure to traumatic event: (check all that apply)	
	Directly experienced	L42
	Witnessed happening to others in person	L43
	Learning about actual or threatened violence or accidental death of a close family member or friend	L44
	Repeated or extreme exposure to aversive details of traumatic events (e.g., police officers repeatedly exposed to details of child abuse)	L45
F UNKNOWN: How old were you at the ime?	Age at time of event:	L46
F UNKNOWN: Did this happen more than nce?	Indicate type of exposure : (circle the appropriate number)	
	1 - Single event	L47
	2 – Prolonged or repeated exposure to same trauma (e.g., witnessing repeated episodes of parental domestic violence over years	

DETAILS FOR PAST EVENT #3

Description of traumatic event:	L48
Indicate type of traumatic event: (check all that apply)	
Death, actual	L49
Death, threatened	L50
Serious Injury, actual	L51
Serious injury, threatened	L52
Sexual violence, actual	L53
Sexual violence, threatened	L54
Indicate mode of exposure to traumatic event: (check all that apply)	
Directly experienced	L55
Witnessed happening to others in person	L56
Learning about actual or threatened violence or accidental death of a close family member or friend	L57
Repeated or extreme exposure to aversive details of traumatic events (e.g., police officers repeatedly exposed to details of child abuse)	L58
Age at time of event:	L59
Indicate type of exposure: (circle the appropriate number)	
1 – Single event	L60
 2 – Prolonged or repeated exposure to same trauma (e.g., witnessing repeated episodes of parental domestic violence over years 	
	Indicate type of traumatic event: (check all that apply) Death, actual Death, threatened Serious Injury, actual Serious injury, threatened Sexual violence, actual Sexual violence, threatened Indicate mode of exposure to traumatic event: (check all that apply) Directly experienced Witnessed happening to others in person Learning about actual or threatened violence or accidental death of a close family member or friend Repeated or extreme exposure to aversive details of traumatic events (e.g., police officers repeatedly exposed to details of child abuse) Age at time of event: Indicate type of exposure: (circle the appropriate number) 1 - Single event 2 - Prolonged or repeated exposure to same trauma (e.g., witnessing repeated episodes of parental domestic

POSTTRAUMATIC STRESS DISORDER **POSTTRAUMATIC STRESS DISORDER CRITERIA**

FOR FOLLOWING QUESTIONS, FOCUS ON THE THREE MOST SEVERE TRAUMATIC EVENT(S) DESCRIBED ON PAGES L.3-L.5.

L87 IF ALL TRAUMAS ARE CONFINED TO THE PAST MONTH, CHECK HERE __ AND SKIP TO *ADJUSTMENT DISORDER*. A. Exposure to actual or threatened death, serious injury, or sexual violence in one (or more) of the following ways: 3 L88 IF MORE THAN ONE TRAUMATIC EVENT IS 1. Directly experiencing the traumatic event(s). ? 1 REPORTED: Which of these do you think affected you the most? 2. Witnessing, in person, the event(s) as it ? 1 2 3 L89 occurred to others. IF SELECTED EVENT IS ULTIMATELY NOT ASSOCIATED WITH THE FULL PTSD SYNDROME, 2 3 L90 1 3. Learning that the traumatic event(s) occurred CONSIDER RE-ASSESSING THE ENTIRE PTSD to a close family member or close friend. In CRITERIA SET (PAGES L.11-L.17) FOR OTHER cases of actual or threatened death of a family REPORTED TRÀUMAS. member or friend, the event(s) must have been violent or accidental. 4. Experiencing repeated or extreme exposure to 3 L91 1 2 aversive details of the traumatic event(s) (e.g., first responders collecting human remains; police officers repeatedly exposed to details of child abuse). Note: Criterion A4 does not apply to exposure through electronic media, television, movies, or pictures, unless the exposure is work-related. 3 L92 AT LEAST ONE A ITEM CODED "3" GO TO *ADJUSTMENT **DISORDER*** Now I'd like to ask a few questions about specific B. Presence of one (or more) of the following ways that (TRAUMATIC EVENT) may have affected intrusion symptoms associated with the you in the past year, since (1 YEAR AGO). traumatic events), beginning after the traumatic event(s) occurred: 1. Recurrent, involuntary, and intrusive 2 3 L93 For example, in the past year... 1 distressing memories of the traumatic event(s). ...have you had memories of (TRAUMATIC EVENT), including feelings, physical sensations, sounds, smells, or images, when you didn't expect to or want to? (How often has this happened?) Past month L94 IF PAST YEAR RATING OF "3": Has this also 1 2 3 happened in the past month, since (1 MONTH AGO)? How many times? (IN THE PAST YEAR...) what about having 2. Recurrent distressing dreams in which the 2 3 L95 1 upsetting dreams that reminded you of content and/or affect of the dream are related (TRAUMATIC EVENT)? Tell me about that. to the traumatic event. IF PAST YEAR RATING OF "3": Has this also Past month happened in the past month? How many L96 3 1 2 times?

SCID-RV (for DSM-5®) (Version 1.0.0)

(IN THE PAST YEAR...)

...what about having found yourself acting or feeling as if you were back in the situation? (Have you had "flashbacks' of [TRAUMATIC EVENT]?)

IF PAST YEAR RATING OF "3": Has this also happened in the past month? How many times?

(IN THE PAST YEAR...) have you had a strong emotional or physical reaction when something reminded you of (TRAUMATIC EVENT)? Give me some examples of the kinds of things that would have triggered this reaction. (Things like...seeing a person who resembles the person who attacked you, hearing the screech of brakes if you were in a car accident, hearing the sound of helicopters if you were in combat, any kind of physically intimacy in someone who was raped?)

NOTE: IF DENIES EMOTIONAL OR PHYSICAL REACTION TO REMINDERS, CODE "1" FOR BOTH B.4 (EMOTIONAL REACTION) AND B.5 (PHYSICAL REACTION).

IF YES: What kind of reaction did you have? Did you get very upset or stay upset for a while, even after the reminder had gone away?


IF PAST YEAR RATING OF "3": Has this also happened in the past month, since (1 MONTH AGO)? How many times?

IF ACKNOWLEDGES STRONG EMOTIONAL OR PHYSICAL REACTION: What about having physical symptoms—like breaking out in a sweat, breathing heavily or irregularly, or feeling your heart pound or race when something reminded you of (TRAUMATIC EVENT)? How about feeling tense or shaky?


IF PAST YEAR RATING OF "3": Has this also happened in the past month? How many times?

PTSD Trauma and Stressor-Related Disorders L.57

3. Dissociative reactions (e.g., flashbacks) in which the individual feels or acts as if the traumatic event(s) were recurring. (Such reactions may occur on a continuum, with the most extreme expression being a complete loss of awareness of present surroundings.)




 Intense or prolonged psychological distress at exposure to internal or external cues that symbolize or resemble an aspect of the traumatic event(s).




L100

Marked physiological reactions to internal or external cues that symbolize or resemble an aspect of the traumatic event(s).



AT LEAST ONE "B" SX IS CODED "3."



In the past year...

C. Persistent avoidance of stimuli associated with the traumatic event(s), beginning after the traumatic event(s) occurred, as evidenced by one or both of the following:

...have you done things to avoid remembering or thinking about (TRAUMATIC EVENT) like keeping yourself busy, distracting yourself like by playing computer or video games or watching TV, or using drugs or alcohol to "numb" yourself or try to forget what happened? (Since [TRAUMATIC EVENT], how long has this gone on?)

IF NO: How about doing things to avoid having feelings similar to those you had during (TRAUMATIC EVENT)? (Since [TRAUMATIC EVENT], how long has this gone on?)


IF PAST YEAR RATING OF "3": Has this also happened in the past month, since (1 MONTH AGO)? How many times?

(IN THE PAST YEAR) ...have there been things, places, or people that you have tried to avoid because it brought up upsetting memories, thoughts, or feelings about (TRAUMATIC EVENT)? (Since [TRAUMATIC EVENT], how long has this gone on?)

IF NO: How about avoiding certain activities, situations, or topics of conversation? (Since [TRAUMATIC EVENT], how long has this gone on?)

IF PAST YEAR RATING OF "3": Has this also happened in the past month? How many times?

 Avoidance of, or efforts to avoid distressing memories, thoughts, or feelings about or closely associated with the traumatic event(s).




2


1

3 L105

 Avoidance of or efforts to avoid external reminders (people, places, conversations, activities, objects, situations), that arouse distressing memories, thoughts, or feelings about or closely associated with the traumatic event(s).



AT LEAST ONE "C" SX IS CODED "3."



In the past year...

D. Negative alterations in cognitions and mood associated with the traumatic event(s), beginning or worsening after the traumatic event(s) occurred, as evidenced by two (or more) of the following:

...have you been unable to remember some important part of what happened? (Tell me about that.)

IF YES: Did you get a head injury during (TRAUMATIC EVENT)? Were you drinking a lot or were taking any drugs at the time of (TRAUMATIC EVENT)?

IF PAST YEAR RATING OF "3": Has this also happened in the past month, since (1 MONTH AGO)? How many times?

(IN THE PAST YEAR) ...has there been a change in how you think about yourself? (Like feeling you are "bad," or permanently damaged or "broken?" Tell me about that. Since this started, have you felt this way most of the time?)

IF NO: Has there been a change in how you see other people or the way the world works? (Like you can't trust anyone anymore? Like the world is a completely dangerous place? Tell me about that. Since this started, have you felt this way most of the time?)

IF PAST YEAR RATING OF "3": Has this also happened in the past month? How much of the time?

(IN THE PAST YEAR) ...have you blamed yourself for the (TRAUMATIC EVENT) or how it affected your life? (Like feeling that (TRAUMATIC EVENT) was your fault or that you should have done something to prevent it? Like feeling that you should have gotten over it by_now?)

IF YES: Tell me about that. (Since this started, have you felt this way most of the time?)

IF NO: Have you blamed someone else for (TRAUMATIC EVENT)? Tell me about that. (What did they have to do with [TRAUMATIC EVENT]?)

IF PAST YEAR RATING OF "3": Has this also happened in the past month? How much of the time?

(IN THE PAST YEAR) ...have you had bad feelings much of the time, like feeing sad, angry, afraid, guilty, ashamed, "in shock"? (Tell me about that.)

IF YES: Is this different from the way you
were before (TRAUMATIC EVENT)?

IF PAST YEAR RATING OF "3": Has this also happened in the past month? How many times?


 Inability to remember an important aspect of the traumatic event(s) (typically due to dissociative amnesia and not to other factors such as head injury, alcohol, or drugs).


 Persistent and exaggerated negative beliefs or expectations about oneself, others, or the world (e.g., "I am bad," "No one can be trusted," "The world is completely dangerous," "My whole nervous system is permanently ruined").


cause or consequences of the traumatic event(s) that lead the individual to blame himself/herself or others.


3. Persistent, distorted cognitions about the

Persistent negative emotional state (e.g., fear, horror, anger, guilt, or shame).









(IN THE PAST YEAR...)

...have you been less interested in things that you were interested in before (TRAUMATIC EVENT), like spending time with family or friends, reading books, watching TV, cooking, or sports? (Tell me about that.)

IF NO LOSS OF INTEREST: Are you still doing as many activities as you used to?

IF PAST YEAR RATING OF "3": Has this also happened in the past month? How many times?

(IN THE PAST YEAR) ...have you felt distant or disconnected from others or have you closed yourself off from other people? (Tell me about that.)


IF PAST YEAR RATING OF "3": Has this also happened in the past month, since (1 MONTH AGO)? How often?

(IN THE PAST YEAR) ...have you been unable to experience good feelings, like feeling happy, joyful, satisfied, loving, or tender towards other people? (Tell me about that.)

IF YES: Is this different from the way you
were before (TRAUMATIC EVENT)?


IF PAST YEAR RATING OF "3": Has this also happened in the past month? How often?

5. Markedly diminished interest or participation in significant activities.




Feelings of detachment or estrangement from others.

 Persistent inability to experience positive emotions (e.g., inability to experience happiness, satisfaction, or loving feelings).



AT LEAST THREE "D" SXS ARE CODED "3."



Past month

1 2 3

L124

In the past year...

E. Marked alterations in arousal and reactivity associated with the traumatic event(s), beginning or worsening after the traumatic event(s) occurred, as evidenced by two (or more) of the following:


...have you lost control of your anger, so that you threatened or hurt someone or damaged something? Tell me what happened. (Was it over something little or even nothing at all?)

IF NO: Since (TRAUMATIC EVENT), have you been more quick-tempered or had a shorter "fuse" than before?

IF YES TO EITHER: How different is this from the way you were before (TRAUMATIC EVENT)?

IF PAST YEAR RATING OF "3": Has this also happened in the past month, since (1 MONTH AGO)? How often?

 Irritable behavior and angry outbursts (with little or no provocation) typically expressed as verbal or physical aggression toward people or objects.



(IN THE PAST YEAR...)

...have you done reckless things, like driving dangerously, or drinking or using drugs without caring about the consequences?

IF NO: How about hurting yourself on purpose or trying to kill yourself? (What did vou do?)

IF YES TO EITHER: How different is this from the way you were before (TRAUMATIC EVENT)?

IF PAST YEAR RATING OF "3": Has this also happened in the past month? How often?

2. Reckless or self-destructive behavior.

NOTE: Any current suicidal thoughts, plans, or actions should be thoroughly assessed by the clinician and action taken if necessary.

> Past month L130 1 2 3

2

3 L129

?

1

(IN THE PAST YEAR) ... have you noticed that you have been more watchful or on guard? (What are some examples?)

IF NO: Have you been extra aware of your surroundings and your environment?

IF PAST YEAR RATING OF "3": Has this also happened in the past month, since (1 MONTH AGO)? How often?

(IN THE PAST YEAR) ... have you been jumpy or easily startled, like by sudden noises? (Is this a change from before [TRAUMATIC EVENT]?)

IF PAST YEAR RATING OF "3": Has this also happened in the past month? How often?

(IN THE PAST YEAR) ...have you had trouble concentrating? (What are some examples? (Is this a change from before [TRAUMATIC EVENT]?)

IF PAST YEAR RATING OF "3": Has this also happened in the past month? How often?

(IN THE PAST YEAR) ...how have you been sleeping since (TRAUMATIC EVENT)? (Is this a change from before [TRAUMATIC EVENT]?)

IF PAST YEAR RATING OF "3": Has this also happened in the past month? How often? 3. Hypervigilance.

4. Exaggerated startle response.

5. Problems with concentration.

6. Sleep disturbances (e.g., difficulty falling or staying asleep or restless sleep).

AT LEAST TWO "E" SXS ARE CODED "3."

3 L131 1 2 Past month 1 2 3 L132

Past month L134 1 2 3 2 1 L135

2

1

3 L133

L136

1 2 3 L137 Past month 3 L138 1 2

Past month

1 2 3

3 L139 GO TO *ADJUSTMENT **DISORDER*** CRITERION E MET L140 PAST MONTH 1 3

SCID-RV (for DSM-5®) (Version 1.0.0)

About how long did these (PTSD SYMPTOMS CODED "3") **last altogether?**

PTSD Trauma and Stressor-Related Disorders L.62

F. Duration of the disturbance (symptoms in criteria B, C, D, and E) is more than 1 month.

	?	1 	2	3 _{L141}
*		ISTME RDER*	NT	

IF UNKNOWN: What effect did (PTSD SXS) have on your life in the past year?

ASK THE FOLLOWING QUESTIONS <u>AS NEEDED</u> TO RATE CRITERION G:

How have (PTSD SXS) affected your relationships or your interactions with other people? (Have they caused you any problems in your relationships with your family, romantic partner or friends?)

How have (PTSD SXS) affected your work/school? (How about your attendance at work or school? Have they affected the quality of your work/schoolwork?)

How have they affected your ability to take care of things at home? What about being involved in things that are important to you like religious activities, physical exercise, or hobbies?


Have (PTSD SXS) affected any other important part of your life?

IF HAVE NOT INTERFERED WITH LIFE IN PAST YEAR: How much have you been bothered or upset by (PTSD SXS)?

IF PAST YEAR RATING OF "3": How have (PTSD SXS) affected your life in the past month, since (1 MONTH AGO)?

G. The disturbance causes clinically significant distress or impairment in social, occupational, or other important areas of functioning.

CRITERION H HAS BEEN OMITTED.



CRITERION G MET PAST MONTH
? 1 2 3

L143

ADJUSTMENT DISORDER

CONSIDER THIS SECTION ONLY IF THERE ARE SYMPTOMS OCCURRING IN THE PAST YEAR THAT DO NOT MEET THE CRITERIA FOR ANOTHER DSM-5 DISORDER. OTHERWISE, CHECK HERE ___ AND GO TO THE NEXT MODULE. INFORMATION OBTAINED L151 FROM OVERVIEW OF PRESENT ILLNESS WILL USUALLY BE SUFFICIENT TO RATE THE CRITERIA FOR ADJUSTMENT DISORDER.

ADJUSTMENT DISORDER CRITERIA

IF UNKNOWN: Did anything happen to you before (SYMPTOMS) began?

IF YES: Tell me about what happened. Do you think that (STRESSOR) had anything to do with your developing (SXS)?

► IF SINGLE EVENT: How long after (STRESSOR) did you first develop (SXS)? (Was it within 3 months?)

▶ IF CHRONIC STRESSOR: How long after DESCRIBE STRESSOR: (STRESSOR) began did you first develop (SXS)? (Was it within 3 months?)

A. The development of emotional or behavioral symptoms in response to an identifiable stressor(s) occurring within 3 months of the onset of the stressor(s).

DESCRIBE SYMPTOMS:

3 L152 1 GO TO THE **NEXT MODULE**

IF UNKNOWN: What effect did (SXS) have on your life in the past year?

ASK THE FOLLOWING QUESTIONS AS NEEDED TO RATE CRITERION B:

How have (SXS) affected your relationships or your interactions with other people? (Have they caused you any problems in your relationships with your family, romantic partner or friends?)


How have (SXS) affected your work/school? (How about your attendance at work or school? Did [SXS] make it more difficult to do your work/schoolwork? How did [SXS] affect the quality of your work/schoolwork?)

How have they affected your ability to take care of things at home? What about being involved in things that are important to you like religious activities, physical exercise, or hobbies?

Have (SXS) affected any other important part of your life?

IF HAVE NOT INTERFERED WITH LIFE: How much have you been bothered or upset by having (SXS)? How upset are you about (STRESSOR)? (Are you more upset than most other people would be? Have others said that you're more upset than you should be? Have [SXS] lasted longer than you or other people think they should have?)


- B. These symptoms or behaviors are clinically significant as evidenced by one or both of the following:
 - 1. Marked distress that is out of proportion to the severity and intensity of the stressor, taking into account the external context and the cultural factors that might influence symptom severity and presentation.
 - 2. Significant impairment in social, occupational, or other important areas of functioning.



Have you had this kind of reaction many times before?


IF UNKNOWN: Were you having these (SXS) even before (STRESSOR) happened?

C. The stress-related disturbance does not meet the criteria for another mental disorder and is not merely an exacerbation of a preexisting mental [including personality] disorder.




IF UNKNOWN: Did someone close to you die just before (SXS)?


D. The symptoms do not represent normal bereavement.



IF UNKNOWN: How long has it been since (STRESSOR AND ITS CONSEQUENCES) was over?

E. Once the stressor (or its consequences) has terminated, the symptoms do not persist for more than an additional 6 months.





Indicate type based on predominant symptoms: (circle the appropriate number) L158

- 1 **With depressed mood:** Low mood, tearfulness, or feelings of hopelessness are predominant.
- 2 **With anxiety:** Nervousness, worry, jitteriness, or separation anxiety is predominant.
- 3 **With mixed anxiety and depressed mood:** A combination of depression and anxiety is predominant.
- 4 With disturbance of conduct: Disturbance in conduct is predominant.
- 5 **With mixed disturbance of emotions and conduct:** Both emotional symptoms (e.g., depression, anxiety) and a disturbance of conduct are predominant.
- 6 **Unspecified:** For maladaptive reactions that are not classifiable as one of the specific subtypes of adjustment disorder (e.g., physical complaints, social withdrawal, or work or academic inhibition).

GO TO THE NEXT MODULE

F. ANXIETY DISORDERS

PANIC DISORDER

PANIC DISORDER CRITERIA

▶IF SCREENING QUESTION #1 ANSWERED SCREEN Q#1 F1 "NO," SKIP TO *AGORAPHOBIA* F.8. YES NO **▶**IF QUESTION #1 ANSWERED "YES": You've said that you have had an intense rush of anxiety, or what GO TO someone might call a "panic attack," in ITEM the past year when you <u>suddenly</u> felt F23_L150 very frightened, or anxious or suddenly developed a lot of physical symptoms. ▶IF SCREENER NOT USED: In the past year, since (1 YEAR AGO), have you had an intense rush of anxiety, or what someone might call a "panic attack," when you suddenly felt very frightened, or anxious or suddenly developed a lot of physical symptoms? Tell me about that. When was the last bad one? What was it like? How did it begin? F2 IF UNKNOWN: Did the symptoms come on A panic attack is an abrupt surge of intense fear or 3 suddenly? intense discomfort that reaches a peak within minutes. IF YES: How long did it take from when it began to when it got really bad? (Did it happen within a few minutes?) **Note:** The abrupt surge can occur from a calm state or an anxious state. During that attack... 1. Palpitations, pounding heart, or accelerated ...did your heart race, pound or skip? 2 3 F3 heart rate. 2. Sweating. 1 2 3 ...did you sweat? ...did you tremble or shake? 3. Trembling or shaking. 2 3 F5 ...were you short of breath? (Have trouble 4. Sensations of shortness of breath or 2 3 F6 catching your breath? Feel like you were smothering. being smothered?) 3 F7 ...did you feel as if you were choking? 5. Feelings of choking. 2 ...did you have chest pain or pressure? 6. Chest pain or discomfort. 1 2 3 F8 7. Nausea or abdominal distress. 2 F9 ...did you have nausea or upset stomach 1 3 or the feeling that you were going to have diarrhea? F10 ...did you feel dizzy, unsteady, or like you 2 3 8. Feeling dizzy, unsteady, lightheaded or faint. 1 might faint?

SCID-RV (for DSM- $5^{ ext{@}}$) (Version 1.0.0) P	anic Disorder	Anxiety Di	sorder	s F			
did you have flushes, hot flashes, or chills?	9. Chills or heat sensations.		?	1	2	3	F11
During that attack							
did you have tingling or numbness in parts of your body?	10. Paresthesias (numbness o sensations)	r tingling	?	1	2	3	F12
did you have the feeling that you were detached from your body or mind, that time was moving slowly, or that you were an outside observer of your own thoughts or movements?	11. Derealization (feelings of undepersonalization (being doneself).		?	1	2	3	F13
IF NO: How about feeling that everything around you was unreal or that you were in a dream?							
were you afraid you were going crazy or might lose control?	12. Fear of losing control or "g	oing crazy."	?	1	2	3	F14
were you afraid that you were dying?	13. Fear of dying.		?	1	2	3	F15
	AT LEAST FOUR ITEMS CODED "THEIR PEAK WITHIN MINUTES	3" AND REACHED		1		3	F16
					PAN	IIC	
Besides the one you just described, have you had any other attacks in the past year					ATT CON	ACK NTINI H NE	UE
which had even more of the symptoms that I							

IF YES, GO BACK TO PAGE 1 AND ASSESS THE SYMPTOMS OF THAT ATTACK.

just asked you about?

In the past year, since (1 YEAR AGO), have any of these attacks come on out of the blue-in situations where you didn't expect to be nervous or uncomfortable?

F YES: What was going on when the attack(s) happened? (What were you doing at the time? Were you already nervous or anxious at the time or rather were you relatively calm or relaxed?)

► IF NO: How about the very first one you had in the past year. What were you doing at the time? (Were you already nervous or anxious at the time or rather were you relatively calm or relaxed?)

IF ATTACK IS UNEXPECTED: How many of these kinds of attacks have you had? (At least two?)

After any of these attacks...


...were you concerned or worried that you might have another attack or worried that you would feel like you were having a heart attack again, or worried that you would lose control or go crazy?

IF YES: How long did that concern or worry last? (Did it last at least a month? Nearly every day?)


...did you do anything differently because of the attacks (like avoiding certain places or not going out alone)? (What about avoiding certain activities like exercise? What about things like always making sure you're near a bathroom or exit?)

IF YES: How long did that last? (As long as a month?)

A. Recurrent unexpected panic attacks.



- B. At least one of the attacks has been followed by 1 month (or more) of one or both of the following:
 - 1. Persistent concern or worry about additional attacks or their consequences (e.g., losing control, having a heart attack, "going crazy").
 - 2. A significant maladaptive change in behavior related to the attacks (e.g., behaviors designed to avoid having panic attacks, such as avoidance of exercise or unfamiliar situations).




2

1

3 F18

CRITERION B.1 OR B.2 CODED "3"



IF UNKNOWN: When did your panic attacks C. [Primary Anxiety Disorder:] The start?

C. [Primary Anxiety Disorder:] The disturbance is not attributable to

Just before you began having panic attacks, were you taking any drugs, caffeine, diet pills, or other medicines?

(How much coffee, tea, or caffeinated beverages do you drink a day?)

Just before the attacks, were you physically ill?

IF YES: What did the doctor say?

C. [Primary Anxiety Disorder:] The disturbance is not attributable to the physiological effects of a substance (e.g., a drug of abuse, a medication) or another medical condition (e.g. hyperthyroidism, cardiopulmonary disorders).

IF THERE IS ANY INDICATION THAT PANIC ATTACKS MAY BE SECONDARY (I.E., A DIRECT PHYSIOLOGICAL CONSEQUENCE OF A GMC OR SUBSTANCE/MEDICATION), GO TO *GMC/SUBSTANCE* F.33, AND RETURN HERE TO MAKE A RATING OF "1" OR "3."

Etiological medical conditions include: endocrine disease (e.g., hyperthyroidism, pheochromocytoma, hypoglycemia, hyperadrenocortisolism), cardiovascular disorders (e.g., congestive heart failure, pulmonary embolism, arrhythmia such as atrial fibrillation), respiratory illness (e.g., chronic obstructive pulmonary disease, asthma, pneumonia), metabolic disturbances (e.g., vitamin B₁₂ deficiency, porphyria), and neurological illness (e.g., neoplasms, vestibular dysfunction, encephalitis, seizure disorders).

Etiological substances/medications include: alcohol (I/W), caffeine (I), cannabis (I), opioids (W), phencyclidine (I), other hallucinogens (I), inhalants, and stimulants (including cocaine) (I/W), sedatives, hypnotics, and anxiolytics (W); anesthetics and analgesics, sympathomimetics or other bronchodilators, anticholinergics, insulin, thyroid preparations, oral contraceptives, antihistamines, antiparkinsonian medications, corticosteroids, antihypertensive and cardiovascular medications, anticonvulsants, lithium carbonate, antipsychotic medications, antidepressant medications, and exposure to heavy metals and toxins such as organophosphate insecticide, nerve gases, carbon monoxide, carbon dioxide, volatile substances such as gasoline and paint.

D. The disturbance is not better explained by another mental disorder (e.g., the panic attacks do not occur only in response to feared social situations, as in Social Anxiety Disorder; in response to circumscribed phobic objects or situations, as in Specific Phobia; in response to obsessions, as in Obsessive-Compulsive Disorder; in response to reminders of traumatic events, as in Posttraumatic Stress Disorder; or in response to separation from attachment figures, as in

Separation Anxiety Disorder).

PRIMARY
ANXIETY
DISORDER

ALL DUE TO
SUBSTANCE
USE OR GMC
GO TO ITEM
F23_L150

ITEM


CONTINUE

WITH NEXT

F22

IF NECESSARY, RETURN TO THIS ITEM AFTER COMPLETING MODULES FOR OC AND RELATED DISORDERS AND TRAUMA- AND STRESS-RELATED DISORDERS.

A, B, C, AND D ARE CODED "3."



GO TO *AGORAPHOBIA*

AGORAPHOBIA

AGORAPHOBIA CRITERIA

- → IF SCREENING QUESTION #2 ANSWERED "NO," SKIP TO *SOCIAL ANXIETY DISORDER* F.14
- → IF QUESTION #2 ANSWERED "YES": You've said that in the past year you have been very anxious or afraid of situations like going out of the house alone, being in crowds, going to stores, standing in lines, or traveling on buses or trains.
- → IF SCREENER NOT USED: In the past year, since (1 YEAR AGO), have you been very anxious about or afraid of situations like going out of the house alone, being in crowds, going to stores, standing in lines, or traveling on buses or trains?

Tell me about the situations that you've been afraid of.

IF UNKNOWN: Have you been afraid of, or anxious about, travelling in taxi cabs, buses, trains, ships or planes?

IF UNKNOWN: How about being in open spaces, like parking lots, outdoor marketplaces, or bridges?


IF UNKNOWN: How about being in enclosed places like stores, movie theaters, or shopping malls?

IF UNKNOWN: How about standing in a line or being in a crowd?

IF UNKNOWN: How about being outside of the house alone?

- A. Marked fear or anxiety about two (or more) of the following five situations:
 - Using public transportation (e.g., [taxi cabs], buses, trains, ships, planes).
 - 2. Being in open spaces (e.g., parking lots, marketplaces, bridges).
 - 3. Being in enclosed places (e.g., shops, theaters, cinemas).
 - 4. Standing in line or being in a crowd.
 - 5. Being outside of the home alone.

AT LEAST TWO ITEMS ARE CODED "3"



1

1

1

1

1

1

GO TO *SOCIAL
ANXIETY DISORDER*

3 F44

3 F45

3 F46

3 F47

3 F48

F49

2

3

In the past year, since (1 YEAR AGO), why did you avoid (SITUATIONS CODED "3") (What were you afraid would happen?)

(Were you afraid that it might be hard for you to get out of the situation if you absolutely needed to...like if you suddenly developed a panic attack?)

(Or developing something else that would be embarrassing like losing control of your bladder or bowels or vomiting?)

(Or becoming impaired in some way like by falling or passing out?)

(How about being worried that there would be nobody there to help you in case these kinds of things happened?)

In the past year, since (1 YEAR AGO), did you almost always fell frightened or anxious when you were in (SITUATIONS CODED "3" ABOVE)?

B. The individual fears or avoids these situations because of thoughts that escape might be difficult or help might not be available in the event of developing panic-like symptoms or other incapacitating or embarrassing symptoms (e.g., fear of falling in the elderly, fear of incontinence).



 The agoraphobic situations almost always provoke fear or anxiety. ? 1 2 3 F51

GO TO
*SOCIAL
ANXIETY
DISORDER*

Have you gone out of your way to avoid these situations in the past year?

IF NO: Have you been only able to go into one of these situations if you were with someone you knew?

IF NO: When you have had to be in one of these situations, have you felt intensely afraid or anxious?

IF UNKNOWN: Have you felt any danger or threat to your safety when you were in (SITUATIONS CODED "3" ABOVE)? (Tell me about that.)

D. The agoraphobic situations are actively avoided, require the presence of a companion, or are endured with intense fear or anxiety.

GO TO
*SOCIAL
ANXIETY
DISORDER*

1 2 3 F53

2

3 F52

E. The fear or anxiety is out of proportion to the actual danger posed by the agoraphobic situations and the sociocultural context.

NOTE: Code "3" if situations do not pose danger or if fear or anxiety is out of proportion to actual danger or sociocultural context.

How long have you been afraid of or avoided (SITUATIONS CODED "3")? (At least 6 months?)


F. The fear, anxiety, or avoidance is persistent, typically lasting for 6 months or more.

G. The fear, anxiety, or avoidance causes clinically

significant distress or impairment in social,

occupational, or other important areas of

functioning.



IF UNKNOWN: What effect have (AGORAPHOBIC SXS) had on your life in the past year?

ASK THE FOLLOWING QUESTIONS <u>AS NEEDED</u> TO RATE CRITERION G:

How have (AGORAPHOBIC SXS) affected your relationships or your interactions with other people? (Have they caused any problems in your relationships with your family, romantic partner or friends?)


How have (AGORAPHOBIC SXS) affected your ability to work, take care of your family or household needs, or be involved in things that are important to you like religious activities, physical exercise, or hobbies?

Have (AGORAPHOBIC SXS) affected any other important part of your life?

IF HAVE NOT INTERFERED WITH FUNCTIONING: How much have you been bothered or upset by having (AGORAPHOBIC SXS)?


IF A GENERAL MEDICAL CONDITION
CHARACTERIZED BY INCAPACITATING
SYMPTOMS IS PRESENT: Is your avoidance of
(SITUATION) related to your (MEDICAL
CONDITION)? (Tell me about it. How often
has [INCAPACITATING SYMPTOM] actually
happened in [AVOIDED SITUATION]?)

H. If another medical condition (e.g., inflammatory bowel disease, Parkinson's disease) is present, the fear, anxiety, or avoidance is clearly excessive.




IF NECESSARY, RETURN TO THIS ITEM AFTER COMPLETING MODULES FOR OC AND RELATED DISORDERS.

I. The fear, anxiety, or avoidance is not better explained by the symptoms of another mental disorder—for example, the symptoms are not confined to Specific Phobia, situational type; do not involve only social situations (as in Social Anxiety Disorder); and are not related exclusively to obsessions (as in Obsessive-Compulsive Disorder), perceived defects or flaws in physical appearance (as in Body Dysmorphic Disorder), reminders of traumatic events (as in Posttraumatic Stress Disorder), or fear of separation (as in Separation Anxiety Disorder).



NOTE: Consider a diagnosis of Specific Phobia if fear is limited to one or only a few specific situations, or a diagnosis of Social Anxiety Disorder if fear is limited to social situations.

AGORAPHOBIA CRITERIA A, B, C, D, E, F, G, H, AND I ARE CODED "3."



SOCIAL ANXIETY DISORDER

SOCIAL ANXIETY DISORDER CRITERIA

► IF SCREENING QUESTIONS #3 AND #4 ARE BOTH ANSWERED "NO," SKIP TO *SPECIFIC PHOBIA* F.19.

IF QUESTION #3 ANSWERED "YES":

You've said that you have been especially anxious or afraid in social situations in the past year, like having a conversation or meeting unfamiliar people.

IF OUESTION #4 ANSWERED "YES":

You've [also] said that there are things that you have been afraid or felt very uncomfortable doing in front of other people in the past year, like speaking, eating, writing, or using a public bathroom.

IF SCREENER NOT USED: In the past year, since (1 YEAR AGO), have you been especially nervous or anxious in social situations like having a conversation or meeting unfamiliar people?

IF NO: Is there anything that you have been afraid to do or felt very uncomfortable doing in front of other people, like speaking, eating, writing, or using a public bathroom?

IF YES TO ANY OF ABOVE: Tell me about that. Give me some examples of when this has happened. (Situations like having a conversation, meeting people you don't know, being observed eating, drinking or going to the bathroom or performing in front of others?)

In the past year, what were you afraid would happen when you were in (SOCIAL OR PERFORMANCE SITUATION)? (Were you afraid of being embarrassed because of what you might say or how you might act? Were you afraid that this would lead to your being rejected by other people? How about making others uncomfortable or offending them because of what you said or how you acted?)


A. Marked fear or anxiety about one or more social situations in which the person is exposed to possible scrutiny by others. Examples include social interactions (e.g., having a conversation,


meeting unfamiliar people), being observed (e.g.,


eating or drinking), and performing in front of others (e.g., giving a speech).

NOTE: Code ${\rm ``1''}$ if fear or anxiety is limited to public speaking and is within normal limits.

B. The individual fears that he or she will act in a way or show anxiety symptoms that will be negatively evaluated (i.e., will be humiliating or embarrassing; will lead to rejection or offend others).









In the past year, did you almost always feel frightened when you would be in (FEARED SOCIAL OR PERFORMANCE SITUATIONS)?

C. The social situations almost always provoke fear or anxiety.



avoid (FEARED SOCIAL OR PERFORMANCE SITUATIONS) in the past year?

IF UNKNOWN: Did you go out of your way to D. The social situations are avoided or endured with intense fear or anxiety.



2

3 F72

3 F73

IF NO: How hard was it for you to be in (FEARED SOCIAL SITUATION)?

IF UNKNOWN: What would you say would be the likely outcome of (PERFORMING POORLY IN SOCIAL SITUATIONS)? (Were these situations actually dangerous in some way, like avoiding being bullied or tormented by someone?)

E. The fear or anxiety is out of proportion to the actual threat posed by the social situation and to the sociocultural context.


NOTE: Code "3" if no threat posed by social situation or if out of proportion to actual threat or sociocultural context.

GO TO *SPECIFIC **PHOBIA***

1

IF UNCLEAR: How long have (SXS OF SOCIAL ANXIETY DISORDER) lasted? (Have they lasted for at least 6 months or more?)

F. The fear, anxiety, or avoidance is persistent, typically lasting for 6 months or more.



IF UNKNOWN: What effect have (SOCIAL ANXIETY SXS) had on your life in the past year?

ASK THE FOLLOWING QUESTIONS AS NEEDED TO RATE CRITERION G:

How have (SOCIAL ANXIETY SXS) affected your ability to have friends or meet new people? (How about dating?) How have (SOCIAL ANXIETY SXS) affected your interactions with other people, especially unfamiliar people?

How have (SOCIAL ANXIETY SXS) affected your ability to do things at school or at work that require interacting with other people? (How about making presentations or giving talks?)

Have you avoided going to school or to work if you think you will be put in a situation which makes your uncomfortable?

How have (SOCIAL ANXIETY SXS) affected your ability to work, take care of your family or household needs, or be involved in things that are important to you like religious activities, physical exercise, or hobbies?

Have (SOCIAL ANXIETY SXS) affected any other important part of your life?

IF HAVE NOT INTERFERED WITH FUNCTIONING: How much you been bothered or upset by having (SOCIAL ANXIETY SXS)?

IF UNKNOWN: When did you begin having (SOCIAL ANXIETY SXS)?


Just before you began having (SOCIAL ANXIETY SXS), were you taking any drugs, caffeine, diet pills, or other medicines?

(How much coffee, tea, or caffeinated beverages did you drink a day?)

Just before (SOCIAL ANXIETY SXS) began, were you physically ill?

IF YES: What did the doctor say?


G. The fear, anxiety, or avoidance causes clinically significant distress or impairment in social, occupational, or other important areas of functioning.



H. [Primary Anxiety Disorder:] The fear, anxiety, or avoidance is not attributable to the physiological effects of a substance (e.g., a drug of abuse, a medication) or another medical condition.

IF THERE IS ANY INDICATION THAT THE ANXIETY MAY BE SECONDARY (I.E., A DIRECT PHYSIOLOGICAL CONSEQUENCE OF GMC OR SUBSTANCE), GO TO *GMC/SUBSTANCE* F.33, AND RETURN HERE TO MAKE A RATING OF "1" OR "3."

NOTE: Refer to list of etiological medical conditions or substances/medications on page F.4.




IF NECESSARY, RETURN TO THIS ITEM AFTER COMPLETING MODULES FOR OC AND RELATED DISORDERS.

IF A GENERAL MEDICAL CONDITION OR MENTAL DISORDER CHARACTERIZED BY POTENTIALLY EMBARRASSING SYMPTOMS IS PRESENT: Has your avoidance of (SOCIAL SITUATIONS) been related to your (MEDICAL CONDITION OR MENTAL DISORDER)?

IF YES: How have you dealt with your condition?

- The fear, anxiety, or avoidance is not better explained by the symptoms of another mental disorder such as Panic Disorder, Separation Anxiety Disorder, Body Dysmorphic Disorder, or Autism Spectrum Disorder.
- J. If another medical condition (e.g., Parkinson's disease, obesity, disfigurement from burns or injury) [or potentially embarrassing mental disorder] is present, the fear, anxiety, or avoidance is clearly unrelated or is excessive.

SOCIAL ANXIETY DISORDER CRITERIA A, B, C, D, E, F, G, H, I AND J ARE CODED "3."



GO TO *SPECIFIC PHOBIA*

SPECIFIC PHOBIA

SPECIFIC PHOBIA CRITERIA

F88 ▶IF SCREENING QUESTION #5 ANSWERED "NO," SKIP TO *CURRENT GENERALIZED ANXIETY **SCREEN Q#5** DISORDER* F.24. YES → IF QUESTION #5 ANSWERED "YES": IF NO: GO TO *CURRENT You've said that there are other **GENERALIZED ANXIETY** things that have made you especially **DISORDER*** anxious or afraid in the past year, like flying, seeing blood, getting a shot, heights, closed places, or certain kinds of animals or insects... A. Marked fear or anxiety about a specific 2 3 F89 → IF SCREENER NOT USED: Are there any object or situation (e.g., flying, heights, other things that have made you animals, receiving an injection, seeing GO TO especially anxious or afraid in the blood). *CURRENT past year, like flying, seeing blood, **GENERALIZED** getting a shot, heights, closed places, ANXIETY or certain kinds of animals or insects? **DISORDER*** Tell me about that. F90 In the past year, since (1 YEAR AGO), did B. The phobic object or situation almost 2 3 1 you almost always immediately feel always provokes immediate fear or anxiety. frightened or anxious when you were GO TO (CONFRONTED WITH PHOBIC STIMULUS)? *CURRENT **GENERALIZED** ANXIETY **DISORDER*** F91 In the past year, did you go out of your C. The phobic situation(s) is actively avoided, 2 3 way to avoid or endured with intense fear or anxiety. (PHOBIC STIMULUS)? (Are there things you GO TO didn't do because of this fear that you *CURRENT would otherwise have done?) **GENERALIZED** ANXIETY IF NO: How hard was it for you when **DISORDER*** (CONFRONTED WITH PHOBIC STIMULUS)? F92 IF PHOBIC STIMULUS IS POSSIBLY D. The fear or anxiety is out of proportion to 1 2 3 DANGEROUS: How dangerous would vou the actual danger posed by the specific say it actually is to (BE EXPOSED TO PHOBIC object or situation and to the sociocultural GO TO STIMULUS)? context. *CURRENT **GENERALIZED**

NOTE: Code "3" if objects or situations do not

pose danger or if fear or anxiety is out of

proportion to actual danger or sociocultural

Do you think that you have been more

afraid of (PHOBIC STIMULUS) than you

should have been given the actual

danger?

context.

ANXIETY

DISORDER*


IF UNKNOWN: How long have you had these fears? (For 6 months or more?)

E. The fear, anxiety, or avoidance is persistent, typically lasting for 6 months or more.

F. The fear, anxiety, or avoidance causes

areas of functioning.

clinically significant distress or impairment in social, occupational, or other important



IF UNKNOWN: What effect have (PHOBIC SXS) had on your life in the past year?

ASK THE FOLLOWING QUESTIONS <u>AS NEEDED</u> TO RATE CRITERION F:

How have (PHOBIC SXS) affected your relationships with your family, romantic partner or friends?

How have (PHOBIC SXS) affected your work/school? (How about your attendance at work or school?)

How about doing other things that are important to you like religious activities, physical exercise, or hobbies?


IF BLOOD-INJECTION-INJURY TYPE: Have you avoided going to the dentist or doctor because of (PHOBIC SXS)? (How has this affected your health?)


Have (PHOBIC SXS) affected any other important part of your life?

IF HAVE NOT INTERFERED WITH LIFE: How much have you been bothered or upset by having (PHOBIC SXS)?


IF NECESSARY, RETURN TO THIS ITEM AFTER COMPLETING MODULES FOR OC AND RELATED DISORDERS.

G. The disturbance is not better explained by the symptoms of another mental disorder, including fear, anxiety, and avoidance of situations associated with panic like symptoms or other incapacitating symptoms (as in Agoraphobia), objects or situations related to obsessions (as in Obsessive-Compulsive Disorder) reminders of traumatic events (as in Posttraumatic Stress Disorder), separation from home or attachment figures (as in Separation Anxiety Disorder) or social situations (as in Social Anxiety Disorder).





SPECIFIC PHOBIA CRITERIA A, B, C, D, E, F, AND G ARE CODED "3."



GO TO *CURRENT GENERALIZED ANXIETY DISORDER*

*CURRENT GENERALIZED ANXIETY GENERALIZED ANXIETY **DISORDER* DISORDER CRITERIA**

► IF SCREENING QUESTION #6 ANSWERED "NO," SKIP TO *PAST GENERALIZED ANXIETY DISORDER* F.27


→ IF QUESTION #6 ANSWERED "YES": You've said that over the past year you've been feeling anxious and worried for a lot of the time. (Tell me about that.)

IF SCREENER NOT USED: Over the past year, have you been feeling anxious and worried for a lot of the time? (Tell me about that.)

SCREEN Q#6 F111 YES NO GO TO THE NEXT MODULE

What kinds of things have you worried about? (What about your job, your health, your family members, your finances, or other smaller things like being late for appointments?) How much did you worry about (EVENTS OR ACTIVITIES)? What else have you worried about?

A. Excessive anxiety and worry (apprehensive expectation), occurring more days than not for at least 6 of the past 12 months, about a number of events or activities (such as work or school performance).




Have you worried about (EVENTS OR ACTIVITIES) even when there was no reason? (Have you worried more than most people would in your circumstances? Has anyone else thought you worried too much? Have you worried more than you should have given your actual circumstances?)

During the past year, since (1 YEAR AGO), would you say that you have been worrying more days than not?

found that it's hard to stop yourself or to think about anything else?

When you're worrying this way, have you B. The person finds it difficult to control the worry.



F113

F115

Now I am going to ask you some questions about symptoms that often go along with being nervous or worried.

Thinking about those periods since (1 YEAR AGO) when you have been feeling nervous, anxious, or worried...

C. The anxiety and worry are associated with three (or more) of the following six symptoms (with at least some symptoms present for more days than not for at least 6 of the past 12 months):

...have you often felt physically restless, like you couldn't sit still?

- 1. Restlessness or feeling keyed up or on edge.
- F114 1 2

...have you often felt keyed up or on edge?

3=Threshold or true

2. Being easily fatigued.

1=Absent or false

SCID-RV (for DSM-5®) (Version 1.0.0)	Other Specified Anxiety Disorder		Anxie	ety Di	sorders	s F.84
have you often tired easily?		?	1	2	3	
have you often had trouble concentrating or has your mind often gone blank?	Difficulty concentrating or mind going blank.	?	1	2	3	F116
have you often been irritable?	4. Irritability.	?	1	2	3	F117
have your muscles often been tense?	5. Muscle tension.	?	1	2	3	F118
have you often had trouble falling or staying asleep? How about often feeling tired when you woke up because you didn't get a good night's sleep?	Sleep disturbance (difficulty falling or staying asleep, or restless unsatisfying sleep).	?	1	2	3	F119
IF UNCLEAR: Did at least some of these symptoms like (SXS CODED "3") happen for more days than not over a six-month	AT LEAST THREE "C" SXS ARE CODED "3" AND AT LEAST SOME OCCURRED MORE DAYS THAN HOT FOR AT LEAST 6 OF THE PAST 12	?	1	2	3	F120
period in the past year?	MONTHS		GO TO THE NEXT MODULE			
IF UNKNOWN: What effect have (GAD SXS) had on your life in the past year?	D. The anxiety, worry, or physical symptoms cause clinically significant distress or		1	2	3	F121
ASK THE FOLLOWING QUESTIONS <u>AS NEEDED</u> TO RATE CRITERION D:	impairment in social, occupational, or other important areas of functioning.	GO TO THE NEXT MODULE				
How have (CAD SYS) affected your						

How have (GAD SXS) affected your relationships or your interactions with other people? (Have [GAD SXS] caused you any problems in your relationships with your family, romantic partner or friends?)

How have (GAD SXS) affected your work/schoolwork? (How about your attendance at work or school? Have [GAD SXS] made it more difficult to do your work/schoolwork? How have [GAD SXS] affected the quality of your work/schoolwork?)

How have (GAD SXS) affected your ability to take care of things at home? How about doing other things that are important to you like religious activities, physical exercise, or hobbies? Have you avoided doing anything because you felt like you weren't up to it?

Has your anxiety or worry affected any other important part of your life?

IF HAS NOT INTERFERED WITH LIFE: How much have you been bothered or upset by having (GAD SXS)?

IF UNKNOWN: When did (GAD SXS) begin?

Just before you began having (GAD SXS), were you taking any drugs, caffeine, diet pills, or other medicines?

(How much coffee, tea, or caffeinated soda do you drink a day?)

Just before (GAD SXS) began, were you physically ill?

IF YES: What did the doctor say?

E. [Primary Anxiety Disorder:] The disturbance is not attributable to the physiological effects of a substance (e.g., a drug of abuse, a medication) or to another medical condition.

IF THERE IS ANY INDICATION THAT THE ANXIETY MAY BE SECONDARY (I.E., A DIRECT PHYSIOLOGICAL CONSEQUENCE OF GMC OR SUBSTANCE/MEDICATION), GO TO *GMC/SUBSTANCE* F.33 AND RETURN HERE TO MAKE A RATING OF "1" OR "3."

NOTE: Refer to list of etiological medical conditions and substances/medications on page F.4.

F122 **PRIMARY ANXIETY** DISORDER ALL DUE TO GMC OR SUB-STANCE/MED-**ICATION USE** GO TO THE **NEXT MODULE** CONTINUE WITH **NEXT ITEM** F123 3 GO TO THE

IF NECESSARY, RETURN TO THIS ITEM AFTER COMPLETING MODULE FOR OC AND RELATED DISORDERS AND EATING DISORDERS.

F. The disturbance is not better explained by another mental disorder (e.g., anxiety or worry about having a panic attacks in Panic Disorder, negative evaluation in Social Anxiety Disorder, contamination or other obsessions in Obsessive Compulsive Disorder, separation from attachment figures in Separation Anxiety Disorder, gaining weight in Anorexia Nervosa, physical complaints in Somatic Symptom disorder, perceived appearance flaws in Body Dysmorphic Disorder or having a serious illness in Illness Anxiety Disorder, or the content of delusional beliefs in Schizophrenia or Delusional Disorder).

E, AND F ARE CODED "3."

F124 GENERALIZED ANXIETY CRITERIA A, B, C, D, 3 GO TO THE CURRENT **NEXT MODULE** GENERALIZED **ANXIETY DISORDER**

NEXT MODULE

GO TO THE NEXT MODULE F126

GMC/SUBSTANCE AS ETIOLOGY FOR ANXIETY SYMPTOMS

ANXIETY DISORDER DUE TO ANOTHER MEDICAL CONDITION

ANXIETY DISORDER DUE TO ANOTHER MEDICAL CONDITION CRITERIA


IF SYMPTOMS NOT TEMPORALLY ASSOCIATED WITH A GENERAL CONDITION CHECK HERE ____ AND GO TO *SUBSTANCE/MEDICATION-INDUCED ANXIETY DISORDER* F.35

F147

F149

CODE BASED ON INFORMATION ALREADY OBTAINED

- A. Panic attacks or anxiety is predominant in the clinical picture.
- ? 1 3 F148
- B/C. There is evidence from this history, physical examination, or laboratory findings that the disturbance is the direct physiological consequence of another medical condition AND the disturbance is not better accounted for by another mental disorder.



Did the (ANXIETY SXS) start or get much worse only after (GMC) began? How long after (GMC) began did (ANXIETY SXS) start or get much worse?

NOTE: The following factors should be considered and, if present, support the conclusion that a general medical condition is etiologic to the anxiety symptoms.

IF GMC HAS RESOLVED: Did the (ANXIETY SXS) get better once the (GMC) got better?

- There is evidence from the literature of a well-established association between the general medical condition and the anxiety symptoms. (Refer to list of etiological general medical conditions on page F.4.)
- There is a close temporal relationship between the course of the anxiety symptoms and the course of the general medical condition.
- The anxiety symptoms are characterized by unusual presenting features (e.g., late ageat-onset).
- The absence of alternative explanations (e.g., anxiety symptoms as a psychological reaction to the stress of being diagnosed with a general medical condition).

IF UNKNOWN: What effect did (ANXIETY SXS) have on your life?

ASK THE FOLLOWING QUESTIONS <u>AS NEEDED</u> TO RATE CRITERION E:

How did (ANXIETY SXS) affect your relationships or your interactions with other people? (Did [ANXIETY SXS] cause you any problems in your relationships with your family, romantic partner or friends?)

How did (ANXIETY SXS) affect your school/work? (How about your attendance at work or school? Did [ANXIETY SXS] make it more difficult to do your work/schoolwork? How did [ANXIETY SXS] affect the quality of your work/schoolwork?)


How did (ANXIETY SXS) affect your ability to take care of things at home? How about doing other things that are important to you like religious activities, physical exercise, or hobbies? Did you avoid doing anything because you felt like you weren't up to it?


Did your anxiety or worry affect any other important part of your life?

IF HAS NOT INTERFERED WITH LIFE: How much were you bothered or upset by having (ANXIETY SXS)?

E. The disturbance causes clinically significant distress or impairment in social, occupational, or other important areas of functioning.

NOTE: The D criterion (delirium rule-out) has been omitted.





*SUBSTANCE/MEDICATION-**INDUCED ANXIETY DISORDER***

SUBSTANCE/MEDICATION-INDUCED ANXIETY DISORDER CRITERIA

F154

IF SYMPTOMS NOT TEMPORALLY ASSOCIATED WITH SUBSTANCE/MEDICATION USE, CHECK AND RETURN TO DISORDER BEING EVALUATED, CONTINUING WITH THE ITEM FOLLOWING "SYMPTOMS ARE NOT ATTRIBUTABLE TO THE PHYSIOLOGICAL EFFECTS OF A SUBSTANCE OR ANOTHER MEDICAL CONDITION" (SEE PAGE NUMBERS IN BOX TO THE RIGHT).

CODE BASED ON INFORMATION ALREADY **OBTAINED**

IF NOT KNOWN: When did the (ANXIETY SXS) begin? Were you already using (SUBSTANCE/MEDICATION) or had you just stopped or cut down your use?

IF UNKNOWN: How much (SUBSTANCE/ MEDICATION) were you using when you began to have (ANXIETY SXS)?

- A. Panic attacks or anxiety is predominant in the clinical picture.
- B. There is evidence from the history, physical examination, or laboratory findings of both (1) and (2):
 - 1. The symptoms in criterion A developed during or soon after substance intoxication or withdrawal or exposure to a medication.
 - 2. The involved substance/ medication is capable of producing the symptoms in Criterion A.

NOTE: Refer to list of substances/medications on page F.4.

ASK ANY OF THE FOLLOWING QUESTIONS AS C. The disturbance is NOT better accounted NEEDED TO RULE OUT A NON-SUBSTANCE-INDUCED ETIOLOGY:

IF UNKNOWN: Which came first, the (SUBSTANCE/MEDICATION USE) or the (ANXIETY SXS)?

IF UNKNOWN: Have you had a period of time when you stopped using (SUBSTANCE/MEDICATION)?

IF YES: After you stopped using (SUBSTANCE/MEDICATION) did the (ANXIETY SXS) go away or get better?

IF YES: How long did it take for them to get better? Did they go away within a month of stopping?


IF UNKNOWN: Have you had any other episodes of (ANXIETY SXS)?

IF YES: How many? Were you using (SUBSTANCE/MEDICATION) at those times?

for by an anxiety disorder that is not substance-induced. Such evidence of an independent anxiety disorder could include the following:

NOTE: The following three statements constitute evidence that the anxiety symptoms are not substance-induced. Code "1" if any are true. Code "3" only if none are true.

- 1) The symptoms precede the onset of the substance/medication use;
- 2) The symptoms persist for a substantial period of time (e.g., about 1 month) after the cessation of acute withdrawal or severe intoxication; or
- 3) There is other evidence suggesting the existence of an independent nonsubstance/ medication-induced anxiety disorder (e.g., a history of recurrent nonsubstance/ medication-related episodes).



F157 1 3 NOT SUBSTANCE INDUCED RETURN TO DISORDER BEING **EVALUATED**

IF UNKNOWN: What effect did (ANXIETY SXS) have on your life?

ASK THE FOLLOWING QUESTIONS <u>AS NEEDED</u> TO RATE CRITERION E:

How did (ANXIETY SXS) affect your relationships or your interactions with other people? (Did [ANXIETY SXS] cause you any problems in your relationships with your family, romantic partner or friends?)

How did (ANXIETY SXS) affect your work/schoolwork? (How about your attendance at work or school? Did [ANXIETY SXS] make it more difficult to do your work/schoolwork? How did [ANXIETY SXS] affect the quality of your work/schoolwork?)


How did (ANXIETY SXS) affect your ability to take care of things at home? How about doing other things that are important to you like religious activities, physical exercise, or hobbies? Did you avoid doing anything because you felt like you weren't up to it?


Did your anxiety or worry affect any other important part of your life?

IF HAS NOT INTERFERED WITH LIFE: How much were you bothered or upset by having (ANXIETY SXS)?

E. The symptoms cause clinically significant distress or impairment in social, occupational, or other important areas of functioning.

NOTE: The D criterion (delirium rule-out) has been omitted.





OG. OBSESSIVE-COMPULSIVE AND RELATED DISORDERS

*OBSESSIVE-COMPULSIVE **DISORDER***

OBSESSIVE-COMPULSIVE DISORDER CRITERIA

► IF SCREENING OUESTIONS #8, #9, AND #10 ARE ALL ANSWERED "NO" SKIP TO *COMPULSIONS* G.2, (NOTE: BECAUSE SOME SUBJECTS WITH OCD MAY BE RELUCTANT TO CONFIDE THEIR OBSESSIONS DURING THE SCREENING, CONSIDER RE-ASKING SCREENING QUESTIONS BELOW AT THIS POINT IN THE SCID.)

▶ IF QUESTION #8 ANSWERED "YES": In the past year, you've said that you've been bothered by thoughts that kept coming back to you even when you didn't want them to, like being exposed to germs or dirt or needing everything to be lined up in a certain way. What were they?

▶ IF QUESTION #9 ANSWERED "YES": In the past year, you've [also] said that you've had images pop into your head that you didn't want like violent or horrible scenes or something of a sexual nature. What were they?

▶ IF QUESTION #10 ANSWERED "YES": In the past year, you've [also] said that you've had urges to do something that kept coming back to you even though you didn't want them to, like an urge to harm a loved one. What were they?

▶ IF SCREENER NOT USED: In the past year, since (1 YEAR AGO), have you been bothered by thoughts that kept coming back to you even when you

How about having images pop into your head that you didn't want like violent or horrible scenes or something of a sexual nature? (What were they?)

didn't want them to, like being exposed to germs or dirt or needing everything to be lined up in a certain

way? (What were they?)

How about having urges to do something that kept coming back to you even though you didn't want them to, like an urge to harm a loved one? (What were they?)

IF YES TO ANY OF ABOVE: Have these (THOUGHTS/IMAGES/URGES) made you very anxious or upset in the past year?

In the past year, since (1 YEAR AGO), when you had these (THOUGHTS/IMAGES/ URGES) did you try hard to get them out of your head? (What would you try to do?)

A. Presence of obsessions, compulsions, or

Obsessions are defined by (1) and (2):


1. Recurrent and persistent thoughts, urges, or images that are experienced, at some time during the disturbance, as intrusive and unwanted, and that in most individuals cause marked anxiety or distress.

NO OBSESSIONS GO TO *COMPULSIONS* G.2


1

2

2. The individual attempts to ignore or suppress such thoughts, urges, or images, or to neutralize them with some other thought or action (i.e., by performing a compulsion).



DESCRIBE CONTENT OF OBSESSION(S):



G4

COMPULSIONS

→ IF SCREENING QUESTION #11 ANSWERED "NO," GO TO *SKIP OUT IF NEITHER OBSESSIONS NOR COMPULSIONS* G.3 (NOTE: BECAUSE SOME SUBJECTS WITH OCD MAY BE RELUCTANT TO CONFIDE THEIR COMPULSIONS DURING THE SCREENING, CONSIDER RE-ASKING SCREENING QUESTION BELOW AT THIS POINT IN THE SCID.)

→ IF QUESTION #11 ANSWERED "YES": In the past year, since (1 YEAR AGO), you've said that there were things you had to do over and over again and were hard to resist doing, like washing your hands again and again, repeating something over and over again until it "felt right," counting up to a certain number, or checking something many times to make sure that you'd done it right. Tell me about that.

IF SCREENER NOT USED: In the past year, since (1 YEAR AGO), was there anything that you had to do over and over again and was hard to resist doing, like washing your hands again and again, repeating something over and over again until it "felt right," counting up to a certain number, or checking something many times to make sure that you'd done it right?

Tell me about that. (What did you have to do?)

IF UNCLEAR: Why did you have to do (COMPULSIVE ACT)? What would happen if you didn't do it?

IF UNCLEAR: How many times would you do (COMPULSIVE ACT)? Have you been doing (COMPULSIVE ACT) more than really made sense?


GO TO *SKIP OUT IF NEITHER
OBSESSIONS NOR COMPULSIONS* G.3
(TOP OF NEXT PAGE)

Compulsions are defined by (1) and (2):

- Repetitive behaviors (e. g., hand washing, ordering, checking) or mental acts (e.g., praying, counting, repeating words silently) that the individual feels driven to perform in response to an obsession, or according to rules that must be applied rigidly.
- The behaviors or mental acts are aimed at preventing or reducing anxiety or distress, or preventing some dreaded event or situation; however these behaviors or mental acts either are not connected in a realistic way with what they are designed to neutralize or prevent, or are clearly excessive.

SCREEN Q#11
YES | NO

IF NO: GO TO
*SKIP OUT IF
NEITHER
OBSESSIONS NOR
COMPULSIONS*
G.3



DESCRIBE CONTENT OF COMPULSION(S):

SKIP OUT IF NEITHER OBSESSIONS NOR COMPULSIONS

→ IF EITHER OBSESSIONS OR COMPULSIONS, OR BOTH, CONTINU	JE BELOW.	
■ IF NEITHER OBSESSIONS NOR COMPULSIONS, CHECK HERE	AND GO TO THE NEXT MODULE.	G9

IF UNKNOWN: How much time do you
spend on (OBSESSION OR COMPULSION) in
the past year?

IF UNKNOWN: What effect did these (OBSESSIONS OR COMPULSIONS) have on your life in the past year?

ASK THE FOLLOWING QUESTIONS <u>AS NEEDED</u> TO RATE CRITERION B:

How have (OBSESSIONS OR COMPULSIONS) affected your relationships or your interactions with other people in the past year? (Have they caused you any problems in your relationships with your family, romantic partner, roommates or friends?)

How have (OBSESSIONS OR COMPULSIONS) affected your work/school in the past year? (How about your attendance at work or school? Have [OBSESSIONS OR COMPULSIONS] made it more difficult to do your work/ schoolwork)? How have (OBSESSIONS OR COMPULSIONS) affected the quality of your work/schoolwork?)

How have (OBSESSIONS OR COMPUSIONS) affected your ability to take care of things at home in the past year? How about doing other things that are important to you like religious activities, physical exercise, or hobbies?

Have (OBSESSIONS OR COMPULSIONS) affected any other important part of your life in the past year?

IF HAVE NOT INTERFERED WITH LIFE: How much have you been bothered in the past year by having (OBSESSIONS OR COMPULSIONS)?

B. The obsessions or compulsions are time consuming (e.g., take more than 1 hour per day) or cause clinically significant distress or impairment in social, occupational, or other important areas of functioning.

IF UNKNOWN: When did (OBSESSIONS OR COMPULSIONS) begin?

Just before this began, were you physically ill?

IF YES: What did the doctor say?

Just before this began, were you using any medications?

IF YES: Any change in the amount you were using?

Just before this began, were you drinking or using any drugs?


C. [Primary Obsessive-Compulsive Disorder.] The obsessive-compulsive symptoms are not attributable to the physiological effects of a substance/medication or to another medical condition.


IF THERE IS ANY INDICATION THAT THE OBSESSIONS OR COMPULSIONS MAY BE SECONDARY (I.E., A DIRECT PHYSIOLOGICAL CONSEQUENCE OF GMC OR SUBSTANCE), GO TO *GMC/SUBSTANCE*G.11 AND RETURN HERE TO MAKE A RATING OF "1" OR "3."

<u>Etiological medical conditions include</u>: Sydenham's chorea, medical conditions leading to striatal damage, such as cerebral infarction.


<u>Etiological substances/medications include</u>: intoxication with cocaine, amphetamines or other stimulants and exposure to heavy metals.

D. The disturbance is not better explained by the symptoms of another mental disorder (e.g., excessive worries, as in Generalized Anxiety Disorder; preoccupation with appearance, as in Body Dysmorphic Disorder; difficulty discarding or parting with possessions, as in Hoarding Disorder; hair pulling, as in Trichotillomania; skin picking, as in Excoriation Disorder; stereotypies, as in Stereotypic Movement Disorder; ritualized eating behavior, as in Eating Disorders; preoccupation with substances or gambling, as in Substance-Related and Addictive Disorders; preoccupation with having an illness, as in Illness Anxiety Disorder; sexual urges or fantasies, as in Paraphilic Disorders; impulses, as in Disruptive, Impulse-Control, and Conduct Disorders; guilty ruminations, as in Major Depressive Disorder; thought insertion or delusional preoccupations, as in Schizophrenia Spectrum and Other Psychotic Disorders; or repetitive patterns of behavior, as in Autism Spectrum Disorder).





OBSESSIVE COMPULSIVE DISORDER CRITERIA A, B, C, D, AND E ARE CODED "3."



GO TO THE NEXT MODULE.

I. FEEDING AND EATING DISORDERS

ANOREXIA NERVOSA

ANOREXIA NERVOSA CRITERIA

➤ IF SCREENING QUESTION #12 ANSWERED "NO," CHECK HERE ____ AND SKIP TO *BULIMIA NERVOSA* I.4


► IF QUESTION #12 ANSWERED "YES": You've said that there was a time in the

past year when you weighed much less than other people thought you ought to weigh...

IF SCREENER NOT USED: Now I would like to ask you some questions about your eating habits and your weight in the past year. Have you had a time when you weighed much less than other people thought you ought to weigh?

IF YES: Why was that? How much did you weigh? How old were you then? How tall were you?

A. Restriction of energy intake relative to requirements, leading to a significantly low body weight in the context of age, sex, developmental trajectory, and physical health. Significantly low weight is defined as a weight that is less than minimally normal or, for children and adolescents, less than minimally expected.



3 I4

15

2

1

GO TO

BULIMIA NERVOSA

IF PAST YEAR RATING OF "3": During the past 3 months, since (3 MONTHS AGO), what is the lowest your weight has been?

In the past year, since (1 YEAR AGO), were you very afraid that you could become fat?

IF NO: Tell me about your eating habits. (Have you avoided high calorie foods or high fat foods? How strict are you about it? Have you ever thrown up after you eaten? How often? Do you exercise a lot after you eat?)

IF PAST YEAR RATING OF "3": Has this also been the case during the past 3 months, since (3 MONTHS AGO)?


At your lowest weight in the past year, did you still feel too fat or that part of your body was too fat?


IF NO: Did you need to be very thin in order to feel better about yourself?

IF NO AND LOW WEIGHT IS MEDICALLY SERIOUS: When you were that thin, did anybody tell you it could be dangerous to your health to be that thin? (What did you think?)

IF PAST YEAR RATING OF "3": Has this also been the case in the past 3 months, since (3 MONTHS AGO)?

B. Intense fear of gaining weight or of becoming fat, or persistent behavior that interferes with weight gain, even though underweight.





Past 3 months

1 2

ANOREXIA NERVOSA CRITERIA A, B, AND C ARE CODED "3"

IF NO: GO TO
*BULIMIA
NERVOSA*

ANOREXIA
NERVOSA

SCID-RV (for DSM-5®) (Version 1.0.0) Anorexia Nervosa Feeding and Eating Disorders I

When did you last have (ANY SXS OF ANOREXIA NERVOSA)? Number of months prior to interview when last had a symptom of Anorexia Nervosa — — — — I12

Do you have eating binges in which you eat a lot of food in a short period of time and feel that your eating is out of control? (How often?)

IF NO: What kinds of things have you done to keep weight off? (Do you ever make yourself vomit or take laxatives, enemas, or water pills? How often?)

Specify **subtype** for current episode: (circle the appropriate number)

1 - Restricting type:

During the last 3 months, the individual has NOT engaged in recurrent episodes of binge eating or purging behavior (i.e., self-induced vomiting or the misuse of laxatives, diuretics, or enemas). This subtype describes presentations in which weight loss is accomplished primarily through dieting, fasting and/or excessive exercise.

2 - Binge-eating/purging type:

During last 3 months, the individual has engaged in recurrent episodes or binge-eating or purging behavior (i.e., self-induced vomiting or misuse of laxatives, diuretics, or enemas).

I14


BULIMIA NERVOSA

BULIMIA NERVOSA CRITERIA

- → IF SCREENING QUESTION #13 IS ANSWERED "NO," GO TO THE NEXT MODULE.
- IF OUESTION #13 ANSWERED "YES": In the past year, since (1 YEAR AGO), you've said that you've had eating binges, that is, times when you couldn't resist eating a lot of food or stop eating once you've started. Tell me about those times.
- ► IF SCREENER NOT USED: In the past year, since (1 YEAR AGO), have you had eating binges, that is, times when you couldn't resist eating a lot of food or stop eating once you started? Tell me about those times.
- A. Recurrent episodes of binge eating. An episode of binge eating is characterized by BOTH of the following:

During these times, were you unable to control what or how much you were eating?

2. A sense of lack of control over eating during the episode (e.g., a feeling that one cannot stop eating or control what or how much one is eating)



SCREEN Q#13 YES | |

GO TO THE NEXT


MODULE

NO

I15


NOTE: Criterion A.2 (lack of control) precedes criterion A.1 to tie in with screening question.

- During those times in the past year, how much did you eat? Over what period of time? What's the most you might eat at such times? (Does this only happen during celebrations or holidays?)
- 1. Eating, in a discrete period of time (e.g., within any 2-hour period), an amount of food that is definitely larger than most people would eat during a similar period of time and under similar circumstances



CRITERIA A.2 AND A.1 ARE CODED "3"

IF PAST YEAR RATING OF "3" FOR BOTH CRITERIA A.2 AND A.1: During the past 3 months, since (3 MONTHS AGO), have you had such episodes?



In the past year, have you done anything to keep yourself from gaining weight because of the binge eating (like making yourself vomit, taking laxatives, enemas, water pills, or thyroid hormone, strict dieting or fasting, or exercising a lot)? Tell me about that. How often did this occur?

IF PAST YEAR RATING OF "3": Have you done (COMPENSATORY BEHAVIOR[S]) during the past 3 months, since (3 MONTHS AGO)?

In the past year, how often were you binge eating and (COMPENSATORY BEHAVIOR[S])? (At least once a week for at least 3 months?)

C. The binge eating and inappropriate compensatory behaviors both occur, on average, at least once a week for 3 months.

B. Recurrent inappropriate compensatory behavior in order to prevent weight gain, such as: selfinduced vomiting; misuse of laxatives, diuretics, enemas, or other medications; fasting; or excessive exercise. ? 1 2 3 I20
GO TO THE NEXT MODULE

Past 3 months 1 21

? 1 2 3 I22
GO TO THE NEXT MODULE

Past 3 months 1 3

AGO), how often were you binge eating and (COMPENSATORY BEHAVIOR[S])? At least once a week?

IF PAST YEAR RATING OF "3": Since (3 MONTHS

In the past year, has your body shape and weight ever been an important factor in how you felt about yourself?

IF YES: How important?

IF PAST YEAR RATING OF "3": Has this also been the case during the past 3 months?

IF UNKNOWN: In the past year, did you binge eat and then (ENGAGE IN COMPENSATORY BEHAVIOR) only when your weight was very low?

D. Self-evaluation is unduly influenced by body shape and weight.

IF UNKNOWN: **In the past year, did you binge** E. The disturbance does not occur exclusively **eat and then** (ENGAGE IN COMPENSATORY during episodes of Anorexia Nervosa.

? 1 2 3 I24

Past 3 months 125
1 3 I26

GO TO THE NEXT

MODULE

BULIMIA NERVOSA CRITERIA A, B, C, D, AND E ARE CODED "3."




TABLE FOR DETERMINING SEVERITY OF ANOREXIA NERVOSA BASED ON BODY MASS INDEX

Anorexia Nervosa Severity	Mild (BMI≥17)	Moderate (BMI=16-16.99)	Severe (BMI=15-15.99)	Extreme (BMI=<15)
Height cms (inches/feet)	Body Weight kg (pounds)	Body Weight kg (pounds)	Body Weight kg (pounds)	Body Weight kg (pounds)
148 (58" / 4′10")	≥38 (≥84)	35-37 (77-82)	33-34 (72-76)	<33 (<72)
150 (59" / 4′11")	≥39 (≥86)	37-38 (79-81)	35-36 (74-78)	<35 (<74)
153 (60" / 5′)	≥40 (≥90)	38-39 (84-87)	36-37 (77-81)	<36 (<77)
155 (61" / 5´1")	≥41 (≥95)	39-40 (86-90)	37-38 (80-85)	<37 (<80)
158 (62" / 5´2")	≥43 (≥95)	41-42 (89-93)	38-39 (82-88)	<38 (<82)
160 (63" / 5′3")	≥44 (≥97)	42-43 (92-96)	39-40 (85-91)	<39 (<85)
163 (64" / 5 ′ 4")	≥46 (≥101)	44-45 (97-99)	40-41 (88-92)	<40 (<88)
165 (65" / 5´5")	≥47 (≥104)	45-46 (100-102)	41-43 (91-95)	<41 (<91)
168 (66" / 5′6")	≥48 (≥106)	46-47 (100-105)	43-44 (93-99)	<43 (<93)
170 (67" / 5´7")	≥49 (≥108)	47-48 (103-107)	44-46 (95-102)	<44 (<95)
173 (68" / 5′8")	≥51 (≥112)	49-50 (104-109)	46-47 (97-103)	<46 (<97)
175 (69" / 5′9")	≥52 (≥115)	50-51 (106-113)	47-48 (99-105)	<47 (<99)
178 (70" / 5′10")	≥54 (≥119)	52-53 (109-116)	48-50 (102-108)	<48 (<102)
180 (71" / 5´11")	≥55 (≥121)	53-54 (115-123)	51-52 (108-114)	<51 (<108)
183 (72" / 6′0")	≥57 (≥126)	54-55 (119-125)	52-53 (111-118)	<52 (<111)
185 (73" / 6 ´1")	≥58 (≥128)	55-57 (124-129)	53-54 (114-121)	<53 (<114)
188 (74" / 6´2")	≥60 (≥132)	57-59 (125-132)	54-55 (117-124)	<54 (<117)
191 (75" / 6′3")	≥61 (≥134)	59-60(128-136)	55-58 (122-127)	<55 (<122)
193 (76" / 6 ′ 4")	≥63 (≥140)	60-62 (132-140)	58-59 (123-131)	<58 (<123)
Severity	Mild	Moderate	Severe	Extreme

Source: Adapted from Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults: The Evidence Report.

GMC/SUBSTANCE CAUSING OBSESSIVE-COMPULSIVE AND RELATED SYMPTOMS

OBSESSIVE-COMPULSIVE AND OBSESSIVE-COMPULSI RELATED DISORDER DUE TO DUE TO ANOTHER MED ANOTHER MEDICAL CONDITION CONDITION CRITERIA

OBSESSIVE-COMPULSIVE AND RELATED DISORDER DUE TO ANOTHER MEDICAL CONDITION CRITERIA


IF SYMPTOMS NOT TEMPORALLY ASSOCIATED WITH A GENERAL MEDICAL CONDITION, CHECK HERE ___ AND GO TO *SUBSTANCE-INDUCED OC AND RELATED DISORDER* G.14.

G28

CODE BASED ON INFORMATION ALREADY OBTAINED

A. Obsessions, compulsions, preoccupations with appearance, hoarding, skin picking, hair pulling, other body-focused repetitive behaviors, or other symptoms characteristic of obsessive-compulsive and related disorder predominate in the clinical picture. ? 1 2 3 G29
GO TO
*SUBSTANCE
INDUCED*
G.14

B/C. There is evidence from the history, physical examination, or laboratory findings that the disturbance is the direct physiological consequence of another medical condition AND the disturbance is not better accounted for by another mental disorder.



Did (OC AND RELATED SXS) start or get much worse only after (GMC) began? How long after (GMC) began did (OC AND RELATED SXS) start or get much worse?

IF GMC HAS RESOLVED: Did the (OC AND RELATED SYMPTOMS) get better once the (GMC) got better?

NOTE: The following factors should be considered and, if present, support the conclusion that a general medical condition is etiologic to the obsessive-compulsive and related symptoms.

- There is evidence from the literature of a well-established association between the general medical condition and the obsessive-compulsive and related symptoms. (Refer to list of etiological general medical conditions on page G.4.)
- There is a close temporal relationship between the course of the obsessivecompulsive and related symptoms and the course of the general medical condition.
- The obsessive-compulsive and related symptoms are characterized by unusual presenting features (e.g., late age-atonset).
- 4) The absence of alternative explanations (e.g., obsessive-compulsive and related symptoms as a psychological reaction to the stress of being diagnosed with a general medical condition).

IF UNKNOWN: What effect have (OC-RELATED SXS) had on your life?

ASK THE FOLLOWING QUESTIONS <u>AS NEEDED</u> TO RATE CRITERION E.:

How have (OC-RELATED SXS) affected your relationships or your interactions with other people? (Have they caused you any problems in your relationships with your family, romantic partner or friends?)

How have (OC-RELATED SXS) affected your work/school? (How about your attendance at work or school? Have [OC-RELATED SXS] made it more difficult to do your work/schoolwork)? How have [OC-RELATED SXS] affected the quality of your work/schoolwork?)


How have (OC-RELATED SXS) affected your ability to take care of things at home? What about being involved in things that are important to you, like religious activities, physical exercise, or hobbies? Have you avoided situations or people because you didn't want other people to see you doing (OC-RELATED BEHAVIORS)?

Have (OC-RELATED SXS) affected any other important part of your life?


IF HAVE NOT INTERFERED WITH LIFE: How much have your (OC-RELATED SXS) bothered or upset you?

E. The disturbance causes clinically significant distress or impairment in social, occupational, or other important areas of functioning

NOTE: The D criterion (delirium rule-out) has been omitted.



G31



SUBSTANCE-/MEDICATION-INDUCED OC SUBSTANCE-/MEDICATION-INDUCED OC AND RELATED DISORDER AND RELATED DISORDER CRITERIA

IF SYMPTOMS NOT TEMPORALLY ASSOCIATED WITH SUBSTANCE/MEDICATION USE (OR IF SYMPTOMS CONFINED TO HOARDING), CHECK HERE AND RETURN TO EPISODE BEING EVALUATED, CONTINUING WITH THE ITEM FOLLOWING "SYMPTOMS ARE NOT ATTRIBUTABLE TO THE PHYSIOLOGICAL EFFECTS OF A SUBSTANCE OR ANOTHER MEDICAL CONDITION" (SEE PAGE NUMBERS IN BOX TO THE RIGHT).

CODE BASED ON INFORMATION ALREADY OBTAINED.

A. Obsessions, compulsions, skin picking, hair pulling, other body-focused repetitive behaviors, or other symptoms characteristic of the obsessive-compulsive and related disorders predominate in the clinical picture.

EPISODE BEING EVALUATED: OCD G.4 G35

> G36 1 2 3

> > G37

IF NOT KNOWN: When did the (OC AND RELATED SXS) begin? Were you already using (SUBSTANCE/MEDICATION) or had you just stopped or cut down your use?

IF UNKNOWN: How much (SUBSTANCE/ MEDICATION) were you using when you began to have (OC AND RELATED SXS)?

- B. There is evidence from the history, physical examination, or laboratory findings of both (1) and (2):
 - 1. The symptoms in criterion A developed during or soon after substance intoxication or withdrawal or exposure to a medication
 - 2. The involved substance/ medication is capable of producing the symptoms in Criterion A

1 2 3 NOT **SUBSTANCE** INDUCED **RETURN TO EPISODE BEING EVALUATED**

NOTE: Refer to list of etiological substances/medications on page G.4.

ASK ANY OF THE FOLLOWING QUESTIONS AS NEEDED TO RULE OUT A NON-SUBSTANCE-INDUCED ETIOLOGY.

IF UNKNOWN: Which came first, the (SUBSTANCE/MEDICATION USE) or the (OC AND RELATED SXS)?

IF UNKNOWN: Have you had a period of time when you stopped using (SUBSTANCE/MEDICATION)?

IF YES: After you stopped using (SUBSTANCE/MEDICATION) did the (OC AND RELATED SXS) go away or get better?

IF YES: How long did it take for them to get better? Did they go away within a month of stopping?

IF UNKNOWN: Have you had any other episodes of (OC AND RELATED SXS)?

IF YES: How many? Were you using (SUBSTANCE/ MEDICATION) at those times?


C. The disturbance is NOT better accounted for by an obsessive-compulsive and related disorder that is not substanceinduced. Such evidence of an independent obsessive-compulsive disorder and related disorder could include the following:

NOTE: The following three statements constitute evidence that the anxiety symptoms are not substance-induced. Code "1" if any are true. Code "3" only if none are true.

The symptoms precede the onset of the substance/medication use;

The symptoms persist for a substantial period of time (e.g., about 1 month) after the cessation of acute withdrawal or severe intoxication:

There is other evidence suggesting the existence of an independent nonsubstance/medication-induced obsessivecompulsive and related disorder (e.g., a history of recurrent non-substance/ medication-related episodes).



IF UNKNOWN: What effect have (OC-RELATED SXS) had on your life?

ASK THE FOLLOWING QUESTIONS <u>AS NEEDED</u> TO RATE CRITERION E:

How have (OC-RELATED SXS) affected your relationships or your interactions with other people? (Have they caused you any problems in your relationships with your family, romantic partner or friends?)

How have (OC-RELATED SXS) affected your work/school? (How about your attendance at work or school? Have [OC-RELATED SXS] made it more difficult to do your work/schoolwork)? How have [OC-RELATED SXS] affected the quality of your work/schoolwork?)


How have (OC-RELATED SXS) affected your ability to take care of things at home? What about being involved in things that are important to you like religious activities, physical exercise, or hobbies? Have you avoided situations or people because you didn't want other people to see you doing (OC-RELATED BEHAVIOR)?


Have (OC-RELATED SXS) affected any other important part of your life?

IF HAVE NOT INTERFERED WITH LIFE: How much have your (OC-RELATED SXS) bothered or upset you?

E. The symptoms cause clinically significant distress or impairment in social, occupational, or other important areas of functioning.

NOTE: The D criterion (delirium rule-out) has been omitted.





INTERMITTENT EXPLOSIVE DISORDER

INTERMITTENT EXPLOSIVE DISORDER **CRITERIA**

► IF SCREENING QUESTIONS #15a AND #15b ARE BOTH ANSWERED "NO," GO TO next section.

► IF SCREENING QUESTION #15a IS ANSWERED "YES": You've said that in the past year have frequently lost control of your temper and ended up yelling or getting into arguments with others. Tell me about that.

IF SCREENING QUESTION #15b IS ANSWERED "YES": You've (also) said that in the past year, you have lost your temper so that you shoved, hit, kicked or threw something at a person or an animal or damaged someone's property. Tell me about that.

IF SCREENER NOT USED: In the past year, since (1 YEAR AGO), have you frequently lost control of your temper and ended up yelling or getting into arguments with others? (Tell me about that.)

IF NO: In the past year, have you lost your temper so that you shoved, hit, kicked or threw something at a person or an animal or damaged someone's property? (Tell me about that.)

IF THERE IS NO EVIDENCE THAT THE SUBJECT HAS HAD VERBAL OR PHYSICAL AGGRESSION, CHECK HERE ____ AND GO TO next section.

IF UNKNOWN: In the past year, have your A. Recurrent behavioral outbursts angry outbursts resulted in someone getting physically hurt? (Tell me about that.)

IF UNKNOWN: In the past year, have you physically injured an animal in anger?

IF UNKNOWN: In the past year, have your outbursts resulted in damaging things, breaking things, smashing windows, punching a hole in a wall, or other damage to property?

IF YES TO ANY OF THESE: During the past year have you had at least 3 such outbursts?

impulses as manifested by either of the


representing a failure to control aggressive

2. Three behavioral outbursts involving damage or destruction of property and/or physical assault involving physical injury against animals or other individuals occurring within a 12-month period.

NOTE: Physical injury includes, at a minimum, a scratch or bruise, whether or not medical attention is sought.

DESCRIBE:

following:



OK3

OK4 2

?=inadequate information

1=absent or false

2=subthreshold

3=threshold or true

IF UNKNOWN: In the past year, have you had angry outbursts in which you shoved, kicked, hit, or threw something without anything or anyone being damaged or injured?

IF UNKNOWN: In the past year have you also had angry outbursts that involved heated arguments, yelling at people, having temper tantrums, or going on "rants," but without physically hurting anyone or damaging anything?

IF YES TO EITHER: If you were to include all the kinds of angry outbursts that we just talked about in the past year (both verbal and physical), did they altogether ever happen as often as twice a week, on average, for at least 3 months?

 Verbal aggression (e.g., temper tantrums, tirades, verbal arguments or fights) or physical aggression toward property, animals, or other individuals, occurring twice weekly, on average, for a period of 3 months. The physical aggression does not result in damage or destruction of property and does not result in physical injury to animals or other individuals.

Check if:

- Verbal aggression (e.g., tantrums, tirades, verbal arguments or fights) twice weekly for 3 months
- Physical aggression without damage or destruction of property (e.g., throwing clothes or books around that do not get damaged) twice weekly for 3 months

EITHER CRITERION A.2 OR A.1 IS CODED "3"

1 3 OK8

GO TO THE

NEXT

MODULE

2

1

3

OK5

OK6


OK7

What kinds of things have set you off? (Do you think your reactions have been much stronger than they should have been given the circumstances? Has anyone told you that your reactions were way off-base given the situation in question?)

IF UNCLEAR: Have all of these outbursts been "on purpose," that is, in order to intimidate someone or force someone to give you what you want?

- B. The magnitude of aggressiveness expressed during the recurrent outbursts is grossly out of proportion to the provocation or to any precipitating psychosocial stressors.
- C. The recurrent aggressive outbursts are not premeditated (i.e., they are impulsive and/or anger-based) and are not committed to achieve some tangible objective (e.g., money, power, intimidation).

NOTE: Code "1" if all outbursts are premeditated or intended to achieve a tangible objective.



IF UNKNOWN: What effect have your outbursts had on your life in the past year?

ASK THE FOLLOWING QUESTIONS <u>AS NEEDED</u> TO RATE CRITERION D:

Have you gotten into trouble because of them? (For example, has anyone called the police or a supervisor because of these outbursts? Have you ever been arrested as a result of your outbursts? Have you ever had to pay a lot of money to compensate someone for the damage you caused?)

How have your outbursts affected your relationships or your interactions with other people? (Have they caused you any problems in your relationships with your family, romantic partner or friends?)

How have they affected your work/school? (How about getting fired from a job or expelled from school or getting "written up" for disciplinary action because of your outbursts?)

Have your outbursts affected any other important part of your life?


IF DOES NOT INTERFERE WITH LIFE: How much have you been bothered or upset by your outbursts?

IF HX OF MANIA, DEPRESSION, OR PSYCHOSIS: Did these outbursts happen only when you were feeling excited, irritable, or depressed, or only when you were having (PSYCHOTIC SXS)?

IF HX OF PTSD: Did you have any outbursts like this prior to exposure to (TRAUMATIC EVENT)?

IF HX OF ADHD: Have you gotten any treatment specifically for the aggressive outbursts?


D. The recurrent aggressive outbursts cause either marked distress in the individual or impairment in occupational or interpersonal functioning, or are associated with financial or legal consequences.



NOTE: Criterion E regarding minimum chronological age has been omitted.

F. The recurrent aggressive outbursts are not better explained by another mental disorder (e.g., Major Depressive Disorder, Bipolar Disorder, [Posttraumatic Stress Disorder], Disruptive Mood Dysregulation Disorder, a Psychotic Disorder, Antisocial Personality Disorder, Borderline Personality Disorder)...

Note: This diagnosis can be made in addition to the diagnosis of Attention-Deficit/ Hyperactivity Disorder when recurrent impulsive aggressive outbursts are in excess of those usually seen in this disorder and warrant independent clinical attention.




Do you have these outbursts only when you've been drinking, using drugs, or taking medications?

IF UNKNOWN: Have you ever had a head injury, seizure, stroke, or some other kind of neurological illness?

IF YES: Have these outbursts occurred
only during (ILLNESS MENTIONED
ABOVE)?

...and are not attributable to another medical condition (e.g., head trauma, Alzheimer's disease) or to the physiological effects of a substance (e.g., alcohol, phencyclidine, cocaine and other stimulants, barbiturates, inhalants, or a medication).

CRITERIA A, B, C, D, AND F ARE CODED "3"



GO TO THE NEXT MODULE

THIS ITEM HAS ALREADY BEEN ASKED AS PART OF THE OVERVIEW AND HAS BEEN PREPOPULATED BASED ON INFORMATION PREVIOULSY OBTAINED:

IF UNKNOWN: How were you affected by the coronavirus pandemic? (Did you or someone close to you need to be hospitalized for treatment? Did you lose someone whom you were close to? How about the financial implications for people close to you related to the crisis?)

QUESTIONS TO DETERMINE POSSIBLE ASSOCIATION OF EACH PAST 12 MONTH DIAGNOSIS WITH **CORONAVIUS AND ASSOCIATED STRESSORS:** (FILL OUT THIS PAGE SEPARATELY FOR EACH PAST 12 MONTH DIAGNOSIS) FOR EACH DISORDER DIAGNOSED IN PAST 12 **MONTHS:** IF UNKNOWN: When did [SXS OF DIAGNOSED DISORDER] start? → IF ONSET SINCE START OF CORONAVIRUS PANDEMIC IN JANUARY 2020: IF UNKNOWN: What was going on in your life when (SXS) started? Do you think (SXS) were due to the effects of the coronavirus pandemic on your life? IF ONSET PRIOR TO START OF **CORONAVIRUS PANDEMIC IN JANUARY** 2020: Did (SXS) become worse since the start of the pandemic? IF YES: When did they get worse? How much worse? Do you think they got worse because of the effects of the coronavirus pandemic on your life? BASED ON ALL AVAILABLE INFORMATION, INDICATE FOR EACH 12-MONTH DIAGNOSIS THE LIKELIHOOD THAT DISORDER OCCURRING IN PAST 12 MONTHS WAS DUE TO THE EFFECTS OF CORONAVIRUS PANDEMIC: (INCLUDING ECONOMIC EFFECTS) 2 10 1 7 Not at all likely Somewhat likely Very likely