

Hemovigilance Module Adverse Reaction Transfusion Associated Circulatory Overload

*Required for saving

*Facility ID#: _____ NHSN Adverse Reaction #: _____

Patient Information

*Patient ID: _____ *Gender: ☐ M ☐ F ☐ Other *Date of Birth: ____/____/____Sex at Birth: ☐ M ☐ F ☐ Unknown

Gender Identity (Specify) _____

Social Security #: _____

Secondary ID: _____

Medicare #: _____

Last Name: _____

First Name: _____

Middle Name: _____

Ethnicity ☐ Hispanic or Latino ☐ Not Hispanic or Not LatinoRace ☐ American Indian/Alaska Native ☐ Asian ☐ Black or African American☐ Native Hawaiian/Other Pacific Islander ☐ White*Blood Group: ☐ A- ☐ A+ ☐ B- ☐ B+ ☐ AB- ☐ AB+ ☐ O- ☐ O+ ☐ Blood type not done☐ Transitional ABO / Rh + ☐ Transitional ABO / Rh - ☐ Transitional ABO / Transitional Rh☐ Group A/Transitional Rh ☐ Group B/Transitional Rh ☐ Group O/Transitional Rh ☐ Group AB/Transitional Rh

Patient Medical History

List the patient's admitting diagnosis. (Use ICD-10 Diagnostic codes/descriptions)

Code: _____ Description: _____

Code: _____ Description: _____

Code: _____ Description: _____

List the patient's underlying indication for transfusion. (Use ICD-10 Diagnostic codes/descriptions)

Code: _____ Description: _____

Code: _____ Description: _____

Code: _____ Description: _____

List the patient's comorbid conditions at the time of the transfusion related to the adverse reaction. (Use ICD-10 Diagnostic codes/descriptions)

☐ UNKNOWN☐ NONE

Code: _____ Description: _____

Code: _____ Description: _____

Code: _____ Description: _____

Assurance of Confidentiality: The voluntarily provided information obtained in this surveillance system that would permit identification of any individual or institution is collected with a guarantee that it will be held in strict confidence, will be used only for the purposes stated, and will not otherwise be disclosed or released without the consent of the individual, or the institution in accordance with Sections 304, 306 and 308(d) of the Public Health Service Act (42 USC 242b, 242k, and 242m(d)).

Public reporting burden of this collection of information is estimated to average 21 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering, and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC, Reports Clearance Officer, 1600 Clifton Rd., MS H21-8, Atlanta, GA 30333, ATTN: PRA (0920-0666).

List the patient's relevant medical procedure including past procedures and procedures to be performed during the current hospital or outpatient stay. (Use ICD-10 Procedure codes/descriptions) ☐ UNKNOWN ☐ NONE

Code: _____ Description: _____
Code: _____ Description: _____
Code: _____ Description: _____

Additional Information _____

Transfusion History

Has the patient received a previous transfusion? ☐ YES ☐ NO ☐ UNKNOWN
Blood Product: ☐ WB ☐ RBC ☐ Platelet ☐ Plasma ☐ Cryoprecipitate ☐ Granulocyte
Date of Transfusion: ____/____/____ ☐ UNKNOWN
Was the patient's adverse reaction transfusion-related? ☐ YES ☐ NO
If yes, provide information about the transfusion adverse reaction.
Type of transfusion adverse reaction: ☐ Allergic ☐ AHTR ☐ DHTR ☐ DSTR ☐ FNHTR
☐ HTR ☐ TTI ☐ PTP ☐ TACO ☐ TAD ☐ TA-GVHD ☐ TRALI ☐ UNKNOWN
☐ OTHER Specify _____

Reaction Details

*Date reaction occurred: ____/____/____ *Time reaction occurred: ____:____ ☐ Time unknown
*Facility location where patient was transfused: _____
Is this reaction associated with an incident? ☐ Yes ☐ No If Yes, Incident #: _____

Investigation Results

*☐ **Transfusion associated circulatory overload (TACO)**

*Case Definition

Check all that occurred within 12 hours of cessation of transfusion (new onset or exacerbation):

- ☐ Acute respiratory distress (dyspnea, orthopnea, cough)
- ☐ Elevated brain natriuretic peptide (BNP)
- ☐ Elevated central venous pressure (CVP)
- ☐ Evidence of left heart failure
- ☐ Evidence of positive fluid balance
- ☐ Radiographic evidence of pulmonary edema

Other signs and symptoms: (check all that apply)

Generalized:	☐ Chills/rigors	☐ Fever	☐ Nausea/vomiting
Cardiovascular:	☐ Blood pressure decrease	☐ Shock	
Cutaneous:	☐ Edema	☐ Flushing	☐ Jaundice
	☐ Other rash	☐ Pruritus (itching)	☐ Urticaria (hives)
Hemolysis/Hemorrhage:	☐ Disseminated intravascular coagulation	☐ Hemoglobinemia	
	☐ Positive antibody screen		
Pain:	☐ Abdominal pain	☐ Back pain	☐ Flank pain
			☐ Infusion site pain
Renal:	☐ Hematuria	☐ Hemoglobinuria	☐ Oliguria
Respiratory:	☐ Bilateral infiltrates on chest x-ray	☐ Bronchospasm	☐ Cough
	☐ Hypoxemia	☐ Shortness of breath	

☐ Other: (specify) _____

***Severity**

Did the patient receive or experience any of the following?

- | | |
|---|---|
| ☐ No treatment required | ☐ Symptomatic treatment only |
| ☐ Hospitalization, including prolonged hospitalization | ☐ Life-threatening reaction |
| ☐ Disability and/or incapacitation | ☐ Congenital anomaly or birth defect(s) of the fetus |
| ☐ Other medically important conditions | ☐ Death ☐ Unknown or not stated |

***Imputability**

Which best describes the relationship between the transfusion and the reaction?

- ☐ No other explanations for circulatory overload are possible.
☐ Transfusion is a likely contributor to circulatory overload
☐ The patient has a history of a pre-existing condition(s) that most likely explains circulatory overload.
☐ Evidence is clearly in favor of a cause other than the transfusion, but transfusion cannot be excluded.
☐ There is conclusive evidence beyond reasonable doubt of a cause other than the transfusion.
☐ The relationship between the adverse reaction and the transfusion is unknown or not stated.

Did the transfusion occur at your facility? ☐ YES ☐ NO

Does the patient have a history of cardiac insufficiency?

- ☐ Yes, the patient has a history of cardiac insufficiency that could explain the circulatory overload, but transfusion is just as likely to have caused the circulatory overload.
☐ Yes, the patient has a history of pre-existing cardiac insufficiency that most likely explains circulatory overload.
☐ No, the patient does not have a history of cardiac insufficiency.

Did the patient received other fluids in addition to the transfusion? ☐ YES ☐ NO

Module-generated Designations

NOTE: Designations for case definition, severity, and imputability will be automatically assigned in the NHSN application based on responses in the corresponding investigation results section above.

***Do you agree with the case definition designation?** ☐ YES ☐ NO

^Please indicate your designation _____

***Do you agree with the severity designation?** ☐ YES ☐ NO

^Please indicate your designation _____

***Do you agree with the imputability designation?** ☐ YES ☐ NO

^Please indicate your designation _____

Patient Treatment

Did the patient receive treatment for the transfusion reaction? ☐ YES ☐ NO ☐ UNKNOWN

If yes, select treatment(s):

- ☐ Medication (*Select the type of medication*)
- | | | | | |
|---|---|---|---|------------------------------------|
| ☐ Antipyretics | ☐ Antihistamines | ☐ Inotropes/Vasopressors | ☐ Bronchodilator | ☐ Diuretics |
| ☐ Intravenous Immunoglobulin | ☐ Intravenous steroids | ☐ Corticosteroids | ☐ Antibiotics | |
| ☐ Antithymocyte globulin | ☐ Cyclosporin | ☐ Other | | |

Page 4 of 4