

The table below shows the expected coefficients of variation (CV) for the estimates for adults by NCHS for presentation criteria. Current NCHS presentation standards\*\* moved away from C criteria. The current standards generally allow for lower prevalence estimates to be released.

CV were calculated assuming a design effect of 2.5 CV in RED are estimates that would not have

Table 1. Sample Adults: Coefficients of Variation for estimation variables with varying expected

<b>SUBGROUP Prevalence:</b>		All Adults (100%)	90%	80%	70%
<b>SUBGROUP Sample Size:</b>		30,000	27,000	24,000	21,000
<b>Prevalence of estimation variable:</b>	Design Effect:	2.5	2.5	2.5	2.5
Variables with prevalence .5%	0.5%	12.88	13.57	14.40	15.39
Variables with prevalence 1%	1%	9.08	9.57	10.16	10.86
Variables with prevalence 2%	2%	6.39	6.74	7.14	7.64
Variables with prevalence 3%	3%	5.19	5.47	5.80	6.20
Variables with prevalence 4%	4%	4.47	4.71	5.00	5.35
Variables with prevalence 5%	5%	3.98	4.19	4.45	4.76
Variables with prevalence 6%	6%	3.61	3.81	4.04	4.32
Variables with prevalence 7%	7%	3.33	3.51	3.72	3.98
Variables with prevalence 8%	8%	3.10	3.26	3.46	3.70
Variables with prevalence 9%	9%	2.90	3.06	3.25	3.47
Variables with prevalence 10%	10%	2.74	2.89	3.06	3.27
Variables with prevalence 20%	15%	2.17	2.29	2.43	2.60
Variables with prevalence 15%	20%	1.83	1.92	2.04	2.18
Variables with prevalence 25%	25%	1.58	1.67	1.77	1.89
Variables with prevalence 30%	30%	1.39	1.47	1.56	1.67
Variables with prevalence 40%	40%	1.12	1.18	1.25	1.34
Variables with prevalence 45%	45%	1.01	1.06	1.13	1.21

and childrens and for smaller samples for subgroups. CV of 30% has been used as a threshold CV and are based on exact confidence intervals but were established to be consistent with prior

met the CV criteria for presentation.

d prevalence by expected sample sizes of covariates

	60%	50%	40%	30%	20%	15%	10%	5%	4%	3%	2%
	18,000	15,000	12,000	9,000	6,000	4,500	3,000	1,500	1,200	900	600
	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5
	16.62	18.21	20.36	23.51	28.80	33.25	40.72	57.59	64.39	74.35	91.06
	11.73	12.85	14.36	16.58	20.31	23.45	28.72	40.62	45.41	52.44	64.23
	8.25	9.04	10.10	11.67	14.29	16.50	20.21	28.58	31.95	36.89	45.18
	6.70	7.34	8.21	9.48	11.61	13.40	16.41	23.21	25.95	29.97	36.70
	5.77	6.32	7.07	8.16	10.00	11.55	14.14	20.00	22.36	25.82	31.62
	5.14	5.63	6.29	7.26	8.90	10.27	12.58	17.80	19.90	22.97	28.14
	4.66	5.11	5.71	6.60	8.08	9.33	11.43	16.16	18.07	20.86	25.55
	4.30	4.71	5.26	6.07	7.44	8.59	10.52	14.88	16.64	19.21	23.53
	4.00	4.38	4.89	5.65	6.92	7.99	9.79	13.84	15.48	17.87	21.89
	3.75	4.11	4.59	5.30	6.49	7.49	9.18	12.98	14.51	16.76	20.53
	3.54	3.87	4.33	5.00	6.12	7.07	8.66	12.25	13.69	15.81	19.36
	2.81	3.07	3.44	3.97	4.86	5.61	6.87	9.72	10.87	12.55	15.37
	2.36	2.58	2.89	3.33	4.08	4.71	5.77	8.16	9.13	10.54	12.91
	2.04	2.24	2.50	2.89	3.54	4.08	5.00	7.07	7.91	9.13	11.18
	1.80	1.97	2.20	2.55	3.12	3.60	4.41	6.24	6.97	8.05	9.86
	1.44	1.58	1.77	2.04	2.50	2.89	3.54	5.00	5.59	6.45	7.91
	1.30	1.43	1.60	1.84	2.26	2.61	3.19	4.51	5.05	5.83	7.14

1%

300

2.5

128.78

90.83

63.90

51.91

44.72

39.79

36.13

33.27

30.96

29.03

27.39

21.73

18.26

15.81

13.94

11.18

10.09

prevalence of modality

Chiropractor	10%
Acupuncture	6%
Meditation	14%
Massage	6%
Naturopathy	6%
Guided imagery or progressive relaxation	14%
Yoga	14%

subgroup % at minimum prevalence (.14)	required subgroup size
0.7%	7%
0.7%	12%
0.7%	5%
0.7%	12%
0.7%	12%
0.7%	5%
0.7%	5%

The table below shows the expected coefficients of variation (CV) for the estimates for adults and children. Current NCHS presentation standards\*\* moved away from CV and are based on exact confidence interval estimates to be released.

CV were calculated assuming a design effect of 2.5 CV in RED are estimates that would not have met the

Table 1. Sample Children: Coefficients of Variation for estimation variables with varying expected prevalence

<b>SUBGROUP Prevalence:</b>		100%	90%	80%	70%
<b>SUBGROUP Sample Size:</b>		10,000	9,000	8,000	7,000
<b>Prevalence of estimation variable:</b>	<b>Design Effect:</b>	2.5	2.5	2.5	2.5
Variables with prevalence 2%	2%	11.07	11.67	12.37	13.23
Variables with prevalence 3%	3%	8.99	9.48	10.05	10.75
Variables with prevalence 5%	5%	6.89	7.26	7.71	8.24
Variables with prevalence 7%	7%	5.76	6.07	6.44	6.89
Variables with prevalence 10%	10%	4.74	5.00	5.30	5.67
Variables with prevalence 20%	20%	3.16	3.33	3.54	3.78
Variables with prevalence 15%	15%	3.76	3.97	4.21	4.50
Variables with prevalence 25%	25%	2.74	2.89	3.06	3.27
Variables with prevalence 30%	30%	2.42	2.55	2.70	2.89
Variables with prevalence 40%	40%	1.94	2.04	2.17	2.31
Variables with prevalence 45%	45%	1.75	1.84	1.95	2.09

ens and for smaller samples for subgroups. CV of 30% has been used as a threshold by NCHS for present  
 als but were established to be consistent with prior criteria. The current standards generally allow for low

CV criteria for presentation.

ence by expected sample sizes of covariates

	60%	50%	40%	30%	20%	15%	10%	5%	2%
	6,000	5,000	4,000	3,000	2,000	1,500	1,000	500	200
	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5
	14.29	15.65	17.50	20.21	24.75	28.58	35.00	49.50	78.26
	11.61	12.71	14.22	16.41	20.10	23.21	28.43	40.21	63.57
	8.90	9.75	10.90	12.58	15.41	17.80	21.79	30.82	48.73
	7.44	8.15	9.11	10.52	12.89	14.88	18.22	25.77	40.75
	6.12	6.71	7.50	8.66	10.61	12.25	15.00	21.21	33.54
	4.08	4.47	5.00	5.77	7.07	8.16	10.00	14.14	22.36
	4.86	5.32	5.95	6.87	8.42	9.72	11.90	16.83	26.61
	3.54	3.87	4.33	5.00	6.12	7.07	8.66	12.25	19.36
	3.12	3.42	3.82	4.41	5.40	6.24	7.64	10.80	17.08
	2.50	2.74	3.06	3.54	4.33	5.00	6.12	8.66	13.69
	2.26	2.47	2.76	3.19	3.91	4.51	5.53	7.82	12.36

ation criteria.  
ver prevalence



The table below shows the expected coefficients of variation (CV) for the estimates for adults and intervals but were established to be consistent with prior criteria. The current standards generally

CV were calculated assuming a design effect of 2.5 CV in RED are estimates that would not have m

Table 1. Sample Children: Coefficients of Variation for estimation variables with varying expected

<b>SUBGROUP Prevalence:</b>		100%	90%	80%	70%
<b>SUBGROUP Sample Size:</b>		7,650	6,885	6,120	5,355
<b>Prevalence of estimation variable:</b>	<b>Design Effect:</b>	2.5	2.5	2.5	2.5
	1%	17.99	18.96	20.11	21.50
Variables with prevalence 2%	2%	12.65	13.34	14.15	15.12
Variables with prevalence 3%	3%	10.28	10.84	11.49	12.29
	4%	8.86	9.34	9.90	10.59
Variables with prevalence 5%	5%	7.88	8.31	8.81	9.42
	6%	7.16	7.54	8.00	8.55
Variables with prevalence 7%	7%	6.59	6.95	7.37	7.88
	8%	6.13	6.46	6.85	7.33
	9%	5.75	6.06	6.43	6.87
Variables with prevalence 10%	10%	5.42	5.72	6.06	6.48
	11%	5.14	5.42	5.75	6.15
	12%	4.90	5.16	5.47	5.85
	13%	4.68	4.93	5.23	5.59
	14%	4.48	4.72	5.01	5.36
Variables with prevalence 15%	15%	4.30	4.54	4.81	5.14
	16%	4.14	4.37	4.63	4.95
	17%	3.99	4.21	4.47	4.77
	18%	3.86	4.07	4.31	4.61
	19%	3.73	3.93	4.17	4.46
Variables with prevalence 20%	20%	3.62	3.81	4.04	4.32
Variables with prevalence 25%	25%	3.13	3.30	3.50	3.74
Variables with prevalence 30%	30%	2.76	2.91	3.09	3.30
Variables with prevalence 40%	40%	2.21	2.33	2.48	2.65
Variables with prevalence 45%	45%	2.00	2.11	2.23	2.39

childrens and for smaller samples for subgroups. CV of 30% has been used as a threshold by NCHS for p  
 / allow for lower prevalence estimates to be released.

et the CV critieria for presentation.

prevalence by expected sample sizes of covariates

60%	50%	40%	30%	20%	19%	18%	17%	16%	15%	10%
4,590	3,825	3,060	2,295	1,530	1,454	1,377	1,301	1,224	1,148	765
2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5
23.22	25.44	28.44	32.84	40.22	41.26	42.40	43.62	44.97	46.44	56.88
16.34	17.90	20.01	23.10	28.30	29.03	29.83	30.69	31.64	32.67	40.02
13.27	14.54	16.25	18.77	22.99	23.58	24.23	24.93	25.70	26.54	32.51
11.43	12.52	14.00	16.17	19.80	20.32	20.87	21.48	22.14	22.87	28.01
10.17	11.14	12.46	14.39	17.62	18.08	18.57	19.11	19.70	20.35	24.92
9.24	10.12	11.31	13.06	16.00	16.42	16.87	17.35	17.89	18.47	22.63
8.51	9.32	10.42	12.03	14.73	15.12	15.53	15.98	16.47	17.01	20.84
7.91	8.67	9.69	11.19	13.71	14.06	14.45	14.87	15.33	15.83	19.39
7.42	8.13	9.09	10.49	12.85	13.19	13.55	13.94	14.37	14.84	18.18
7.00	7.67	8.57	9.90	12.13	12.44	12.78	13.15	13.56	14.00	17.15
6.64	7.27	8.13	9.39	11.50	11.80	12.12	12.47	12.86	13.28	16.26
6.32	6.92	7.74	8.94	10.95	11.23	11.54	11.87	12.24	12.64	15.48
6.04	6.61	7.39	8.54	10.46	10.73	11.02	11.34	11.69	12.07	14.79
5.78	6.34	7.08	8.18	10.02	10.28	10.56	10.87	11.20	11.57	14.17
5.56	6.09	6.80	7.86	9.62	9.87	10.14	10.44	10.76	11.11	13.61
5.35	5.86	6.55	7.56	9.26	9.50	9.76	10.05	10.36	10.69	13.10
5.16	5.65	6.32	7.29	8.93	9.16	9.41	9.69	9.99	10.31	12.63
4.98	5.46	6.10	7.04	8.63	8.85	9.09	9.36	9.65	9.96	12.20
4.82	5.28	5.90	6.81	8.35	8.56	8.80	9.05	9.33	9.64	11.80
4.67	5.11	5.72	6.60	8.08	8.29	8.52	8.77	9.04	9.34	11.43
4.04	4.43	4.95	5.72	7.00	7.18	7.38	7.59	7.83	8.08	9.90
3.56	3.91	4.37	5.04	6.17	6.34	6.51	6.70	6.90	7.13	8.73
2.86	3.13	3.50	4.04	4.95	5.08	5.22	5.37	5.54	5.72	7.00
2.58	2.83	3.16	3.65	4.47	4.58	4.71	4.85	5.00	5.16	6.32

resentation criteria. Current NCHS presentation standards\*\* moved away from CV and are based on e

9%	8%	7%	6%	5%	4%	3%	2%	1%	0.9%	0.8%
689	612	536	459	383	306	230	153	77	69	61
2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5
59.96	63.59	67.98	73.43	80.44	89.93	103.85	127.19	179.87	189.60	201.10
42.18	44.74	47.83	51.66	56.59	63.27	73.06	89.48	126.54	133.39	141.48
34.26	36.34	38.85	41.97	45.97	51.40	59.35	72.69	102.79	108.35	114.93
29.52	31.31	33.47	36.16	39.61	44.28	51.13	62.62	88.56	93.35	99.01
26.27	27.86	29.78	32.17	35.24	39.40	45.49	55.72	78.80	83.06	88.10
23.85	25.30	27.04	29.21	32.00	35.78	41.31	50.60	71.55	75.42	80.00
21.96	23.30	24.90	26.90	29.47	32.95	38.04	46.59	65.89	69.46	73.67
20.43	21.67	23.17	25.03	27.42	30.65	35.39	43.35	61.30	64.62	68.54
19.16	20.32	21.73	23.47	25.71	28.74	33.19	40.65	57.48	60.59	64.27
18.08	19.17	20.50	22.14	24.25	27.12	31.31	38.35	54.23	57.17	60.63
17.14	18.18	19.44	20.99	23.00	25.71	29.69	36.36	51.42	54.20	57.49
16.32	17.31	18.50	19.99	21.89	24.48	28.26	34.62	48.95	51.60	54.73
15.59	16.53	17.68	19.09	20.91	23.38	27.00	33.07	46.77	49.30	52.29
14.93	15.84	16.93	18.29	20.04	22.40	25.87	31.68	44.80	47.23	50.09
14.34	15.21	16.27	17.57	19.25	21.52	24.85	30.43	43.03	45.36	48.11
13.81	14.64	15.66	16.91	18.52	20.71	23.91	29.29	41.42	43.66	46.31
13.31	14.12	15.10	16.31	17.86	19.97	23.06	28.24	39.94	42.10	44.66
12.86	13.64	14.58	15.75	17.26	19.29	22.28	27.28	38.58	40.67	43.14
12.44	13.20	14.11	15.24	16.69	18.66	21.55	26.39	37.33	39.34	41.73
12.05	12.78	13.67	14.76	16.17	18.08	20.87	25.57	36.16	38.11	40.42
10.44	11.07	11.83	12.78	14.00	15.66	18.08	22.14	31.31	33.00	35.01
9.20	9.76	10.44	11.27	12.35	13.81	15.94	19.53	27.61	29.11	30.87
7.38	7.83	8.37	9.04	9.90	11.07	12.78	15.66	22.14	23.34	24.75
6.66	7.07	7.55	8.16	8.94	9.99	11.54	14.13	19.99	21.07	22.34

xact confidence

0.7%	0.6%	0.5%
54	46	38

2.5	2.5	2.5
-----	-----	-----

214.98	232.21	254.37
151.25	163.37	178.96
122.86	132.71	145.37
105.85	114.33	125.24
94.18	101.73	111.44
85.52	92.37	101.19
78.76	85.07	93.19
73.27	79.14	86.70
68.71	74.21	81.29
64.82	70.01	76.70
61.46	66.38	72.72
58.51	63.20	69.23
55.90	60.37	66.14
53.55	57.84	63.36
51.43	55.56	60.86
49.51	53.47	58.58
47.74	51.57	56.49
46.12	49.81	54.57
44.61	48.19	52.79
43.21	46.68	51.13
37.42	40.42	44.28
33.00	35.65	39.05
26.46	28.58	31.31
23.89	25.80	28.26

The table below shows the expected coefficients of variation (CV) for the estimates for adults and children presentation criteria. Current NCHS presentation standards\*\* moved away from CV and are based on existing standards generally allow for lower prevalence estimates to be released.

CV were calculated assuming a design effect of 2.5 CV in RED are estimates that would not have met the

Table 1. Sample Adolescents: Coefficients of Variation for estimation variables with varying expected prevalence

<b>SUBGROUP Prevalence:</b>		100%	90%	80%	70%
<b>SUBGROUP Sample Size:</b>		3,600	3,240	2,880	2,520
<b>Prevalence of estimation variable:</b>	<b>Design Effect:</b>	2.5	2.5	2.5	2.5
Variables with prevalence 2%	2%	18.45	19.44	20.62	22.05
Variables with prevalence 3%	3%	14.98	15.80	16.75	17.91
Variables with prevalence 5%	5%	11.49	12.11	12.84	13.73
Variables with prevalence 7%	7%	9.61	10.12	10.74	11.48
Variables with prevalence 10%	10%	7.91	8.33	8.84	9.45
Variables with prevalence 15%	15%	6.27	6.61	7.01	7.50
Variables with prevalence 20%	20%	5.27	5.56	5.89	6.30
Variables with prevalence 25%	25%	4.56	4.81	5.10	5.46
Variables with prevalence 30%	30%	4.03	4.24	4.50	4.81
Variables with prevalence 40%	40%	3.23	3.40	3.61	3.86
Variables with prevalence 45%	45%	2.91	3.07	3.26	3.48

ens and for smaller samples for subgroups. CV of 30% has been used as a threshold by NCHS for  
 act confidence intervals but were established to be consistent with prior criteria. The current

CV criteria for presentation.

ivalence by expected sample sizes of covariates

	60%	50%	40%	30%	20%	15%	10%	5%	2%
	2,160	1,800	1,440	1,080	720	540	360	180	72
	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5
	23.81	26.09	29.17	33.68	41.25	47.63	58.33	82.50	130.44
	19.34	21.19	23.69	27.36	33.51	38.69	47.39	67.01	105.96
	14.83	16.24	18.16	20.97	25.69	29.66	36.32	51.37	81.22
	12.40	13.58	15.19	17.54	21.48	24.80	30.37	42.96	67.92
	10.21	11.18	12.50	14.43	17.68	20.41	25.00	35.36	55.90
	8.10	8.87	9.92	11.45	14.03	16.20	19.84	28.05	44.36
	6.80	7.45	8.33	9.62	11.79	13.61	16.67	23.57	37.27
	5.89	6.45	7.22	8.33	10.21	11.79	14.43	20.41	32.27
	5.20	5.69	6.36	7.35	9.00	10.39	12.73	18.00	28.46
	4.17	4.56	5.10	5.89	7.22	8.33	10.21	14.43	22.82
	3.76	4.12	4.61	5.32	6.51	7.52	9.21	13.03	20.60