

Child 9-11

Timestamp

Form Approved
OMB NO. 0920-24EG
Exp. Date XX/XX/20XX

Public reporting burden of this collection of information is estimated to average 30 minutes, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Reports Clearance Officer; 1600 Clifton Road NE, MS H21-8, Atlanta, Georgia 30333; ATTN: PRA (0920-24EG).

Completing the survey is voluntary. If you are not comfortable answering a question, just leave it blank.

Please complete the following questions about yourself.

How old are you, in years?

If you are not between the ages of 9-11 years, please request an alternative form from the project staff.

Below is a list of sentences that describe how people feel. Read each phrase and decide if it is "Not True or Hardly Ever True" or "Somewhat True or Sometimes True" or "Very True or Often True" for you. Then for each sentence, select the response that seems to describe you for the last 3 months.

When I feel frightened, it is hard to breathe.

- ☐ Not true or hardly ever true
☐ Somewhat true or sometimes true
☐ Very true or often true

I get headaches when I am at school.

- ☐ Not true or hardly ever true
☐ Somewhat true or sometimes true
☐ Very true or often true

I don't like to be with people I don't know well.

- ☐ Not true or hardly ever true
☐ Somewhat true or sometimes true
☐ Very true or often true

I get scared if I sleep away from home.

- ☐ Not true or hardly ever true
☐ Somewhat true or sometimes true
☐ Very true or often true

I worry about other people liking me.

- ☐ Not true or hardly ever true
☐ Somewhat true or sometimes true
☐ Very true or often true

When I get frightened, I feel like passing out.

- ☐ Not true or hardly ever true
☐ Somewhat true or sometimes true
☐ Very true or often true

I am nervous.

- ☐ Not true or hardly ever true
☐ Somewhat true or sometimes true
☐ Very true or often true

I follow my mother or father wherever they go.

- ☐ Not true or hardly ever true
☐ Somewhat true or sometimes true
☐ Very true or often true

People tell me that I look nervous.

- ☐ Not true or hardly ever true
☐ Somewhat true or sometimes true
☐ Very true or often true

I feel nervous with people I don't know well.

- ☐ Not true or hardly ever true
☐ Somewhat true or sometimes true
☐ Very true or often true

I get stomachaches at school.

- ☐ Not true or hardly ever true
☐ Somewhat true or sometimes true
☐ Very true or often true

When I get frightened, I feel like I am going crazy.

- ☐ Not true or hardly ever true
☐ Somewhat true or sometimes true
☐ Very true or often true

I worry about sleeping alone.

- ☐ Not true or hardly ever true
☐ Somewhat true or sometimes true
☐ Very true or often true

I worry about being as good as other kids.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
When I get frightened, I feel like things are not real.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
I have nightmares about something bad happening to my parents.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
I worry about going to school.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
When I get frightened, my heart beats fast.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
I get shaky.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
I have nightmares about something bad happening to me.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
I worry about things working out for me.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
When I get frightened, I sweat a lot.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
I am a worrier.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
I get really frightened for no reason at all.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
I am afraid to be alone in the house.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
It is hard for me to talk with people I don't know well.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
When I get frightened, I feel like I am choking.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true

People tell me that I worry too much.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
I don't like to be away from my family.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
I am afraid of having anxiety (or panic) attacks.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
I worry that something bad might happen to my parents.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
I feel shy with people I don't know well.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
I worry about what is going to happen in the future.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
When I get frightened, I feel like throwing up.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
I worry about how well I do things.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
I am scared to go to school.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
I worry about things that have already happened.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
When I get frightened, I feel dizzy.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
I feel nervous when I am with other children or adults and I have to do something while they watch me (for example: read aloud, speak, play a game, play a sport.)	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
I feel nervous when I am going to parties, dances, or any place where there will be people that I don't know well.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true
I am shy.	☐ Not true or hardly ever true ☐ Somewhat true or sometimes true ☐ Very true or often true

Did anyone help you complete this survey?

- ☐ Yes, someone helped me, but I completed most of the survey on my own.
 - ☐ Yes, someone helped me with all or most of the survey.
 - ☐ No, I completed the survey on my own.
-

You skipped some questions on this page. Please review and complete those questions before going to the next page. If you intentionally skipped the questions, you can go to the next page.

ASQ

Please complete the survey below. Thank you!

Medical Record Number

Timestamp

Please ask the following questions only for those ages 9 and up.

Note to person administering the ASQ: Please provide the following information to the respondent before asking the questions.

This survey asks about mental health and emotional well-being. If you answer that you have had suicidal thoughts or behaviors, or purposely tried to hurt yourself, we may inform your doctor or other clinic staff. This would be to ensure your safety and provide you with support and care.

By completing this survey, you accept and consent to this protocol. If you have concerns or need immediate help, please tell the clinic staff.

1) In the past few weeks, have you wished you were dead?

- ☐ Yes
☐ No
☐ Refused to answer

2) In the past few weeks, have you felt that you or your family would be better off if you were dead?

- ☐ Yes
☐ No
☐ Refused to answer

3) In the past week, have you been having thoughts about killing yourself?

- ☐ Yes
☐ No
☐ Refused to answer

4) Have you ever tried to kill yourself?

- ☐ Yes
☐ No
☐ Refused to answer

4a) How?

4b) When?

The patient answered "No" to questions 1 through 4; therefore, screening is complete, and it is not necessary to ask question #5. No intervention is necessary; however, clinical judgment can always override a negative screen.

Do you want to ask the patient question #5 (Are you having thoughts of killing yourself right now?) or finish the ASQ?

- ☐ Ask question #5
☐ Finish the ASQ

This patient is considered a positive screen. Ask question #5 to assess acuity.

5) Are you having thoughts of killing yourself right now?

- ☐ Yes
☐ No

5b) Please describe:

Patient is acute positive screen (imminent risk identified)

Patient requires a STAT safety/full mental health evaluation. Patient cannot leave until evaluated for safety.

Keep patient in sight. Remove all dangerous objects from room. Alert physician or clinician responsible for patient's care.

Provide resources to all patients: 988 Suicide and Crisis Lifeline, 988 (call, text),
<https://988lifeline.org/>

Patient is non-acute positive screen (potential risk identified).

Patient requires a brief suicide safety assessment to determine if a full mental health evaluation is needed. If a patient (or parent/guardian) refuses the brief assessment, this should be treated as an "against medical advice" (AMA) discharge.

Alert physician or clinician responsible for patient's care.

Provide resources to all patients: 988 Suicide and Crisis Lifeline, 988 (call, text),
<https://988lifeline.org/>

Initials of person (staff/professional) completing ASQ

Optional: Provide any comments on clinical information entered on this form. Please do not use any patient identifiers.

Overview of ASQ - this information is included above, within skip logic, and only included here for reference.

If patient answers "No" to all questions 1 through 4, screening is complete (not necessary to ask question #5). No intervention is necessary (*Note: Clinical judgment can always override a negative screen).

If patient answers "Yes" to any of questions 1 through 4, or refuses to answer, they are considered a positive screen. Ask question #5 to assess acuity.

"Yes" to question #5 = acute positive screen (imminent risk identified)

Patient requires a STAT safety/full mental health evaluation. Patient cannot leave until evaluated for safety.

Keep patient in sight. Remove all dangerous objects from room. Alert physician or clinician responsible for patient's care.

"No" to question #5 (but "Yes" or "Refused" to one of questions 1-4) = non-acute positive screen (potential risk identified)

Patient requires a brief suicide safety assessment to determine if a full mental health evaluation is needed. If a patient (or parent/guardian) refuses the brief assessment, this should be treated as an "against medical advice" (AMA) discharge.

Alert physician or clinician responsible for patient's care.

Provide Resources to All Patients:

- 988 Suicide and Crisis Lifeline, Call or Text 988
- Visit <https://988lifeline.org> to chat

YGTSS

Please complete the survey below. Thank you!

Medical Record Number

Timestamp

This instrument to be completed by trained professional.

Motor Tics

Age of first motor tics, in years

Describe first motor tic:

Was tic onset sudden or gradual?

Age of worst motor tics, in years?

Motor Tic Symptom Checklist

Please select if the patient currently (during the past week) has each tic OR if they ever (but not currently) had the tic. State age of onset (in years) if patient has had this behavior.

The patient has experienced, or others have noticed, involuntary and apparently purposeless bouts of:

eye movements

eye blinking, squinting, a quick turning of the eyes, rolling of the eyes to one side, or opening eyes wide very briefly.

☐ Current
☐ Ever

What was the age of onset of this behavior?

eye gestures such as looking surprised or quizzical, or looking to one side for a brief period of time, as if s/he heard a noise.

☐ Current
☐ Ever

What was the age of onset of this behavior?

nose, mouth, tongue movements, or facial grimacing

nose twitching, biting the tongue, chewing on the lip or licking the lip, lip pouting, teeth baring, or teeth grinding.

☐ Current
☐ Ever

What was the age of onset of this behavior?

broadening the nostrils as if smelling something,
smiling, or other gestures involving the mouth,
holding funny expressions, or sticking out the tongue.

☐ Current
☐ Ever

What was the age of onset of this behavior?

head jerks/movements

☐ Current
☐ Ever

touching the shoulder with the chin or lifting the
chin up.

What was the age of onset of this behavior?

throwing the head back, as if to get hair out of the
eyes.

☐ Current
☐ Ever

What was the age of onset of this behavior?

shoulder jerks/movements

☐ Current
☐ Ever

jerking a shoulder.

What was the age of onset of this behavior?

shrugging the shoulder as if to say "I don't know."

☐ Current
☐ Ever

What was the age of onset of this behavior?

arm or hand movements

☐ Current
☐ Ever

quickly flexing the arms or extending them, nail
biting, poking with fingers, or popping knuckles.

What was the age of onset of this behavior?

passing hand through the hair in a combing like
fashion, or touching objects or others, pinching, or
counting with fingers for no purpose, or writing
tics, such as writing over and over the same letter
or word, or pulling back on the pencil while writing.

☐ Current
☐ Ever

What was the age of onset of this behavior?

leg, foot, or toe movements

☐ Current
☐ Ever

kicking, skipping, knee-bending, flexing or extension
of the ankles; shaking, stomping or tapping the foot.

What was the age of onset of this behavior?

taking a step forward and two steps backward,
squatting, or deep knee-bending.

☐ Current
☐ Ever

What was the age of onset of this behavior?

abdominal/trunk/pelvis movements

☐ Current
☐ Ever

tensing the abdomen, tensing the buttocks.

What was the age of onset of this behavior?

other simple motor tics.

☐ Current
☐ Ever

Please write example(s):

What was the age of onset of this behavior?

Other complex motor tics

☐ Current
☐ Ever

touching

What was the age of onset of this behavior?

tapping

☐ Current
☐ Ever

What was the age of onset of this behavior?

picking

☐ Current
☐ Ever

What was the age of onset of this behavior?

evening-up

☐ Current
☐ Ever

What was the age of onset of this behavior?

reckless behaviors

☐ Current
☐ Ever

What was the age of onset of this behavior?

stimulus-dependent tics (a tic which follows, for example, hearing a particular word or phrase, seeing a specific object, smelling a particular odor).

☐ Current
☐ Ever

What was the age of onset of this behavior?

Please write example(s):

rude/obscene gestures; obscene finger/hand gestures.

☐ Current
☐ Ever

What was the age of onset of this behavior?

unusual postures.

☐ Current
☐ Ever

What was the age of onset of this behavior?

bending or gyrating, such as bending over.

☐ Current
☐ Ever

What was the age of onset of this behavior?

rotating or spinning on one foot.

☐ Current
☐ Ever

What was the age of onset of this behavior?

copying the action of another (echopraxia)

☐ Current
☐ Ever

What was the age of onset of this behavior?

sudden tic-like impulsive behaviors.

☐ Current
☐ Ever

What was the age of onset of this behavior?

Please describe this behavior.

tic-like behaviors that could injure/mutilate others.

☐ Current
☐ Ever

What was the age of onset of this behavior?

Please describe this behavior.

self-injurious tic-like behavior(s).

☐ Current
☐ Ever

What was the age of onset of this behavior?

Please describe this behavior.

other involuntary and apparently purposeless motor
tics (that do not fit in any previous categories).

Please describe any other patterns or sequences of
motor tic behaviors:

Phonic (Vocal) Tics

Age of first vocal tics, in years

Describe first vocal tic:

Was tic onset sudden or gradual?

Age of worst vocal tics, in years

Phonic Tic Symptom Checklist

Please select if the patient currently (during the past week) has each tic OR if they ever (but not currently) had the tic. State age of onset (in years) if patient has had this behavior.

The patient has experienced, or others have noticed, involuntary and apparently purposeless bouts of:

coughing.

☐ Current
☐ Ever

What was the age of onset of this behavior?

throat clearing.

☐ Current
☐ Ever

What was the age of onset of this behavior?

sniffing.

☐ Current
☐ Ever

What was the age of onset of this behavior?

whistling.

- ☐ Current
☐ Ever

What was the age of onset of this behavior?

animal or bird noises.

- ☐ Current
☐ Ever

What was the age of onset of this behavior?

other simple phonic tics.

- ☐ Current
☐ Ever

What was the age of onset of this behavior?

Please list:

syllables.

- ☐ Current
☐ Ever

What was the age of onset of this behavior?

Please list:

words.

- ☐ Current
☐ Ever

What was the age of onset of this behavior?

Please list:

rude or obscene words or phrases.

- ☐ Current
☐ Ever

What was the age of onset of this behavior?

Please list:

repeating what someone else said, either sounds, single words or sentences. Perhaps repeating what's said on TV (echolalia).

- ☐ Current
☐ Ever

What was the age of onset of this behavior?

repeating something the patient said over and over again (palilalia).

- ☐ Current
☐ Ever

What was the age of onset of this behavior?

other tic-like speech problems, such as sudden changes in volume or pitch.

- ☐ Current
☐ Ever

What was the age of onset of this behavior?

Please describe:

Describe any other patterns or sequences of phonic tic behaviors:

What was the age of onset of this behavior?

Severity Ratings: Number (Past 7-10 days)

Current Motor Number

Current Phonic Number

Rating Scale

(0) None (no tics)

(1) Single tic

(2) Multiple discrete tics (2-5)

(3) Multiple discrete tics (>5)

(4) Multiple discrete tics plus at least one orchestrated pattern of multiple simultaneous or sequential tics where it is difficult to distinguish discrete tics.

(5) Multiple discrete tics plus several (>2) orchestrated paroxysms of multiple simultaneous or sequential tics where it is difficult to distinguish discrete tics.

Severity Ratings: Frequency (Past 7-10 days)

Current Motor Frequency

Current Phonic Frequency

Rating Scale

(0) None: No evidence of specific tic behaviors.

(1) Rarely: Specific tic behaviors have been present during previous week. These behaviors occur infrequently, often not on a daily basis. If bouts of tics occur, they are brief and uncommon.

(2) Occasionally: Specific tic behaviors are usually present on a daily basis, but there are long tic-free intervals during the day. Bouts of tics may occur on occasion and are not sustained for more than a few minutes at a time.

(3) Frequently: Specific tic behaviors are present on a daily basis. Tic free intervals as long as 3 hours are not uncommon. Bouts of tics occur regularly but may be limited to a single setting.

(4) Almost Always: Specific tic behaviors are present virtually every waking hour of every day, and periods of sustained tic behaviors occur regularly. Bouts of tics are common and are not limited to a single setting.

(5) Always: Specific tic behaviors are present virtually all the time. Tic free intervals are difficult to identify and do not last more than 5 to 10 minutes at most.

Severity Ratings: Intensity (Past 7-10 days)

Current Motor Intensity

Current Phonic Intensity

Rating Scale

(0) Absent: Tics are not present at all

(1) Minimal: Tics are not visible or audible (based solely on patient's private experience) or tics are less forceful than comparable voluntary actions and are typically not noticed because of their intensity.

(2) Mild: Tics are not more forceful than comparable voluntary actions or utterances and are typically not noticed because of their intensity.

(3) Moderate: Tics are more forceful than comparable voluntary actions, but are not outside the range of normal expression for comparable voluntary actions or utterances. They may call attention to the individual because of their forceful character.

(4) Marked: Tics are more forceful than comparable voluntary actions or utterances and typically have an "exaggerated" character. Such tics frequently call attention to the individual because of their forceful and exaggerated character.

(5) Severe: Tics are extremely forceful and exaggerated in expression. These tics call attention to the individual and may result in risk of physical injury (accidental, provoked, or self-inflicted) because of their forceful expression.

Severity Ratings: Complexity (Past 7-10 days)

Current Motor Complexity

Current Phonic Complexity

Rating Scale

(0) None: Tics are not present OR if present, all tics are clearly "simple" (sudden, brief, purposeless) in character.

(1) Borderline: Some tics are not clearly "simple" in character.

(2) Mild: Some tics are clearly "complex" (purposeful in appearance) and mimic brief "automatic" behaviors, such as grooming, syllables, or brief meaningful utterances such as "ah huh", or "hi", that could be readily camouflaged.

(3) Moderate: Some tics are more "complex" (more purposeful and sustained in appearance) and may occur in orchestrated bouts that would be difficult to camouflage, but could be rationalized or "explained" as normal behavior or speech (picking, tapping, saying "you bet" or "honey", brief echolalia).

(4) Marked: Some tics are very "complex" in character and tend to occur in sustained orchestrated bouts that would be difficult to camouflage and could not be easily rationalized as normal behavior or speech because of their duration and/or their unusual, inappropriate, bizarre or obscene character (a lengthy facial contortion, touching genitals, echolalia, speech atypicalities, longer bouts of saying "what do you mean" repeatedly or saying "fu" or "sh").

(5) Severe: Some tics involve lengthy bouts of orchestrated behavior or speech that would be impossible to camouflage or successfully rationalize as normal because of their duration and/or extremely unusual, inappropriate, bizarre or obscene character (lengthy displays or utterances often involving copropraxia, self-abusive behavior, or coprolalia).

Severity Ratings: Interference (Past 7-10 days)

Current Motor Interference

Current Phonic Interference

Rating Scale

(0) None: This means there are no tics present at all.

(1) Minimal: When tics are present, they do not interrupt the flow of behavior or speech.

(2) Mild: When tics are present, they occasionally interrupt the flow of behavior or speech.

(3) Moderate: When tics are present, they frequently interrupt the flow of behavior or speech.

(4) Marked: When tics are present, they frequently interrupt the flow of behavior or speech, and they occasionally disrupt intended action or communication.

(5) Severe: When tics are present, they frequently disrupt intended action or communication.

Severity Ratings: Impairment (Past 7-10 days)

Current Motor Impairment

Current Phonic Impairment

Rating Scale

(0) None

(10) Minimal: Tics associated with subtle difficulties in self-esteem, family life, social acceptance or school/job functioning (infrequent upset or concern about tics vis a vis the future, periodic, slight increase in family tensions because of tics; friend or acquaintances may occasionally notice or comment about tics in an upsetting way.

(20) Mild: Tics associated with minor difficulties in self-esteem, family life, social acceptance, or school/job functioning.

(30) Moderate: Tics associated with some clear problems in self-esteem, family life, social acceptance, or school/job functioning (episodes of dysphoria, periodic distress and upheaval in the family, frequent teasing by peers or episodic social avoidance, periodic interference in school/job performance because of tics).

(40) Marked: Tics associated with major difficulties in self-esteem, family life, social acceptance, or school/job functioning.

(50) Severe: Tics associated with extreme difficulties in self-esteem, family life, social acceptance, or school/job functioning (severe depression with suicidal ideation, disruption of the family [separation/divorce, residential placement], disruption of social ties, severely restricted life because of social stigma and social avoidance, removal from school/job).

Initials of person completing this form.

Optional: Provide any comments on clinical information entered on this form. Please do not use any patient identifiers.

For the person conducting the assessment: How familiar are you with this individual (being assessed with YGTSS)?

- ☐ Not familiar (for example: this was my first encounter with this individual, or previous encounters were very brief)
- ☐ Somewhat familiar (for example: I have interacted with this individual on more than one occasion and for more than just a brief encounter)
- ☐ Very familiar (I have interacted with this individual on several occasions AND am very familiar with their tic symptoms)