

Routine Testing in Prioritized Health Care Settings

Aggregate Data Tables for Diagnose Indicators 1A1.1A, 1A1.1B, and 1A1.1C: Percentage of persons tested for HIV (new and previously positives) in health care facilities identified as a priority for routine opt-out screening.

Contextual Question (provide response in the PS20-2010 EPMP where indicated): Please briefly explain the 1) 2010-funded health care facilities that have been prioritized for routine opt-out screening and 2) how HIV screening was implemented.

The following data table should be completed for each EHE county and each of the 7 EHE states:

Table 1A1.A Routine Opt-Out HIV Testing in Prioritized Health Care Facilities: Enter the name of each prioritized health care facility in column A. Select the facility type from the drop-down options in column B. In columns C through H, enter the requested aggregate data (e.g., number of persons served, number of HIV-positive persons, etc.). Please note that in Column D you will need to enter the total number of persons whose HIV test were directly funded by PS20-2010. You may add additional rows as needed.

Prioritized Health Care Facility Name	Facility Type	Number of Persons Served ¹		Number of HIV-Positive Persons ⁴	Previously Diagnosed HIV-Positive Persons
		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated

¹ Includes persons, regardless of age, who attended at least one medical appointment at the health care facility during the reporting period.

² Includes tests that are supported in any way by PS20-2010-funded resources (e.g., funding, test kits, personnel, training and technical assistance, record enhancements).

³ Includes persons who had a positive or negative test result.

⁴ Includes newly and previously diagnosed infections and those with unknown prior history

⁵ Includes newly identified HIV diagnosis that have been confirmed in surveillance. For persons included in this column, the HIV surveillance system diagnosis is found and 2) there is no indication of a previous diagnosis by either client self-report (if the client was asked) or review of other data

⁶ Includes persons for whom a new diagnosis of HIV is determined by self-report or provider report.

CDC Calculations for Percentages: Percentage of persons tested (Column D%)=(D#/C#)*100; Percentage of HIV-Positive Persons (Column E%)=(E#/D#)*100; Percentage of HIV-Positive Persons (Column F%)=(F#/D#)*100; New Positivity—Confirmed in Surveillance (Column G%)=(G#/D#)*100; New Positivity—Not Confirmed in Surveillance (Column H%)=(H#/D#)*100

ettings

and diagnosed with HIV (new and

criteria used to select the PS20-
ning is conducted in each facility.

health care facility in column A and
ta in cells labelled “DDE” (direct data
l or indirectly funded (e.g., EMR

Newly Diagnosed HIV-positive Persons Confirmed in Surveillance ⁵	Newly Diagnosed HIV-positive Persons Identified by Self-Report ⁶
DDE	DDE
% Calculated	% Calculated
DDE	DDE
% Calculated	% Calculated
DDE	DDE
% Calculated	% Calculated
DDE	DDE
% Calculated	% Calculated
DDE	DDE
% Calculated	% Calculated

e, laboratory support, electronic medical

em is checked and 1) no prior report of HIV
sources (if other data sources were checked).

in E%)=(E#/D#)*100; **Percentage of
y—Self Report (Column H%)=(H#/D#)*100.**

Table 1A1.B. Persons Tested and Positivity in Health

	Total Number of Persons Served ¹	Number o Test
	N	N
Age Group		
<15 Years		
15-19 Years		
20-29 Years		
30-65 Years		
≥ 66 Years		
Unknown		
Gender		
Male		
Female		
Transgender ⁷		
Unknown		
Race and Hispanic Origin⁸		
Hispanic or Latino		
Not Hispanic or Latino	American Indian or Alaska Native alone	
	Asian alone	
	Black or African American alone	
	Native Hawaiian or Pacific Islander alone	

	White alone		
	Two or more races		
	Unknown		
Population Groups			
	MSM/IDU		
	MSM		
	PWID		
	Heterosexual males		
	Heterosexual females		
	Other		
	Unknown		
	Total		

¹ Includes persons who received one or more services at the health

² Includes tests that are supported in any way by PS20-2010-funded enhancements).

³ Includes persons who had a positive or negative test result.

⁴ Includes newly and previously diagnosed infections and those with

⁵ Includes newly identified HIV diagnosis that have been confirmed (1) found and 2) there is no indication of a previous diagnosis by either

⁶ Includes self-report and provider report of new HIV diagnosis.

⁷ Transgender includes all persons whose gender identity or expression is woman, transgender male or transgender man, or other gender not

⁸ Race and ethnicity are to be collected at the local level in accordance with aggregate using these categories. Hispanic or Latino persons can be

CDC Calculations for Percentages: *Percentage of persons tested (Column D%)=(D#/B#) * 100; New Positivity—Confirmed in Surveillance*

Health Care Facilities Conducting Routine Opt-Out HIV Testing

[illegible]

Routine Testing in Prioritized Jails and Corre

Aggregate Data Tables for Diagnose Indicators 1A2.1A, 1A2.1B, 1A2.1C: Percentage of persons incarcerated in are tested for and diagnosed with HIV (new and previous positives).

Contextual Question (provide response in the PS20-2010 EPMP where indicated): Please briefly explain the 1) PS20-2010-funded jails and correctional facilities and 2) how HIV screening is conducted in each facility.

The following data table should be completed for each EHE county and each of the 7 EHE states:

Table 1A2.A Routine Opt-Out HIV Testing in Prioritized Jails and Correctional Facilities: Enter the name of each p column A and select when HIV screening is conducted from the drop-down options in column B. In columns C through G, ent cells labelled “DDE” (direct data entry). Please note that in Column D you will need to enter the total number of persons w indirectly funded (e.g., EMR enhancements, testing staff, etc.) by PS20-2010. You may add additional rows as needed.

Prioritized Jail or Correctional Facility Name	HIV Screening Conducted	Number of Persons Incarcerated ¹	Number of Incarcerated Persons Tested ^{2,3}	Number of HIV-positive Persons ⁴	Previously Diagnosed HIV-positive Persons
		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated

¹ Includes persons detained or incarcerated at the prioritized jail or correctional facility during the reporting period.

² Includes tests that are supported in any way by PS20-2010-funded resources (e.g., funding, test kits, personnel, training and technical a medical record enhancements).

³ Includes persons who received at least one HIV test during their detainment or incarceration. Testing can occur at intake, release, or an HIV testing occurs more than once during detainment or incarceration, then report results from most recent HIV test.

⁴ Includes newly and previously diagnosed infections and those with unknown prior history.

⁵ Includes newly identified HIV diagnosis that have been confirmed in surveillance. For persons included in this column, the HIV surveillar of HIV diagnosis is found and 2) there is no indication of a previous diagnosis by either client self-report (if the client was asked) or review were checked).

⁶ Includes self-report and provider report of new HIV diagnosis.

CDC Calculations for Percentages: *Percentage of persons tested (Column D%)*=(D#/C#)* 100; *Percentage of HIV-Positive Persons (Column F%)*=(F#/D#)* 100; *New Positivity—Confirmed in Surveillance (Column G%)*=(G#/D#)* 100; *New Positivity—Self R*

ectional Facilities

jails and correctional facilities who

criteria used to select the prioritized

prioritized jail or correctional facility in ter the requested aggregate data in hose HIV test were directly funded or	
Newly Diagnosed HIV-positive Persons Confirmed in Surveillance ⁵	Newly Diagnosed HIV-positive Persons Identified by Self-Report ⁶
DDE	DDE
% Calculated	% Calculated
DDE	DDE
% Calculated	% Calculated
DDE	DDE
% Calculated	% Calculated
DDE	DDE
% Calculated	% Calculated
DDE	DDE
% Calculated	% Calculated

ssistance, laboratory support, electronic

y time during detainment or incarceration. If

nce system is checked and 1) no prior report
v of other data sources (if other data sources

in E%)=(E#/D#)*100; Percentage of Previous
eport (Column H%)=(H#/D#)*100.

Table 1A2.B. Persons Tested and Positivity in Ja

		Total Number of Persons Detained or Incarcerated ¹
		N
Age Group		
<15 Years		
15-19 Years		
20-29 Years		
30-65 Years		
≥ 66 Years		
Unknown		
Gender		
Male		
Female		
Transgender ⁷		
Unknown		
Race and Hispanic Origin ⁸		
Hispanic or Latino		
Hispanic or Latino	American Indian or Alaska Native alone	
	Asian alone	
	Black or African American alone	

Not Hispanic or Latino	Native Hawaiian or Pacific Islander alone	
	White alone	
	Two or more races	
	Unknown	
Population Groups		
	MSM/IDU	
	MSM	
	PWID	
	Heterosexual males	
	Heterosexual females	
	Other	
	Unknown	
	Total	

¹ Includes persons detained or incarcerated at the prioritized j

² Includes tests that are supported in any way by PS20-2010-fu enhancements).

³ Includes persons who had a positive or negative test result.

⁴ Includes newly and previously diagnosed infections and thos

⁵ Includes newly identified HIV diagnosis that have been confir found and 2) there is no indication of a previous diagnosis by c

⁶ Includes self-report and provider report of new HIV diagnosi:

⁷ Transgender' includes all persons whose gender identity or e transgender male or transgender man, or other gender non-bi

⁸Race and ethnicity are to be collected at the local le reported to CDC in the aggregate using these catego

CDC Calculations for Percentages: *Percentage of persons test (Column D%)=(D#/B#) * 100; New Positivity—Confirmed in Sur*

Prisons and Correctional Facilities Conducting Routine Opt-Out HIV Testing

[illegible]

[illegible]

Testing in Non-Traditional Venues:

Aggregate Data Tables for Diagnose indicators 1B1.1A, 1B1.1B, and 1B1.1C: Percentage of HIV-positive and previous positives) in non-traditional venues identified as a priority for FHE testing services.

Contextual Question 1B1 (provide response in the PS20-2010 EPMP where indicated): Please list all the non-traditional testing venues where PS20-2010- funded testing is conducted.

The following data table should be completed for each EHE county and each of the 7 EHE states:

Table 1B1.A Testing in Non-Traditional Venues: Enter the name of each non-traditional venue category in setting type (health care or non-healthcare) from the drop-down options in column B. In columns C through F, enter aggregate data in cells labelled "DDE" (direct data entry). You may add additional rows as needed.

Non-Traditional Venue Category ¹	Setting Type	Number of Tests	Number of HIV-positive Tests ²	Previously Diagnosed HIV-positive Persons	Newly Diagnosed HIV-positive Persons Confirmed in Surveillance ³
		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated

¹ Aggregate counts for each non-traditional venue category (e.g., retail, pharmacy, mobile, etc.) should be reported in this column. Recipients conducting HIV testing in 6 different non-traditional pharmacy venues (e.g., CVS, Walgreens, Rite Aid, etc.) and 7 different non-traditional retail venues (e.g., Ross, Marshalls, Dollar General etc.) would report aggregate counts for two non-traditional venue categories—pharmacy and retail. Recipients will not need to report individual aggregate counts for each of the 13 venues noted in the example.

² Includes newly and previously diagnosed infections and those with unknown prior history

³ Includes newly identified HIV diagnosis that have been confirmed in surveillance. For persons included in this column, the system is checked and 1) no prior report of HIV diagnosis is found and 2) there is no indication of a previous diagnosis by either the client (if the client was asked) or review of other data sources (if other data sources were checked).

⁴ Includes self-report and provider report of new HIV diagnosis.

CDC Calculations for Percentages: *Percentage of HIV-Positive Persons (Column D%)*=(D#/C#)* 100; *Percentage of Previously Diagnosed HIV-Positive Persons (Column E%)*=(E#/C#)* 100; *New Positivity—Confirmed in Surveillance (Column F%)*=(F#/C#)* 100; *New Positivity—Self Report (Column G%)*=(G#/C#)* 100

***Drop down options:** healthcare setting, non-healthcare setting.

live tests (new
and briefly describe

column A and select enter the requested
Newly Diagnosed HIV-positive Persons Identified by Self-Report ⁴
DDE
% Calculated
DDE
% Calculated
DDE
% Calculated
DDE
% Calculated
DDE
% Calculated

column. For example, a
different retail venues
agency and retail.

HIV surveillance
other client self-report

HIV-Positives (Column
H%)=(G#/C#)*100.

Table 1B1.B HIV Testing and Positive Tests in Non-Traditional Settings			
	Number of HIV Tests	Number of HIV- positive Tests ¹	
	N	N	%
Age Group			
<15 Years			
15-19 Years			
20-29 Years			
30-65 Years			
≥ 66 Years			
Unknown			
Test Setting			
Health Care Setting			
Non-Health Care Setting			
Unknown			
Testing History			
First-Time Tester			
Repeat Tester			
Unknown			
Gender			
Male			
Female			

Transgender ⁴				
Unknown				
Race and Hispanic Origin⁵				
Hispanic or Latino				
Not Hispanic or Latino	American Indian or Alaska Native alone			
	Asian alone			
	Black or African American alone			
	Native Hawaiian or Pacific Islander alone			
	White alone			
	Two or more races			
Unknown				
Population Groups				
MSM/IDU				
MSM				
PWID				
Heterosexual males				
Heterosexual females				
Other				
Unknown				
Total				

¹ Includes newly and previously diagnosed infections and those with unknown status.

² Includes newly identified HIV diagnosis that have been confirmed in surveillance data sources and 2) there is no indication of a previous HIV diagnosis (data sources were checked).

³ Includes self-report and provider report of new HIV diagnosis.

⁴ Transgender' includes all persons whose gender identity or expression is transgender woman, transgender male or transgender man, or other transgender identity.

⁵ Race and ethnicity are to be collected at the local level in accordance with the OMB standards (https://www.govinfo.gov/content/pkg/FR-1997-10-30/pdf/97-28653.pdf). These categories. Hispanic or Latino persons can be of any race.

CDC Calculations for Percentages: Percentage of HIV-Positive Persons with New Positivity—Confirmed in Surveillance (Column D%)=(D#/A#) * 100. Result (Column F%)=(F#/A#) * 100.

Additional Venues							
Number of Previously Diagnosed HIV-positive Persons		Newly Diagnosed HIV-positive Persons Confirmed in Surveillance ²		Newly Diagnosed HIV-positive Persons Identified by Self-Report ³		Test Result Unknown	
N	%	N	%	N	%	N	%

HIV Self-Test

Aggregate Data Tables for Diagnose indicators 1B2.1 and 1C4.1: Number of HIV self-test kits distributed in health department and number of persons who received at least one HIV self-test kit.

Contextual Question (provide response in the PS20-2010 EPMP where indicated): Please briefly describe how to or ordered by testers. Please indicate if, and how test results are provided to the health department.

The following data table should be completed for each EHE county and each of the 7 EHE states:

Table 1C4. HIV Self-Tests and Positive Tests

		Number of HIV Self-Test Kits Distributed	Number of Persons Receiving ≥ 1 HIV Self-Test Kit		Newly Diagnosed HIV-positive Persons Confirmed in Surveillance ¹		Newly Diagnosed HIV-positive Person Self-Reported
		N	N %		N %		N %
Age Group							
<15 Years							
15-19 Years							
20-29 Years							
30-65 Years							
≥ 66 Years							
Unknown							
Test Setting							
Health Care Setting							
Non-Health Care Setting							
Unknown							
Testing History							
First-Time Tester							
Repeat Tester							
Unknown							
Gender							
Male							
Female							
Transgender ³							
Unknown							
Race and Hispanic Origin⁴							
Hispanic or Latino							
Hispanic or Latino	American Indian or Alaska Native alone						
	Asian alone						
	Black or African American alone						

Not Hisp	Native Hawaiian or Pacific Islander alone						
	White alone						
	Two or more races						
Unknown							
Population Groups							
MSM/IDU							
MSM							
PWID							
Heterosexual males							
Heterosexual females							
Other							
Unknown							
Total							

¹ Includes newly identified HIV diagnosis that have been confirmed in surveillance. For persons included in this column, the HIV surveillance system there is no indication of a previous diagnosis by either client self-report (if the client was asked) or review of other data sources (if other data sources were available).

² Includes self-report and provider report of new HIV diagnosis.

³ Transgender' includes all persons whose gender identity or expression is different from their sex assigned at birth. Transgender persons may be male or transgender man, or other gender non-binary person.

⁴ Race and ethnicity are to be collected at the local level in accordance with OMB standards (<https://www.govinfo.gov/content/pkg/FR-1997-10-01/page/10-01.html>). Hispanic or Latino persons can be of any race.

CDC Calculations for Percentages: Percentage of Persons Receiving at least 1 HIV Self-Test Kit (Column B%)=(B#/A#) * 100; New Positivity—Self Report (Column D%)=(D#/B#) * 100; Percentage of Previous HIV-Positives (Column E%)=(E#/B#) * 100; Percentage of Test Results (Column F%)=(F#/B#) * 100.

lth care settings and non-health care settings and

HIV self-test kits funded by PS20-2010 are distributed

[illegible]

Provision of PrEP Services:

Data should include all PrEP services funded by PS20-2010 regardless of the funding source for the HIV test reported as a cascade and will be calculated as specified in the numerator and denominator for each indicator in Table in the PS20-2010 EPMP).

Aggregate Data Tables for Prevent Indicator 3A1.3: *Number and percentage of HIV-negative persons with PrEP provider*

Note: PrEP cascade is reported as follows:

The following data table should be completed for each EHE county and each of the 7 EHE states:

Table 3A1. PrEP in STD Clinics							
	HIV-Negative Persons ¹	Screened for PrEP Need ^{2,3}		Eligible for PrEP Referral ^{4,5}		Referred to PrEP Provider ^{6,7}	
	N	N	%	N	%	N	%
Age Group							
<15 Years							
15-19 Years							
20-29 Years							
30-65 Years							
≥ 66 Years							
Unknown							
Test Setting							
STD Clinic							
Other Health Care Setting							
Non-Health Care Setting							
Unknown							
Gender							
Male							
Female							
Transgender ¹²							
Unknown							
Race and Hispanic Origin¹³							
Hispanic or Latino							
Not Hispanic or Latino	American Indian or Alaska Native alone						
	Asian alone						
	Black or African American alone						
	Native Hawaiian or Pacific Islander alone						
	White alone						
	Two or more races						
Unknown							

Population Groups							
MSM/IDU							
MSM							
PWID							
Heterosexual males							
Heterosexual females							
Other							
Unknown							
Total							

¹ Includes HIV-negative persons who received HIV testing or PrEP services (e.g., screening, referral, linkage, prescription) that were in at 2010 funds. Denominator for Column B

^{2,3} Screening for PrEP eligibility is the process of conducting an initial assessment regarding a person's eligibility (e.g., behavioral risk screening, prophylaxis (PrEP) and determining whether a more thorough evaluation is warranted. A person is screened if an initial assessment of eligibility is completed. For further discussion on PrEP screening, see Centers for Disease Control and Prevention: US Public Health Service: PrEP: prevention of HIV infection in the United States—2017 Update: a clinical practice guideline. <https://www.cdc.gov/hiv/pdf/risk/prep/cdc-prEP-guideline.pdf> Denominator is Column A (number of HIV-negative persons). All persons reflected in Column B should also be reflected in Column C.

⁴ Eligible for PrEP Referral refers to a person's status with regard to whether or not he or she meets appropriate criteria for using PrEP; not he or she is HIV-negative and at risk for HIV, as defined locally or by CDC guidelines for PrEP.

⁵ Denominator is Column B (number of persons screened for PrEP). Persons reported as eligible for PrEP referral are those who have seen a provider. These individuals should also be reflected in the counts reported for columns A-B.

⁶ Referral to a PrEP provider is a process involving the provision of information on who the providers are, what documents referred per person, how to get to the provider's agency, and what to expect from the referral process.

⁷ Denominator is Column C (number of persons eligible for PrEP). Persons who are referred to a PrEP provider should also be reflected in the counts reported for columns A-C.

⁸ Linkage to PrEP provider is the process through which a person at risk for becoming infected with HIV is helped to access a healthcare provider for evaluation and management for PrEP. Linked to a PrEP provider refers to the outcome of the referral or linkage of a PrEP eligible person to a provider indicated by the person's attendance at the first appointment and verified through reviews of medical records or other data systems or other data systems.

⁹ Denominator is Column D (number of persons referred). Persons who are linked to a PrEP provider should also be reflected in the counts reported for columns A-D.

¹⁰ Prescribed PrEP refers to a person who has been adequately evaluated and received a prescription for PrEP. Prescription information can be obtained from medical records, pharmacy data, or self-report by client.

¹¹ Denominator is Column E (number of persons linked to PrEP). Persons who are prescribed PrEP should also be reflected in the counts reported for columns A-E.

¹² 'Transgender' includes all persons whose gender identity or expression is different from their sex assigned at birth. Transgender persons include transgender female or transgender woman, transgender male or transgender man, or other gender non-binary person.

¹³ Race and ethnicity are to be collected at the local level in accordance with OMB standards (<https://www.govinfo.gov/content/pkg/FR-2016-06-28/pdf/2016-06-28.pdf>) and reported to CDC in the aggregate using these categories. Hispanic or Latino persons can be of any race.

CDC Calculations for Percentages: Percentage of Persons Screened for PrEP (Column B)=(B#/A#) * 100; Percentage of Persons Eligible for PrEP (Column C)=(C#/B#) * 100; Percentage of Persons Referred for PrEP (Column D)=(D#/C#) * 100; Percentage of Persons Linked to a PrEP Provider (Column E)=(E#/D#) * 100; Percentage of Persons Prescribed PrEP (Column F)=(F#/E#) * 100.

est. PrEP data will be
licator (see Prevent

10 are linked to a

[illegible]

ny way supported by PS20-

eening) for pre-exposure
eligibility for PrEP services
posure prophylaxis for the
c-hiv-prep-guidelines-
lumn A.

specifically, whether or

qual or drug risk behaviors.

son should take with them,

n the counts reported for

provider who offers
n to a PrEP provider, as
self-report by the client.

ints reported for columns

may come from clinical

s reported for columns A-E.

ons may self-identify as

-1997-10-30/pdf/97-

Eligible for PrEP (Column
Provider (Column

Linkage to SSP:

Aggregate Data Tables for Diagnose Indicators 1A5.1, 1B4.3: *Percentage of persons who inject drugs (PWID) tested for HIV in an STD clinic or non-health care setting who are linked to a syringe services program (SSP).*

The following data table should be completed for each EHE county and each of the 7 EHE states:

Table 1A5.1A PWID Tested for HIV and Linked to SSP					
	Total Number of PWID Screened for HIV	Referred to SSP ¹		Linked to SSP ²	
	N	N	%	N	%
Age Group					
<15 Years					
15-19 Years					
20-29 Years					
30-65 Years					
≥ 66 Years					
Unknown					
Test Setting					
STD Clinic					
Other Health Care Setting					
Non-health Care Setting					
Gender					
Male					
Female					
Transgender ³					
Unknown					

Race and Hispanic Origin ⁴						
Hispanic or Latino						
Not Hispanic or Latino	American Indian or Alaska Native alone					
	Asian alone					
	Black or African American alone					
	Native Hawaiian or Pacific Islander alone					
	White alone					
	Two or more races					
Unknown						
Population Groups						
MSM						
Heterosexual males						
Heterosexual females						
Other						
Unknown						
Total						

¹ Persons who are linked to an SSP are those who attend an appointment at the SSP where they are seen by a provider. Linkage to an SSP may be determined by provider report, record review, or client self-report.

²Active referrals to SSP may include, but are not limited to, activities such as making appointments for the client, providing transportation, using a case manager or peer navigator to help with access to services, providing the organization to which the client is referred with information collected about the client, etc.

³'Transgender' includes all persons whose gender identity or expression is different from their sex assigned at birth. Transgender persons may self-identify as transgender female or transgender woman, transgender male or transgender man, or other gender non-binary person.

⁴Race and ethnicity are to be collected at the local level in accordance with OMB standards (<https://www.govinfo.gov/content/pkg/FR-1997-10-30/pdf/97-28653.pdf>) and reported to CDC in the aggregate using these categories. Hispanic or Latino persons can be of any race.

CDC Calculations for Percentages: Percentage of PWID Referred to SSP (Column B)=(B#/A#)* 100; Percentage of PWID Linked to SSP (Column C)=(C#/B#)* 100.

Syringe Services Program (SSP) Locations and Ser

Aggregate Data Tables for Prevent indicators 3B1.1: Number of SSPs operating in the jurisdiction

The following data table should be completed for each EHE county and each of the 7 EHE states:

Table 3B1.A SSPs in the jurisdiction				
Total Number of SSPs Operating in the Jurisdiction	Number of Fixed Site SSP Locations		Number of Mobile/Outreach SSP Locations	
N	N	%	N	%
DDE	DDE	Calculated	DDE	Calculated

CDC Calculations for Percentages: *Percentage of Fixed Location SSP (Column B)*=(B#/A#) * 100; *Percentage of Mobile/Outreach Location SSP (Column C)*=(C#/A#) * 100.

Aggregate Data Tables for Treat indicators 3B1.2: Number of encounters that received at least one standard SSP service

Contextual Question (provide response in the PS20-2010 EPMP where indicated): Please list and briefly describe the services provided by each of the SSPs operating in your jurisdiction. If the SSP provides outreach services, please briefly describe the method(s) used to conduct outreach (e.g. mobile vans, delivery services, etc.).

Aggregate Data Collection (DDE= direct data entry):

Table 3B1.B Provision of Standard SSP Services in the jurisdiction					
Total Number of SSP Encounters	SSP Encounters with 1 or more Standard Services Provided ¹				
	Number of SSP Encounters with 1 or more Standard Services Provided ¹	Number of Encounters at Fixed Site SSP Locations		Number of Encounters via Mobile/Outreach SSP Locations	
N	N	N	%	N	%
DDE	DDE	DDE	Calculated	DDE	Calculated

¹ Sterile needles, syringes, injection equipment; condoms; syringe disposal; HIV testing; linkage to HIV care ≤30 days of diagnosis; HCV testing; linkage to HCV care; linkage to a PrEP provider; Naloxone distribution, linkage to medication-assistance treatment; infectious disease prevention, detection, care, and treatment; substance use disorder care and treatment; essential support services.

CDC Calculations for Percentages: *Percentage of Encounters at Fixed Location SSP (Column B)*=(B#/A#) * 100; *Percentage of Encounters at Mobile/Outreach Location SSP (Column C)*=(C#/A#) * 100.

Aggregate Data Tables for Treat indicators 3B3.1

Table 3B3.1 SSPs providing or actively referring to key SSP services

Total Number of SSPs Operating in the Jurisdiction	Provide/Actively Refer to Infectious Disease Prevention, Detection, Care, and Treatment		Provide/Actively Refer to Substance Use Disorder Care and Treatment		Provide/Actively Refer to Essential Services
	N	%	N	%	N
Calculated	DDE	Calculated	DDE	Calculated	DDE

CDC Calculations for Percentages: *Percentage of SSP Providing/Referring to Infectious Disease, Prevention, Detection, Care, and Treatment (Column B)=(B#/A#) * 100; Percentage of SSP Providing/Referring to Substance Use Care and Treatment (Column C)=(C#/A#) * 100; Percentage of SSP Providing/Referring to Essential Services (Column D)=(D#/A#) * 100.*

vice Provision:

atively Refer al Support ices
%
Calculated

ention,
› Substance
al Support

Integrated HIV Screening

Aggregate Data Table for Integrated Screening Indicator J1: Percentage of HIV screenings conducted that are integrated with other screenings (i.e., viral hepatitis/TB/STD).

Contextual Question (provide response in the APR and EOY): Please briefly describe where and how integrated screening funded under PS20-2010 is conducted.

The following data table should be completed for each EHE county and each of the 7 EHE states

(DDE= direct data entry):

Table J1: Percentage of HIV screenings conducted that are integrated with other screenings (i.e., viral hepatitis/TB/STD).				
	Number of HIV Tests	Number of HIV tests integrated with viral hepatitis tests ¹	Number of HIV tests integrated with TB tests ²	Number of HIV tests integrated with STD tests ³
Age Group	N	N %	N %	N %
<15 Years	DDE	DDE	DDE	DDE
		% Calculated	% Calculated	% Calculated
15-19 Years	DDE	DDE	DDE	DDE
		% Calculated	% Calculated	% Calculated
20-29 Years	DDE	DDE	DDE	DDE
		% Calculated	% Calculated	% Calculated
30-65 Years	DDE	DDE	DDE	DDE
		% Calculated	% Calculated	% Calculated
≥ 66 Years	DDE	DDE	DDE	DDE
		% Calculated	% Calculated	% Calculated
Unknown	DDE	DDE	DDE	DDE
		% Calculated	% Calculated	% Calculated
Test Setting				
STD Clinic	DDE	DDE	DDE	DDE

STD Clinic		DDE	% Calculated	% Calculated	% Calculated
Other Health Care Setting		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
Non-health Care Setting		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
Gender					
Male		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
Female		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
Transgender ⁴		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
Unknown		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
Race and Hispanic Origin ⁵					
Hispanic or Latino		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
Not Hispanic or Latino	American Indian or Alaska Native alone	DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
	Asian alone	DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
	Black or African American alone	DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
	Native Hawaiian or Pacific Islander alone	DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
	White alone	DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
Two or more races	DDE	DDE	DDE	DDE	
		% Calculated	% Calculated	% Calculated	
Unknown		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
Population Groups					
MSM/IDU		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
MSM		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
PWID		DDE	DDE	DDE	DDE
			% Calculated	% Calculated	% Calculated
Heterosexual males		DDE	DDE	DDE	DDE

Heterosexual males	DDE	% Calculated	% Calculated	% Calculated
Heterosexual females	DDE	DDE	DDE	DDE
		% Calculated	% Calculated	% Calculated
Other	DDE	DDE	DDE	DDE
		% Calculated	% Calculated	% Calculated
Unknown	DDE	DDE	DDE	DDE
		% Calculated	% Calculated	% Calculated
Total				
		% Calculated	% Calculated	% Calculated

¹ Includes HIV tests conducted along with Hepatitis C tests

² Includes HIV tests conducted along with Tuberculosis tests

³ Includes HIV tests conducted along with tests for other sexually transmitted diseases (syphilis, gonorrhea, and chlamydia)

⁴ Transgender¹ includes all persons whose gender identity or expression is different from their sex assigned at birth. Transgender persons may self-identify as transgender female or transgender woman, transgender male or transgender man, or other gender non-binary person.

⁵Race and ethnicity are to be collected at the local level in accordance with OMB standards (<https://www.govinfo.gov/content/pkg/FR-1997-10-30/pdf/97-28653.pdf>) and reported to CDC in the aggregate using these categories. Hispanic or Latino persons can be of any race.

CDC Calculation: *Percentage of HIV-Viral Hepatitis Integrated Tests (Column C%)*=(C#/B)*100; *HIV-TB Integrated Tests (Column D%)*=(D#/B)*100; *Percentage of HIV-STD Integrated Tests (Column E%)*=(E#/B)*100

TelePrEP

Aggregate Data Table for TelePrEP Indicator J2: Percentage of persons who received TelePrEP services

The following data table should be completed for each EHE county and each of the 7 EHE states

(DDE= direct data entry):

Table J2: Percentage of persons who received TelePrEP services		
	Persons who received PrEP Services ¹	Persons who received TelePrEP Services ²
Age Group	N	N %
<15 Years	DDE	DDE
		% Calculated
15-19 Years	DDE	DDE
		% Calculated
20-29 Years	DDE	DDE
		% Calculated
30-65 Years	DDE	DDE
		% Calculated
≥ 66 Years	DDE	DDE
		% Calculated
Unknown	DDE	DDE
		% Calculated
Test Setting		
STD Clinic	DDE	DDE
		% Calculated

Navi

Aggregate assistance

The follow

(DDE= dire

Table J3: P	
Age Group	
<15 Year	
15-19 Ye	
20-29 Ye	
30-65 Ye	
≥ 66 Yea	
Unknow	
Test Setting	
STD Clini	

STD Clinic		DDE	% Calculated
Other Health Care Setting		DDE	DDE
			% Calculated
Non-Health Care Setting		DDE	DDE
			% Calculated
Gender			
Male		DDE	DDE
			% Calculated
Female		DDE	DDE
			% Calculated
Transgender ³		DDE	DDE
			% Calculated
Unknown		DDE	DDE
			% Calculated
Race and Hispanic Origin⁴			
Hispanic or Latino		DDE	DDE
			% Calculated
Not Hispanic or Latino	American Indian or Alaska Native alone	DDE	DDE
			% Calculated
	Asian alone	DDE	DDE
			% Calculated
	Black or African American alone	DDE	DDE
			% Calculated
	Native Hawaiian or Pacific Islander alone	DDE	DDE
			% Calculated
White alone		DDE	DDE
			% Calculated
Two or more races		DDE	DDE
			% Calculated
Unknown		DDE	DDE
			% Calculated
Population Groups			
MSM/IDU		DDE	DDE
			% Calculated
MSM		DDE	DDE
			% Calculated
PWID		DDE	DDE
			% Calculated
Heterosexual males		DDE	DDE

STD Clinic	
Other Health Care Setting	
Non-health Care Setting	
Gender	
Male	
Female	
Transgender	
Unknown	
Race and Hispanic Origin⁴	
Hispanic or Latino	
Not Hispanic or Latino	
Unknown	
Population Groups	
MSM/IDU	
MSM	
PWID	
Heterosexual males	

Heterosexual males	DDE	% Calculated
Heterosexual females	DDE	DDE
		% Calculated
Other	DDE	DDE
		% Calculated
Unknown	DDE	DDE
		% Calculated
Total		
		% Calculated

Heterosexual	
Heterosexual	
Other	
Unknown	

¹ An outcome of the referral or linkage of a PrEP eligible person to a PrEP provider, indicated by attendance at the first appointment and verified through reviews of medical records or other data systems or self-report by the client.

² An outcome of the referral or linkage of a PrEP eligible person to a PrEP provider, indicated by attendance at the first **telehealth** appointment and verified through reviews of medical records or other data systems or self-report by the client. Denominator is Column A (number of persons who received PrEP services).

¹ An outcome records or oth

³ Transgender¹ includes all persons whose gender identity or expression is different from their sex assigned at birth. Transgender persons may self-identify as transgender female or transgender woman, transgender male or transgender man, or other gender non-binary person.

² An encounter HCV testing; li treatment; su

⁴Race and ethnicity are to be collected at the local level in accordance with OMB standards (<https://www.govinfo.gov/content/pkg/FR-1997-10-30/pdf/97-28653.pdf>) and reported to CDC in the aggregate using these categories. Hispanic or Latino persons can be of any race.

³ Transgende transgender v

CDC Calculation: Percentage of HIV-Viral Hepatitis Integrated Tests (Column C%)
=(C#/B)*100; HIV-TB Integrated Tests (Column D%)=(D#/B)*100; Percentage of HIV-STD Integrated Tests (Column E%)=(E#/B)*100

⁴Race and eth 28653.pdf) an

CDC Calculat Percentage c

Navigator Assistance for HIV/STD Screenings

Data Table for Navigator Assistance Indicator J3: Percentage of persons who were linked to HIV/STD screenings/PrEP/SSP with the assistance of navigators

This data table should be completed for each EHE county and each of the 7 EHE states

(For direct data entry):

Percentage of persons who were linked to HIV/STD screenings/PrEP/SSP with the assistance of navigators				
	Number of persons who received navigator assistance for HIV/STD screenings	Number of persons screened for HIV/STD	Number of persons who received navigator assistance for PrEP	Number of persons who were linked to PrEP Provider ¹
	N	N %	N	N %
Alabama	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
Arkansas	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
California	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
Colorado	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
Connecticut	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
Delaware	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
Florida	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
Grand Total				
		DDE		DDE
Notes:	DDE		DDE	

ic	DDE	% Calculated	DDE	% Calculated
ealth Care Setting	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
ith Care Setting	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
nder ³	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
n	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
ispanic Origin ⁴				
or Latino	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
American Indian or Alaska Native alone	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
Asian alone	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
Black or African American alone	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
Native Hawaiian or Pacific Islander alone	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
White alone	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
Two or more races	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
n	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
Groups				
U	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
	DDE	DDE	DDE	DDE
		% Calculated		% Calculated
exual males	DDE	DDE	DDE	DDE

Sexual males		DDE	% Calculated		DDE	% Calculated
Sexual females		DDE	DDE		DDE	DDE
			% Calculated			% Calculated
		DDE	DDE		DDE	DDE
			% Calculated			% Calculated
n		DDE	DDE		DDE	DDE
			% Calculated			% Calculated
Total						
			% Calculated			% Calculated

of the referral or linkage of a PrEP eligible person by navigators to a PrEP provider, indicated by attendance at the first appointment, data from provider data systems or self-report by the client.

or at an SSP to receive SSP services including Sterile needles, syringes, injection equipment; condoms; syringe disposal; HIV testing; linkage to HCV care; linkage to a PrEP provider; Naloxone distribution, linkage to medication-assistance treatment; infectious disease treatment; substance use disorder care and treatment; essential support services

er' includes all persons whose gender identity or expression is different from their sex assigned at birth. Transgender persons include transgender woman, transgender male or transgender man, or other gender non-binary person.

Demographic data are to be collected at the local level in accordance with OMB standards (<https://www.govinfo.gov/content/pkg/FR-1997-05-01/page/10000.pdf>) and reported to CDC in the aggregate using these categories. Hispanic or Latino persons can be of any race.

Formula: Percentage of Persons who were screened for HIV/STD (Column B%) = (B#/A)*100; Percentage of persons who were linked to SSP services (Column F%)=Column F#/E)*100

'/STD screenings/PrEP/SSP with the

Aggregate Data Table for Social Marketing Indicator .
HIV/STD screenings/PrEP/SSP (e.g., print; digital/intel
MSM, youth, MSM of color, Hispanics/Latinos, Blacks/

Contextual Question J4 (provide response in the APR was conducted

The following data table should be completed for each

(DDE= direct data entry):

navigators	
Number of persons who received navigator assistance for SSP services	Number of persons who were linked to SSP services ²
N	N %
DDE	DDE
	% Calculated
DDE	DDE
	% Calculated
DDE	DDE
	% Calculated
DDE	DDE
	% Calculated
DDE	DDE
	% Calculated
DDE	DDE
	% Calculated
DDE	DDE
	% Calculated
DDE	DDE

Table J4: Percentage of impressions from social marketing ca

Social Marketing Campaign Name	Advertising Type ²	Priority Group ³

¹ Out-of-home (OOH) advertising is also known as outdoor billboards, transit ads on buses or trains, wallscapes, and point-of-purchase advertising, such as those in medical centers, airports, stores, and restaurants.

² Includes print advertising; digital/internet-based advertising

³ Includes MSM; youth; MSM of color; Hispanics/Latinos; BI

⁴ In this column report the total number of impressions for reported in this column may reflect repeated exposures to

	DDE	% Calculated
	DDE	DDE
	DDE	% Calculated
	DDE	DDE
	DDE	% Calculated
	DDE	DDE
	DDE	% Calculated
	DDE	DDE
	DDE	% Calculated
	DDE	DDE
	DDE	% Calculated
	DDE	DDE
	DDE	% Calculated
	DDE	DDE
	DDE	% Calculated
	DDE	DDE
	DDE	% Calculated
	DDE	DDE
	DDE	% Calculated
	DDE	DDE
	DDE	% Calculated
	DDE	DDE
	DDE	% Calculated
	DDE	DDE
	DDE	% Calculated
	DDE	DDE
	DDE	% Calculated
	DDE	DDE
	DDE	% Calculated
	DDE	DDE
		% Calculated
	DDE	DDE
		% Calculated
	DDE	DDE

Calculation: Overall impressions specific to population gro
 =(H#/F) * 100.

	DDE	% Calculated
	DDE	DDE
	DDE	% Calculated
	DDE	DDE
	DDE	% Calculated
	DDE	DDE
	DDE	% Calculated
		% Calculated

ntment and verified through reviews of medical

sting; linkage to HIV care ≤30 days of diagnosis;
isease prevention, detection, care, and

may self-identify as transgender female or

7-10-30/pdf/97-

*vere linked to PrEP (Column D%)=(D#/C)*100;*

Campaigns

14: Percentage of impressions from social marketing campaign(s) advertising conducted in (internet-based; radio; television; out-of-home advertising) targeting specified priority groups (e.g., African Americans, and Transgender persons)

15 and EOY): Please list and briefly describe the PS20-2010-funded social marketing campaign that

for each EHE county and each of the 7 EHE states:

Campaign(s) advertising conducted in HIV/STD screenings/PrEP/SSP				
Number of overall impressions ⁴	Number of unique impressions	Number of persons in the priority group	Number of overall impressions specific to priority group	Number of unique impressions specific to priority group
N	N	N	N %	N %
DDE	DDE	DDE	DDE	DDE
			% Calculated	% Calculated
DDE	DDE	DDE	DDE	DDE
			% Calculated	% Calculated
DDE	DDE	DDE	DDE	DDE
			% Calculated	% Calculated
DDE	DDE	DDE	DDE	DDE
			% Calculated	% Calculated
DDE	DDE	DDE	DDE	DDE
			% Calculated	% Calculated

advertising or outdoor media and is generally advertising experienced outside of the home. These include posters that may be seen while “on the go” or in the community. This may also include “place-based” messages, or other buildings or facilities.

Internet-based; radio advertising; television; out-of-home

African Americans; Transgender persons; other group(s) (specified by recipient).

advertising activities targeted to the specific demographic group of interest in the campaign. Counts include direct messaging/advertising.

Imp (Column G%) = $(G\#/F) * 100$; *Unique impressions specific to population group (Column H%)*

Extragenital Testing in STD Clinics

Aggregate Data Tables for Performance Measure: STD-1: *Percentage of persons tested for recta*

Gray = auto-calculated cell; does not represent a reporting burden

TableXXX. Extragenital testing and testing in STD Clinics

		Number of persons tested for rectal GC/CT		Number of persons testing positive for rectal GC/CT	
		N	%	N	%
Age Group					
<15 Years					
15-19 Years					
20-29 Years					
30-65 Years					
≥ 66 Years					
Unknown					
Gender					
Male					
Female					
Transgender					
Unknown					
Race and Hispanic Origin¹					
Hispanic or Latino					
Not Hispanic or Latino	American Indian or Alaska Native alone				
	Asian alone				
	Black or African American alone				
	Native Hawaiian or Pacific Islander alone				
	White alone				
	Two or more races				
Unknown					
Population Groups					
MSM/IDU					
MSM					
PWID					
Heterosexual males					
Heterosexual females					
Other					
Unknown					
Total					

¹Race and ethnicity are to be collected at the local level in accordance with OMB standards (<https://www.govinfo.gov/content/pkg/FR-1997-10-30/pdf/97-28653.pdf>) and reported to CDC in the aggregate using these categories. Hispanic or Latino persons can be of any race.

I GC/CT who are positive for rectal GC/CT

Routine Testing in Prioritized Health Care Settings: STI

Aggregate Data Tables for Diagnose Indicators 1A1.1A, 1A1.1B, and 1A1.1C: Percentage of persc

Gray = auto-calculated cell; does not represent a reporting burden

Table 1A1.B. Persons Tested and Positivity in STD Clinics						
	Total Number of Persons Served ¹	Number of Persons Tested ^{2,3}		Number of HIV- Positive Persons ⁴		Previously Pers
	N	N	%	N	%	N
Age Group						
<15 Years						
15-19 Years						
20-29 Years						
30-65 Years						
≥ 66 Years						
Unknown						
Gender						
Male						
Female						
Transgender ⁷						
Unknown						
Race and Hispanic Origin⁸						
Hispanic or Latino						
Not Hispanic or Latino	American Indian or Alaska Native alone					
	Asian alone					
	Black or African American alone					
	Native Hawaiian or Pacific Islander alone					
	White alone					
	Two or more races					
Unknown						

Population Groups					
MSM/IDU					
MSM					
PWID					
Heterosexual males					
Heterosexual females					
Other					
Unknown					
Total					

¹ Includes persons who received one or more services at the health care facility during the reporting period.

² Includes tests that are supported in any way by P520-2010-funded resources (e.g., funding, test kits, personnel, training and enhancements).

³ Includes persons who had a positive or negative test result.

⁴ Includes newly and previously diagnosed infections and those with unknown prior history

⁵ Includes newly identified HIV diagnosis that have been confirmed in surveillance. For persons included in this column, the HIV found and 2) there is no indication of a previous diagnosis by either client self-report (if the client was asked) or review of other records.

⁶ Includes self-report and provider report of new HIV diagnosis.

⁷ Transgender includes all persons whose gender identity or expression is different from their sex assigned at birth. Transgender woman, transgender male or transgender man, or other gender non-binary person.

⁸ Race and ethnicity are to be collected at the local level in accordance with OMB standards (<https://www.govinfo.gov/content/pkg/FR-2017-06-16/pdf/2017-06-16.pdf>). Aggregate using these categories. Hispanic or Latino persons can be of any race.

CDC Calculations for Percentages: *Percentage of persons tested (Column B%)*=(B#/A#)*100; *Percentage of HIV-Positive Persons (Column D%)*=(D#/B#)*100; *New Positivity—Confirmed in Surveillance (Column E%)*=(E#/B#)*100; *New Positivity—Self Report (Column F%)*=(F#/B#)*100

D Clinic

ons tested for and diagnosed with HIV (new and previous positives) in health care facilities identified as a priority

[illegible]

technical assistance, laboratory support, electronic medical record

✓ surveillance system is checked and 1) no prior report of HIV diagnosis is
or data sources (if other data sources were checked).

er persons may self-identify as transgender female or transgender

:/pkg/FR-1997-10-30/pdf/97-28653.pdf) and reported to CDC in the

ons (Column C%)=(C#/B#) * 100; *Percentage of Previous HIV-Positives*
rt (Column F%)=(F#/B#) * 100.

for routine opt-out screening.

Provision of PrEP Services: STD Clinics

Aggregate Data Tables for Prevent Indicator 3A1.3: Number and percentage of HIV-negative persons who are eligible for PrEP at the STD clinic

The following data table should be completed for each reporting STD clinic

Gray = auto-calculated cell; does not represent a reporting burden

Table XX. PrEP in STD Clinics

		HIV-Negative Persons ¹	Added for STD clinics: Already using PrEP	Screened for PrEP Need ^{2,3}		Eligible for PrEP Referral ^{4,5}		Referred Provider ⁶
		N	N	N	%	N	%	N
Age Group								
<15 Years								
15-19 Years								
20-29 Years								
30-65 Years								
≥ 66 Years								
Unknown								
Test Setting								
STD Clinic								
Other Health Care Setting								
Non-Health Care Setting								
Unknown								
Gender								
Male								
Female								
Transgender ¹²								
Unknown								
Race and Hispanic Origin¹³								
Hispanic or Latino								
Not Hispanic or Latino	American Indian or Alaska Native alone							
	Asian alone							
	Black or African American alone							
	Native Hawaiian or Pacific Islander alone							
	White alone							
	Two or more races							
Unknown								
Population Groups								
MSM/IDU								
MSM								

PWID							
Heterosexual males							
Heterosexual females							
Other							
Unknown							
Total							

¹ Includes HIV-negative persons who received HIV testing or PrEP services (e.g., screening, referral, linkage, prescription) that were in any way Denominator for Column B

^{2,3} Screening for PrEP eligibility is the process of conducting an initial assessment regarding a person's eligibility (e.g., behavioral risk screening and determining whether a more thorough evaluation is warranted). A person is screened if an initial assessment of eligibility for PrEP service PrEP screening, see Centers for Disease Control and Prevention: US Public Health Service: Preexposure prophylaxis for the prevention of HIV Update: a clinical practice guideline. <https://www.cdc.gov/hiv/pdf/risk/prep/cdc-hiv-prep-guidelines-2017.pdf>³Denominator is Column A (number of persons reflected in Column B should also be reflected in Column A).

⁴ Eligible for PrEP Referral refers to a person's status with regard to whether or not he or she meets appropriate criteria for using PrEP; specifically, not at risk for HIV, as defined locally or by CDC guidelines for PrEP

⁵Denominator is Column B (number of persons screened for PrEP). Persons reported as eligible for PrEP referral are those who have sexual activity and should also be reflected in the counts reported for columns A-B.

⁶ Referral to a PrEP provider is a process involving the provision of information on who the providers are, what documents referred person should bring, and what to expect from the referral process.

⁷Denominator is Column C (number of persons eligible for PrEP). Persons who are referred to a PrEP provider should also be reflected in the counts reported for columns A-D.

⁸ Linkage to PrEP provider is the process through which a person at risk for becoming infected with HIV is helped to access a healthcare provider for management for PrEP. Linked to a PrEP provider refers to the outcome of the referral or linkage of a PrEP eligible person to a PrEP provider, at the first appointment and verified through reviews of medical records or other data systems or self-report by the client.

⁹ Denominator is Column D (number of persons referred). Persons who are linked to a PrEP provider should also be reflected in the counts reported for columns A-E.

¹⁰ Prescribed PrEP refers to a person who has been adequately evaluated and received a prescription for PrEP. Prescription information may be obtained from medical records, pharmacy data, or self-report by client.

¹¹ Denominator is Column E (number of persons linked to PrEP). Persons who are prescribed PrEP should also be reflected in the counts reported for columns A-F.

¹² 'Transgender' includes all persons whose gender identity or expression is different from their sex assigned at birth. Transgender persons may be transgender woman, transgender male or transgender man, or other gender non-binary person.

¹³Race and ethnicity are to be collected at the local level in accordance with OMB standards (<https://www.govinfo.gov/content/pkg/FR-2017-06-16/pdf/2017-06-16.pdf>) and reported to CDC in the aggregate using these categories. Hispanic or Latino persons can be of any race.

CDC Calculations for Percentages: Percentage of Persons Screened for PrEP (Column B)=(B#/A#) * 100; Percentage of Persons Eligible for PrEP (Column C)=(C#/A#) * 100; Percentage of Persons Referred for PrEP (Column D)=(D#/C#) * 100; Percentage of Persons Linked to a PrEP Provider (Column E)=(E#/D#) * 100; Percentage of Persons Prescribed PrEP (Column F)=(F#/E#) * 100.

are linked to a PrEP provider and

Linked to PrEP Provider ^{6,7}	Linked to a PrEP Provider ^{8,9}		Added for STD Clinics: Prescribed PrEP at/by the STD clinic	
%	N	%	N	%

y supported by PS20-2010 funds.

ig) for pre-exposure prophylaxis (PrEP)
 es is completed. For further discussion on
 infection in the United States—2017
 umber of HIV-negative persons). All

fically, whether or not he or she is HIV-

r drug risk behaviors. These individuals

hould take with them, how to get to the

counts reported for columns A-C.

ider who offers evaluation and
 as indicated by the person's attendance

ported for columns A-D.

come from clinical records, pharmacy

rted for columns A-E.

ay self-identify as transgender female or

R-1997-10-30/pdf/97-28653.pdf) and

le for PrEP (Column C)=(C#/B#))* 100;
#/D#))* 100; Percentage of Persons

STD Clinic Service Capacity

Data Table for STD clinic capacity performance measures STD-3, STD-4, STD-5, and STD-6: Number of [X categ
Gray = auto-calculated cell; does not represent a reporting burden

Clinic service capacity domains
Prevention and referrals
Risk assessment and education for pre-exposure prophylaxis (PrEP) for HIV prevention
Referral to PrEP providers
Linkage to PrEP providers
Risk assessment and education for non-occupational post-exposure prophylaxis (nPEP)
Referral or linkage to nPEP providers
Referral or linkage to HIV care for people with HIV, if indicated
Clinic capacity PM-STD-3: # of prevention and referral services offered routinely by STD clinics
Laboratory capacity
At time of patient visit: Test for HIV (rapid or laboratory)
Through clinical laboratory: Extragenital (pharynx and rectum) NAAT for gonorrhea and chlamydia
Through clinical laboratory: Fourth generation antigen/antibody HIV test
Through clinical laboratory: Laboratory tests needed for providing nPEP and PrEP, as per clinical protocol
Viral Load testing
Clinic capacity PM-STD-4: # laboratory services offered routinely by STD clinics
Treatment
On site provision (starter pack or 3 month supply): PrEP
On site provision (7 day or full course): nPEP
On site: treatment for acute or new diagnosis of HIV care
By prescription: PrEP
By prescription: nPEP
By prescription: treatment for acute or new diagnosis of HIV care
Clinic capacity PM-STD-5: # of treatment services offered routinely by STD clinics
Service models
Ongoing care for patients taking PrEP
STD express visit
Telemedicine
Provision or linkage to kits for self collection for GC/CT testing outside the clinic
Provision or linkage to self-testing kits for HIV testing outside the clinic

Clinic capacity PM-STD-6: # of service models offered routinely by STD clinics

Optional: Any other information about clinic capacity you wish to provide at this time.

ory] services provided by STD clinics

Did your clinic provide this service? Y/N (in the reporting period)	Brief description of major service changes for this particular reporting period (if applicable)
