

# Pesticide Use on Sugarbeets in Minnesota 2021

5/2/2024

**For information regarding this report, please contact**

Naworaj Acharya ([naworaj.acharya@state.mn.us](mailto:naworaj.acharya@state.mn.us)), 651-201-6029

Denton Bruening ([denton.bruening@state.mn.us](mailto:denton.bruening@state.mn.us)), 651-261-1993

Minnesota Department of Agriculture

Pesticide and Fertilizer Management Division

# Table of Contents

|                                                         |          |
|---------------------------------------------------------|----------|
| <b>Table of Contents .....</b>                          | <b>3</b> |
| List of Tables .....                                    | 4        |
| List of Figures .....                                   | 4        |
| Acknowledgements .....                                  | 5        |
| Executive Summary .....                                 | 6        |
| Background.....                                         | 8        |
| Survey Objective, Data Collection and Limitations ..... | 10       |
| Data Interpretation and Findings .....                  | 11       |
| Pesticide Application Overview .....                    | 11       |
| Herbicide Application.....                              | 14       |
| Fungicide Application.....                              | 15       |
| Fungicide Seed Treatment .....                          | 16       |
| Insecticide Application .....                           | 17       |
| Insecticide Seed Treatment .....                        | 18       |
| Potential Implications.....                             | 19       |

## List of Tables

|                                                                                                  |    |
|--------------------------------------------------------------------------------------------------|----|
| Table 1. Herbicide active ingredients used on sugarbeet acres in 2021 .....                      | 11 |
| Table 2. Fungicide active ingredients used on sugarbeet acres in 2021 .....                      | 12 |
| Table 3. Fungicide active ingredients used for seed treatment on sugarbeet acres in 2021 .....   | 12 |
| Table 4. Insecticide active ingredients used on sugarbeet acres in 2021 .....                    | 13 |
| Table 5. Insecticide active ingredients used for seed treatment on sugarbeet acres in 2021 ..... | 13 |
| Table 6. Herbicide application details by active ingredient (a.i.) for sugarbeets .....          | 14 |
| Table 7. Fungicide application details by active ingredient (a.i.) for sugarbeets .....          | 15 |
| Table 8. Fungicide seed treatment details by active ingredient (a.i.) for sugarbeets .....       | 16 |
| Table 9. Insecticide application details by active ingredient (a.i.) for sugarbeets .....        | 17 |
| Table 10. Insecticide seed treatment details by active ingredients (a.i.) for sugarbeets .....   | 18 |

## List of Figures

|                                                                          |   |
|--------------------------------------------------------------------------|---|
| Figure 1. Sugarbeet production in Minnesota from 1970 through 2022 ..... | 8 |
| Figure 2. A map showing sugarbeet production areas in Minnesota .....    | 9 |

## Acknowledgements

---

This report is a result of collaborative efforts between the Minnesota Department of Agriculture (MDA) and Minnesota Sugarbeet Cooperatives. The MDA is very grateful to Southern Minnesota Beet Sugar Cooperative (SMBSC), American Crystal Sugar Company (ACSC) and Minn-Dak Farmers Cooperative (MDFC) for voluntarily providing the sugarbeet pesticide use information. Special thanks go to Joe Hastings (ACSC), Emma Burt (MDFC) and Mark Bloomquist (SMBSC) for coordinating the data compilation and reviewing the report. The MDA is also thankful to Dr. Tom Peters, Associate Professor/Sugarbeet Agronomist, North Dakota State University for providing the avenue to connect with the Minnesota Sugarbeet Cooperatives and reviewing the report. We are also grateful to the sugarbeet growers associated with SMBSC, ACSC and MDFC who were directly or indirectly involved in the data compilation.

## Executive Summary

---

Pesticides are an important tool in managing weeds (herbicides), diseases (fungicides), and insects (insecticides) in sugarbeets. Pesticide use information for corn, soybeans and other important Minnesotan crops has been well documented through the Minnesota Department of Agriculture (MDA) and United States Department of Agriculture's National Agricultural Statistics Service (USDA NASS) collaborative surveys. However, such information has not been available for sugarbeets through the MDA and USDA NASS survey. This survey was designed to collect basic pesticide use information on sugarbeet acres, including the use of pesticide treated seeds, for 2021 growing season. Southern Minnesota Beet Sugar Cooperative (SMBSC), American Crystal Sugar Company (ACSC), and Minn-Dak Farmers Cooperative (MDFC) collaborated for data compilation and provided the MDA a comprehensive pesticide use data set representative of their growers. The data were organized and presented in a form similar to pesticide data collected by the MDA and USDA NASS for other crops in Minnesota. This report represents 425,868 acres of sugarbeets planted in Minnesota by the growers associated with SMBSC, ACSC and MDFC.

This report summarizes herbicide, insecticide and fungicide use<sup>1</sup> on sugarbeet acres in Minnesota for the 2021 growing season. The pesticide use information presented in this report is solely based on the data provided by SMBSC, ACSC and MDFC and does not represent the sugarbeet acres not associated with the cooperatives.

- Sugarbeet growers associated with SMBSC, ACSC and MDFC reported<sup>2</sup> using 12 herbicide active ingredients, 5 insecticide active ingredients and 14 fungicide active ingredients to manage weeds, insects, and diseases excluding seed treatments. Growers used 10 fungicide active ingredients and 2 insecticide active ingredients as seed treatments.
- A total of 4,665,546 pounds of pesticide active ingredients including seed treatments were used on 425,868 acres of reported sugarbeet acres.
- Herbicides were applied to 100% of reported sugarbeet acres. Glyphosate (100%) was applied on the most acres.
- Fungicides were applied to 100% of reported sugarbeet acres. Triphenyltin Hydroxide (97%) was applied on the most acres. This does not include fungicide seed treatments.
- Fungicide treated seeds were used on 100% of reported acres. Thiram (100%) was applied on the most acres as seed treatments.

---

<sup>1</sup> Insecticide and fungicide seed treatments was included in this report. There was no herbicide included on the sugarbeet seed.

<sup>2</sup> Active ingredients used on very few acres were not reported, generally less than 1,000 acres.


- At least 47% of reported acres didn't receive any insecticide applications. Chlorpyrifos<sup>3</sup> (24%) was applied on the most acres. Neonicotinoid insecticide, imidacloprid, was applied to 4% of reported acres as an in-furrow at-plant application.
- About 70% of reported acres didn't receive any insecticide seed treatments. Insecticide seed treatments included only neonicotinoids, clothianidin and thiamethoxam. Clothianidin (29%) was applied on the most acres as seed treatments.

---

<sup>3</sup> Following the 2023 Eighth Circuit Court ruling, the Environmental Protection Agency (EPA) reinstated food and feed tolerances for chlorpyrifos. The MDA has conditionally registered chlorpyrifos products for food and feed uses for 2024 only, requiring [Water Quality Best Management Practices for chlorpyrifos](#) to be accompanied with the sale of each product.

## Background

In 2021, sugarbeet crop was the third most valuable crop in Minnesota after corn and soybeans with a value of 656 million dollars, according to the USDA<sup>4</sup>. This was the highest value of sugarbeet production recorded until this time<sup>4</sup>. In the past 10 years, Minnesota has typically produced between 30 to 35% of all US sugarbeet production (Figure 1)<sup>4</sup>. In each year since 1970, Minnesota was the top sugarbeet producing state. Sugarbeet production in the state is concentrated along the Minnesota River Valley and the Red River Valley (Figure 2)<sup>5</sup>.



**Figure 1. Sugarbeet production in Minnesota from 1970 through 2022 (Source: USDA)**

There are three sugarbeet cooperatives in Minnesota: Southern Minnesota Beet Sugar Cooperative (SMBSC), American Crystal Sugar Company (ACSC), and Minn-Dak Farmers Cooperative (MDFC). Each cooperative is owned by growers who are the shareholders of the company with the right and obligation to grow and deliver sugarbeets to be refined into white granulated sugar. Cooperatives provide technical assistance and supply resources to their growers for the proper management of weeds, diseases, and insects. A total of 425,868 acres of sugarbeets were planted in Minnesota in the 2021 growing season by the growers associated with these cooperatives. These planted acres produced 12.2 million tons of sugarbeets resulting in 28.7 tons per acre<sup>6</sup>.


<sup>4</sup> [https://www.nass.usda.gov/Data\\_and\\_Statistics/index.php](https://www.nass.usda.gov/Data_and_Statistics/index.php)

<sup>5</sup> <https://croplandcros.scinet.usda.gov/>

<sup>6</sup> Based on reported planted acres, not harvested acres

Several weeds, insects and diseases are commonly reported on sugarbeet acres in Minnesota<sup>7</sup>. Example grass and broadleaf weeds include waterhemp, pigweed, kochia, lambsquarter, wild oat, cocklebur, ragweed, thistle, barnyard grass, buffalo bur, foxtail, and nightshade. Sugarbeet root maggot, wireworms, cutworms, springtails and lygus bugs are major insect pests that attack sugarbeets. Diseases such as aphanomyces damping-off, rhizoctonia damping-off, fusarium yellows/yellowing, rhizomania (crazy root), cercospora leaf spot, bacterial leaf spot, and alternaria leaf spot are commonly reported on sugarbeets.

Sugarbeet farming requires intensive management practices that include proper pesticide use for managing a range of pests. Pesticides are an important tool to protect sugarbeets from weeds, diseases, and insects. Pesticide use information for other major Minnesotan crops has been well documented through the Minnesota Department of Agriculture (MDA) and United States Department of Agriculture's National Agricultural Statistics Service (USDA NASS) collaborative surveys. However, such information has not been available for sugarbeets through the MDA and USDA NASS survey. The MDA has been gathering and providing pesticide information for corn, soybeans, wheat, and alfalfa to the public for nearly 20 years in partnership with the USDA NASS (reports are available at [Agricultural Pesticide Sales and Use Reports - Statewide | Minnesota Department of Agriculture](#)).



**Figure 2. A map showing sugarbeet production areas in Minnesota (Source: USDA)**

---

<sup>7</sup> Sugarbeet production guide (<https://www.ndsu.edu/agriculture/sites/default/files/2023-01/a1698-23.pdf>)

## Survey Objective, Data Collection and Limitations

---

This survey was designed to collect basic information on herbicide, fungicide and insecticide use including the use of fungicide and insecticide treated seeds in sugarbeet industry for 2021 growing season. The survey is a result of collaborative efforts between the MDA and Minnesota sugarbeet cooperatives and is the first statewide report of pesticide use on sugarbeet acres in Minnesota from the MDA. The information presented in this report is based on reported planted acres, not harvested acres.

SMBSC, ACSC and MDFC collaborated to provide combined data set in the provided formats. The data represent collective and representative information of sugarbeet growers associated with SMBSC, ACSC and MDFC. The collected data were organized and presented in a form similar to pesticide data collected by the MDA and USDA NASS for other major crops in Minnesota. The provided data were presented as such without weighing them unlike the way the pesticide use data were presented in the MDA's other pesticide use reports (<https://www.mda.state.mn.us/agricultural-pesticide-sales-use-reports-statewide>).

It is not possible to determine which sugarbeet acres received two or more products of different active ingredients though the individual applications and rates were captured based on the provided information. For example, sugarbeet acres that received s-metolachlor applications were captured but it was not possible to report acres that received both s-metolachlor and ethofumesate. Sugarbeet acres treated with an herbicide that contains more than one active ingredient were presented as such without separating them by individual active ingredient.

The pesticide use information presented in this report is solely based on the data provided by SMBSC, ACSC and MDFC and does not represent the sugarbeet acres not associated with the cooperatives.

# Data Interpretation and Findings

---

## Pesticide Application Overview

Growers associated with SMBSC, ACSC and MDFC planted 425,868 acres of sugarbeets in the 2021 growing season. Growers reported 31 pesticide active ingredients excluding seed treatments (12 herbicide active ingredients, five insecticide active ingredients and 14 fungicide active ingredients) to manage weeds, insects, and diseases (Table 1 - 5). Growers reported 10 fungicide active ingredients and two insecticide active ingredients for seed treatments (Table 8 and 10).

**Table 1. Herbicide active ingredients used on sugarbeet acres in 2021**

| Active Ingredient<br>(Herbicide)          | Sites of<br>action group | Example<br>products* |
|-------------------------------------------|--------------------------|----------------------|
| Acetochlor                                | 15                       | Warrant              |
| Aciflourfen                               | 14                       | Ultra-Blazer         |
| Clethodim                                 | 1                        | Select               |
| Clopyralid                                | 4                        | Stinger              |
| Desmedipham + Phenmedipham                | 5 + 5                    | Betamix              |
| Desmedipham + Phenmedipham + Ethofumesate | 5 + 5 + 16               | Progress             |
| Dimethenamid-p                            | 15                       | Outlook              |
| Ethofumesate                              | 16                       | Nortron              |
| Glyphosate                                | 9                        | Glyphosate           |
| Quizalofop                                | 1                        | Assure II            |
| S-metolachlor                             | 15                       | Dual Magnum          |
| Triflurosulfuron                          | 2                        | Upbeet               |

\*Reference to commercial products or trade names is made with the understanding that no discrimination is intended, and no endorsement is implied. The example pesticide products mentioned in the table do not necessarily represent the products used by sugarbeet growers associated with SMBSC, ACSC and MDFC.

**Table 2. Fungicide active ingredients used on sugarbeet acres in 2021**

| Active Ingredient (Fungicide) | Modes of action group | Example products* |
|-------------------------------|-----------------------|-------------------|
| Azoxystrobin                  | 11                    | Quadris           |
| Bixafen                       | 7                     | Lucento           |
| Copper                        | M1                    | Copper            |
| Difenoconazole                | 3                     | Inspire           |
| Flutriafol                    | 3                     | Lucento           |
| Fluxapyroxad                  | 7                     | Priaxor           |
| Mancozeb                      | M3                    | Maneb             |
| Mefentrifluconazole           | 3                     | Provysol          |
| Propiconazole                 | 3                     | Inspire           |
| Prothioconazole               | 3                     | Proline           |
| Pyraclostrobin                | 11                    | Headline          |
| Tetraconazole                 | 3                     | Eminent           |
| Thiophanate Methyl            | 1                     | Topsin M          |
| Triphenyltin Hydroxide        | 30                    | Tin               |

\*Reference to commercial products or trade names is made with the understanding that no discrimination is intended, and no endorsement is implied. The example pesticide products mentioned in the table do not necessarily represent the products used by sugarbeet growers associated with SMBSC, ACSC and MDFC.

**Table 3. Fungicide active ingredients used for seed treatment on sugarbeet acres in 2021**

| Active Ingredient (Fungicide Seed Treatment) | Modes of action group | Example products* |
|----------------------------------------------|-----------------------|-------------------|
| Fludioxonil                                  | 12                    | Maxim 4           |
| Fluxapyroxad                                 | 7                     | Systiva           |
| Hymexazol                                    | 32                    | Tachigaren        |
| Mefenoxam                                    | 4                     | Apron XL          |
| Metalaxyl                                    | 4                     | Allegiance        |
| Metconazole                                  | 3                     | Metlock           |
| Penthiopyrad                                 | 7                     | Kabina            |
| Sedaxane                                     | 7                     | Vibrance          |
| Thiram                                       | M3                    | Thiram            |
| Tolclofos-methyl                             | 14                    | Rizolex           |

\*Reference to commercial products or trade names is made with the understanding that no discrimination is intended, and no endorsement is implied. The example pesticide products mentioned in the table do not necessarily represent the products used by sugarbeet growers associated with SMBSC, ACSC and MDFC.

**Table 4. Insecticide active ingredients used on sugarbeet acres in 2021**

| Active Ingredient<br>(Insecticide) | Mode of<br>action group | Example<br>products* |
|------------------------------------|-------------------------|----------------------|
| Chlorpyrifos                       | 1B                      | Lorsban              |
| Esfenvalerate                      | 3A                      | Asana XL             |
| Imidacloprid                       | 4A                      | Midac FC             |
| Terbufos                           | 1B                      | Counter              |
| Zeta-cypermethrin                  | 3A                      | Mustang max          |

\*Reference to commercial products or trade names is made with the understanding that no discrimination is intended, and no endorsement is implied. The example pesticide products mentioned in the table do not necessarily represent the products used by sugarbeet growers associated with SMBSC, ACSC and MDFC.

**Table 5. Insecticide active ingredients used for seed treatment on sugarbeet acres in 2021**

| Active Ingredient<br>(Insecticide Seed Treatment) | Mode of<br>action group | Example<br>products* |
|---------------------------------------------------|-------------------------|----------------------|
| Clothianidin                                      | 4A                      | Poncho Beta          |
| Thiamethoxam                                      | 4A                      | Cruiser              |

\*Reference to commercial products or trade names is made with the understanding that no discrimination is intended, and no endorsement is implied. The example pesticide products mentioned in the table do not necessarily represent the products used by sugarbeet growers associated with SMBSC, ACSC and MDFC.

## Herbicide Application

A summary of herbicide applications on reported sugarbeet acres is presented in Table 6. Herbicides were applied to 100% of reported sugarbeet acres in 2021 growing season<sup>8</sup>. Glyphosate (100%) was applied on the most acres. Glyphosate and s-metolachlor contributed the most pounds of active ingredient applied per year for reported planted acres.

**Table 6. Herbicide application details by active ingredient (a.i.) for sugarbeets**

| Agricultural Chemical (a.i.)              | Planted Acres Treated Percent | Average Applications Number | Average Rate Per Application Pounds per Acre (a.i.) | Average Rate Per Crop Year Pounds per Acre (a.i.) | Total Applied Per Crop Year Total Pounds (a.i.) |
|-------------------------------------------|-------------------------------|-----------------------------|-----------------------------------------------------|---------------------------------------------------|-------------------------------------------------|
| Acetochlor                                | 6.46                          | 1.00                        | 0.94                                                | 0.94                                              | 25,805                                          |
| Aciflourfen                               | 8.68                          | 1.00                        | 0.25                                                | 0.25                                              | 9,243                                           |
| Clethodim                                 | 11.29                         | 1.00                        | 0.07                                                | 0.07                                              | 3,280                                           |
| Clopyralid                                | 51.95                         | 1.53                        | 0.07                                                | 0.11                                              | 23,801                                          |
| Desmedipham + Phenmedipham                | 3.82                          | 1.00                        | 0.24                                                | 0.24                                              | 3,970                                           |
| Desmedipham + Phenmedipham + Ethofumesate | 0.14                          | 1.00                        | 0.34                                                | 0.34                                              | 202                                             |
| Dimethenamid-p                            | 24.57                         | 1.19                        | 0.56                                                | 0.67                                              | 70,047                                          |
| Ethofumesate Post-emergence*              | 45.34                         | 1.68                        | 0.13                                                | 0.21                                              | 40,549                                          |
| Ethofumesate PPI and Pre-emergence**      | 31.21                         | 1.00                        | 1.65                                                | 1.65                                              | 219,285                                         |
| Glyphosate                                | 100.00                        | 2.08                        | 0.98                                                | 2.05                                              | 871,966                                         |
| Quizalofop                                | 2.49                          | 1.00                        | 0.05                                                | 0.05                                              | 511                                             |
| S-metolachlor Post-emergence *            | 46.34                         | 1.30                        | 0.96                                                | 1.24                                              | 245,009                                         |
| S-metolachlor PPI and Pre-emergence **    | 25.26                         | 1.00                        | 0.72                                                | 0.72                                              | 77,064                                          |
| Triflusalufuron                           | 0.24                          | 1.00                        | 0.03                                                | 0.03                                              | 31                                              |

\*Post includes herbicides that are applied post-emergence.

\*\*PPI (Preplant Incorporated) and pre-emergence includes herbicides that are applied prior to planting and that are incorporated into the soil or applied pre-emergence.

<sup>8</sup> There was no herbicide seed treatment on sugarbeets.

## Fungicide Application

A summary of fungicide applications on sugarbeet acres is presented in Table 7. Fungicides were applied to 100%<sup>9</sup> of reported sugarbeet acres in 2021 growing season excluding seed treatments. Triphenyltin hydroxide (97%) was applied on the most acres. Mancozeb and triphenyltin hydroxide contributed the most pounds of active ingredient applied per year for reported acres excluding seed treatments.

**Table 7. Fungicide application details by active ingredient (a.i.) for sugarbeets**

| Agricultural Chemical (a.i.) | Planted Acres Treated Percent | Average Applications Number | Average Rate Per Application Pounds per Acre (a.i.) | Average Rate Per Crop Year Pounds per Acre (a.i.) | Total Applied Per Crop Year Total Pounds (a.i.) |
|------------------------------|-------------------------------|-----------------------------|-----------------------------------------------------|---------------------------------------------------|-------------------------------------------------|
| Azoxystrobin at Plant*       | 39.28                         | 1.00                        | 0.15                                                | 0.15                                              | 25,875                                          |
| Azoxystrobin Post**          | 13.99                         | 1.00                        | 0.24                                                | 0.24                                              | 14,522                                          |
| Bixafen                      | 3.86                          | 1.00                        | 0.07                                                | 0.07                                              | 1,088                                           |
| Copper                       | 25.50                         | 1.12                        | 0.57                                                | 0.64                                              | 69,017                                          |
| Difconazole                  | 36.50                         | 1.00                        | 0.11                                                | 0.11                                              | 17,684                                          |
| Flutriafol                   | 3.93                          | 1.00                        | 0.11                                                | 0.11                                              | 1,882                                           |
| Fluxapyroxad                 | 4.78                          | 1.00                        | 0.07                                                | 0.07                                              | 1,481                                           |
| Mancozeb                     | 92.98                         | 3.30                        | 1.60                                                | 5.28                                              | 2,090,654                                       |
| Mefentrifluconazole          | 37.71                         | 1.01                        | 0.10                                                | 0.11                                              | 16,932                                          |
| Propiconazole                | 36.50                         | 1.00                        | 0.11                                                | 0.11                                              | 17,684                                          |
| Prothioconazole              | 83.47                         | 1.00                        | 0.18                                                | 0.18                                              | 63,315                                          |
| Pyraclostrobin at Plant*     | 0.17                          | 1.00                        | 0.15                                                | 0.15                                              | 105                                             |
| Pyraclostrobin Post**        | 46.74                         | 1.00                        | 0.15                                                | 0.15                                              | 29,097                                          |
| Tetraconazole                | 8.03                          | 1.00                        | 0.18                                                | 0.18                                              | 6,075                                           |
| Thiophanate Methyl           | 31.58                         | 1.00                        | 0.35                                                | 0.35                                              | 47,281                                          |
| Triphenyltin Hydroxide       | 97.16                         | 2.00                        | 0.25                                                | 0.52                                              | 216,866                                         |

\*At Plant includes fungicides applied at planting.

\*\*Post includes fungicides applied after planting.

<sup>9</sup> Verified by the Coops

## Fungicide Seed Treatment

A summary of fungicide seed treatments on reported sugarbeets acres is presented in Table 8. Fungicide treated seeds were used on 100% of reported acres. A total of 10 different fungicide active ingredients were reported for seed treatments. Thiram (100%) was applied on the most acres as seed treatments. Hymexazol and penthiopyrad contributed the most pounds of active ingredient applied per year for reported acre as seed treatments.

**Table 8. Fungicide seed treatment details by active ingredient (a.i.) for sugarbeets**

| Agricultural Chemical (a.i.) | Planted Acres Treated Percent | Average Rate Per Application Pounds per Acre (a.i.) | Average Rate Per Crop Year Pounds per Acre (a.i.) | Total Applied Per Crop Year Total Pounds (a.i.) |
|------------------------------|-------------------------------|-----------------------------------------------------|---------------------------------------------------|-------------------------------------------------|
| Fludioxonil                  | 0.10                          | 0.0001                                              | 0.0001                                            | <1                                              |
| Fluxapyroxad                 | 17.98                         | 0.0067                                              | 0.0067                                            | 514                                             |
| Hymexazol                    | 91.41                         | 0.0320                                              | 0.0300                                            | 12,199                                          |
| Mefenoxam                    | 1.19                          | 0.0002                                              | 0.0002                                            | <1                                              |
| Metalaxyl                    | 98.81                         | 0.0002                                              | 0.0002                                            | 86                                              |
| Metconazole                  | 3.02                          | 0.0003                                              | 0.0003                                            | 4                                               |
| Penthiopyrad                 | 59.55                         | 0.0188                                              | 0.0188                                            | 4,774                                           |
| Sedaxane                     | 33.43                         | 0.0020                                              | 0.0020                                            | 287                                             |
| Thiram                       | 99.90                         | 0.0034                                              | 0.0034                                            | 1,427                                           |
| Tolclofos-methyl             | 3.02                          | 0.0007                                              | 0.0007                                            | 9                                               |

# Insecticide Application

A summary of insecticide applications on reported sugarbeet acres is presented in Table 9. Chlorpyrifos<sup>10</sup> (24%) was applied on the most acres and also contributed the most pounds of active ingredient applied per year for reported acres excluding seed treatments.

**Table 9. Insecticide application details by active ingredient (a.i.) for sugarbeets**

| Agricultural Chemical (a.i.) | Planted Acres Treated Percent | Average Applications Number | Average Rate Per Application Pounds per Acre (a.i.) | Average Rate Per Crop Year Pounds per Acre (a.i.) | Total Applied Per Crop Year Total Pounds (a.i.) |
|------------------------------|-------------------------------|-----------------------------|-----------------------------------------------------|---------------------------------------------------|-------------------------------------------------|
| Chlorpyrifos at Plant*       | 0.33                          | 1.00                        | 1.50                                                | 1.50                                              | 2,115                                           |
| Chlorpyrifos Post**          | 23.63                         | 1.00                        | 1.00                                                | 1.00                                              | 100,649                                         |
| Esfenvalerate                | 6.27                          | 1.00                        | 0.05                                                | 0.05                                              | 1,323                                           |
| Imidacloprid                 | 4.50                          | 1.00                        | 0.18                                                | 0.18                                              | 3,458                                           |
| Terbufos                     | 10.18                         | 1.00                        | 1.50                                                | 1.50                                              | 65,042                                          |
| Zeta-cypermethrin            | 8.00                          | 1.00                        | 0.03                                                | 0.03                                              | 852                                             |

\*At Plant includes insecticides applied at planting.

\*\*Post includes insecticides applied after planting.

<sup>10</sup> Following the 2023 Eighth Circuit Court ruling, the Environmental Protection Agency (EPA) reinstated food and feed tolerances for chlorpyrifos. The MDA has conditionally registered chlorpyrifos products for food and feed uses for 2024 only, requiring [Water Quality Best Management Practices for chlorpyrifos](#) to be accompanied with the sale of each product.

## Insecticide Seed Treatment

A summary of insecticide seed treatments on reported sugarbeet acres is presented in Table 10. Only about 30% of reported acres received insecticide seed treatments. Neonicotinoids were the only insecticides reported for seed treatments. Clothianidin (29%) was applied on the most acres and also contributed the most pounds of active ingredient applied per year for reported acres as seed treatments.

**Table 10. Insecticide seed treatment details by active ingredients (a.i.) for sugarbeets**

| Agricultural Chemical (a.i.) | Planted Acres Treated <i>Percent</i> | Average Rate Per Application Pounds per Acre (a.i.) | Average Rate Per Crop Year Pounds per Acre (a.i.) | Total Applied Per Crop Year Total Pounds (a.i.) |
|------------------------------|--------------------------------------|-----------------------------------------------------|---------------------------------------------------|-------------------------------------------------|
| Clothianidin                 | 29.40                                | 0.09                                                | 0.09                                              | 8,232                                           |
| Thiamethoxam                 | 0.98                                 | 0.08                                                | 0.08                                              | 333                                             |

## Potential Implications

---

This survey is a result of collaborative efforts between the MDA and Minnesota sugarbeet cooperatives and is the first statewide report of pesticide use on sugarbeet acres from the MDA. In the past, similar information has been collected for corn, soybean, wheat, and other crops through the cooperative efforts between the MDA and USDA NASS.

The survey results will help the MDA understand current pesticide use practices in sugarbeets. It is important that the MDA, as a state regulatory agency, stays updated with pesticide use practices in crop industries and assists them to deal with potential regulatory changes.

The MDA is responsible for the development and promotion of pesticide best management practices (BMPs). The BMPs include both legal requirements and voluntary practices that are designed to prevent and minimize the degradation of Minnesota's water resources. This survey indicates that sugarbeet growers followed BMPs related to the rate and frequency of pesticide applications. It also showcases the good pesticide record keeping practices followed by sugarbeet farmers and cooperatives.

In 2021, sugarbeet growers used 1,590,762 pounds of herbicides, 2,892,780 pounds of fungicides and 182,004 pounds of insecticides including fungicide and insecticide seed treatments, which represent 34, 62 and 4 percent of the total pesticides used on sugarbeet acres. Sugarbeet growers applied 42 different pesticide active ingredients in varying quantities based on total pounds of active ingredient applied per crop year. The amount of pesticide used does not necessarily correlate with its environmental risks, which depend on many factors such as chemical behavior and toxicity of the pesticides, and environmental factors. For instance, glyphosate sticks to soil and does not move around much; while acetochlor, with low soil adsorption and high solubility, can be easily washed away by rain, depending on field conditions. In 2021, a total of 25,805 pounds of acetochlor and 871,966 pounds of glyphosate were used on sugarbeet acres. Glyphosate has consistently been sold in higher quantities than acetochlor in Minnesota since 2000. However, the MDA water monitoring results show multiple instances of the state water quality standard violations due to acetochlor, while no such violations were linked to glyphosate.